

This is a “preproof” accepted article for *Journal of Clinical and Translational Science*.

This version may be subject to change during the production process.

10.1017/cts.2024.512

Evidence-Based Recruitment Strategies for Clinical Research: Study Personnel’s and Research Participants’ Perceptions About Successful Methods of Outreach for a U.S. Autism-Research Cohort

Robin P. Goin-Kochel^{1,2}, Ivana Lozano^{1,2}, Gabrielle Duhon^{1,2}, Gabriela Marzano^{1,2}, Amy Daniels³, J. Kiely Law³, Katharine Diehl³, LeeAnne Green Snyder³, Pamela Feliciano³, Wendy K. Chung^{3,4}, & SPARK Consortium

¹Baylor College of Medicine, Houston, TX

²Texas Children’s Hospital, Houston, TX

³Simons Foundation, New York, NY

⁴Boston Children’s Hospital, Harvard Medical School, Boston, MA

Corresponding author: Robin P. Goin-Kochel, Ph.D., Autism Program, Meyer Center for Developmental Pediatrics and Autism, Texas Children’s Hospital, 8080 N. Stadium Drive, Suite 100, Houston, TX 77018; Telephone: 832-824-3390; Email: kochel@bcm.edu

Running head: EVIDENCE-BASED RECRUITMENT STRATEGIES

Conflicts of Interest: None

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Abstract

Introduction: Under enrollment of participants in clinical research is costly and delays study completion to impact public health. Given that research personnel make decisions about which strategies to pursue and participants are the recipients of these efforts, we surveyed research staff ($n = 52$) and participants ($n = 4,144$) affiliated with SPARK (Simons Foundation Powering Autism for Knowledge)—the largest study of autism in the U.S.—to understand their perceptions of effective recruitment strategies.

Methods: In Study 1, research personnel were asked to report recruitment strategies that they tried for SPARK and to indicate which ones they would and would not repeat/recommend. In Study 2, SPARK participants were asked to indicate all the ways they heard about the study prior to enrollment and which one was most influential in their decisions to enroll.

Results: Staff rated *speaking with a SPARK study-team member* (36.5%), *speaking with a medical provider* (19.2%), *word of mouth* (11.5%), and *a live TV news story* (11.5%) as the most successful strategies. Participants most often heard about SPARK via *social media* (47.0%), *speaking with a medical provider* (23.1%), and an *online search* (20.1%). Research personnel's and participants' views on effective recruitment strategies often differed, with the exception of *speaking with a medical provider*.

Conclusion: Results suggest that a combination of strategies is likely to be most effective in reaching diverse audiences. Findings have implications for the selection of strategies that meet a study's specific needs, as well as recruitment-strategy “combinations” that may enhance the influence of outreach efforts.

Keywords: research recruitment; participant enrollment; autism; parent/caregiver

Evidence-Based Recruitment Strategies for Clinical Research: Study Personnel's and Research Participants' Perceptions About Successful Methods of Outreach for a U.S. Autism-Research Cohort

Recruitment for clinical studies remains one of the biggest challenges to performing studies of all sizes. As many as 75% of studies are unable to enroll the proposed number of participants and 90% fail to enroll within the original timeframe[1]. Study preparation and start-up costs comprise the largest proportion of expenses in clinical research[2], and extensions granted to low-enrolling studies further increase costs and delay results that could have important public-health implications. Unfortunately, many studies are eventually abandoned as a result of failing to meet enrollment goals. A report from Oregon Health and Science University revealed that the university spent nearly \$1 million in fiscal year 2009 on studies that terminated early because of low enrollment numbers[2]. Not only do these investments fail to produce any scientific results[3], they also indirectly prevent other research studies from being conducted that could produce results. For these reasons, research dollars need to be allocated to recruitment activities that yield the greatest return on investment. Understanding which recruitment methods are most likely to successfully enroll participants is thus of significant importance financially, scientifically, and ethically.

Common Recruitment Challenges

Research teams experience a range of challenges that contribute to low enrollment in clinical studies. Factors like the topic of study, invasiveness or riskiness of procedures, duration of participation, and perceived cost/benefit ratio may all influence a potential participant's willingness to participate. For example, during a survey of individuals aged 14-21 with post-traumatic stress disorder who were participating in a clinical trial, approximately 60% of respondents reported the demands of the study to be a significant barrier to participation, including video recordings, problems with transportation, and diagnostic procedures[4]. In studies that rely heavily on referrals from healthcare providers, gatekeeping—or the reluctance to share research opportunities with potentially eligible patients—can hinder recruitment efforts and lead to selection bias[5]. Relatedly, ensuring racial and ethnic representativeness can be difficult, as research has identified several barriers to participation among communities of color. In particular, Black/African American individuals commonly cite concerns about racial discrimination and mistrust of healthcare/research staff as barriers to participation in genetic

research[6-9], while Asian families report language barriers and a lack of information/resources available in their native languages[8]. Fortunately, recent studies leveraging tools within the electronic health record (EHR) for research recruitment demonstrate both (a) improved collaboration between healthcare providers and researchers and (b) increased engagement and retention of families of color in clinical research[10-12]. Efforts like these are critical for identifying effective recruitment strategies that counteract barriers to research participation and expedite study progress.

Effective Recruitment Strategies

Several types of recruitment methods have been described as facilitating recruitment and retention efforts for clinical-research studies, with some of the most common being social media advertising, physician referrals, and study flyers[4,13]. In one study that compared the effectiveness of paid social-media advertising (i.e., Facebook) with a radio-based recruitment campaign, social media demonstrated superior outreach at a substantially lower cost[14]. Similar work that compared Facebook advertising with traditional recruitment methods (e.g., flyers, newspaper ads, radio, word-of-mouth) found that, while the former was more cost effective during initial outreach and screening efforts, it was less cost effective in terms of actual enrollment and study completion; as such, the authors recommended using a variety of recruitment strategies simultaneously[15]. Other strategies described as effective include sending personalized letters to potentially eligible participants, using telephone reminders throughout the study, and providing financial incentives[16-17]. Additionally, being flexible in terms of when study visits are offered (e.g., including evenings and weekends) and actively engaging clinic staff in the identification of potential participants were successful strategies for recruiting parents in behavioral-intervention trials[18]. Furthermore, targeted recruitment of particular racial or ethnic groups should include sociocultural-specific methods[16], such as recruiting through places of worship and having the study team reflect similar racial, ethnic, and language backgrounds of the participants. Collectively, this information provides an evidence-base for the range and relative success of various recruitment strategies. However, the majority of this work focuses on adult-patient populations with mental and medical illnesses who enrolled in treatment-intervention studies. Specific to recruitment in pediatric research, investigators from one study observed that *type* of recruitment method was key, with in-person recruitment being the only factor significantly associated with achieving 80% of enrollment goals[19]. These results were similar

to those of another study that demonstrated active (direct patient outreach) versus passive (flyers, posters, social media, press release) recruitment strategies as superior in engaging families for a pediatric clinical trial addressing obesity, with active strategies effectively recruiting 95% of the total sample[20]. Literature on this topic is sparse, however, and virtually nothing is known about strategies that are most effective for recruitment of pediatric populations with developmental disabilities, where caregivers often decide on behalf of their minor children and/or dependents whether to participate.

Given that (a) investigators and their research teams choose which recruitment strategies to pursue and (b) parents/caregivers are frequently the recipients of these outreach efforts for studies targeting children/dependents, it is valuable to examine these groups' perceptions about which recruitment methods are most successful in facilitating participant enrollment. Investing time and financial resources in strategies that resonate most with the target audience stands to maximize study enrollment while improving efficiency. Study teams can use this information to evaluate their current recruitment practices and determine whether and where changes may be needed. With this in mind, the purpose of the current study was to assess study personnel's and research participants' perspectives about effective recruitment strategies for the national SPARK cohort of individuals with autism and their family members.

SPARK Cohort

As described in *An evaluation of participant recruitment in SPARK, a large, online longitudinal research study of autism*[21], SPARK (Simons Foundation Powering Autism Research for Knowledge) is a national US study led by the Simons Foundation Autism Research Initiative to identify new genetic contributions for autism and accelerate the pace of autism research[22]. A key goal for SPARK is to sequence the DNA of 50,000 individuals with autism and both biological parents (trios). To reach this goal, SPARK enlisted the help of more than 30 clinical sites across the U.S. to recruit participants and assist them with study completion. The clinical site network operated under a single independent regulatory board (Western Institutional Review Board—Copernicus Group [WIRB-CG]). As such, all sites had IRB approval to conduct the same research activities; however, site-specific institutional barriers may have prevented some sites from pursuing particular activities. Each recruitment site was granted between \$150,000 to \$200,000 annually (inclusive of 20% indirect costs) to enroll a certain number of families each year. The budget was allocated at the PIs' discretion for personnel, supplies, and

recruitment/outreach activities. The current assessment specifically focused on participant recruitment at the site level to afford a granular view of these activities relative to the broader outreach efforts of the central SPARK team at the Simons Foundation. For a complete overview of these SPARK central efforts, please see Daniels et al.[21] Consistent with this paper, we refer to the “primary account holder” as the individual in the family who initiated participation in SPARK and “clinical-site affiliation” as the primary account holder’s self-selected association with a clinical site in the SPARK network. Both studies described below were reviewed and approved by the Baylor College of Medicine Internal Review Board.

Studies: Methods and Results

Study 1: Study Teams’ Perceptions of Effective Recruitment Strategies

Participants. We invited 150 current or past coordinators and principal investigators (PIs) from 25 SPARK sites (active as of June 2018) to participate in a survey about the recruitment strategies that their site had used and which they felt were most successful in influencing families to enroll. A total of 52 individuals responded from 23 sites, for a response rate of 35%; 32 (61.5%) were in a coordinator role, 13 (25.0%) were in a PI role, 5 (9.6%) were in some other role, and 2 (3.8%) were no longer working on the project. In terms of their time/effort allocated to SPARK, 17 (32.7%) contributed 75% effort or more, 10 (19.2%) contributed between 25%—74% effort, 6 (11.5%) contributed between 11%—24% effort, and 19 (36.5%) contributed $\leq 10\%$ effort. Before joining SPARK, 39 (75.0%) had prior experience recruiting for a clinical-research study.

Procedure. We developed a REDCap online questionnaire that asked about recruitment strategies that study teams had used for SPARK, which strategies they believed were most influential in participants’ decisions to enroll, perceived cost of each strategy, and which strategies they believed were most effective overall. An initial list of 20 specific recruitment strategies were presented in the questionnaire, with three open-ended spaces for participants to provide additional strategies not listed (see Table 1 for a complete list and description provided to participants). Email invitations to complete this questionnaire were distributed to potential participants, with up to three email reminders sent to non-responders. The survey was open for six days prior to a SPARK all-site investigator meeting, at which aggregate results from the survey were shared. Participants were incentivized with a chance to have their names drawn to receive one of five \$100 gift cards. Data were analyzed using SPSS v. 28 analytical software.

Results. The average number of recruitment strategies that study personnel indicated they had used for SPARK was 11.8 ($SD = 2.9$, range = 4–18). The frequencies and proportions of participants using each type of strategy is shown in Table 1. Other recruitment methods listed in open-ended format that did not fit into a category included advertising prior to previews at movie theaters, smaller email blasts (i.e., < 500), and attending non-autism-specific events (e.g., festivals, state fairs). The methods most frequently endorsed as being most influential in families' decisions to participate included *speaking with a SPARK team study-team member* ($n = 19$, 36.5%), *speaking with a medical provider* ($n = 10$, 19.2%), *word of mouth* ($n = 6$, 11.5%), and a *live TV news story* ($n = 6$, 11.5%).

In terms of perceived costs of recruitment strategies, participants rated most strategies as “least” or “somewhat” costly, with the highest proportions indicating (a) *unpaid social media* ($n = 44$, 97.8%), *mass email blasts* ($n = 39$, 97.5%), *hospital/clinic website* ($n = 38$, 97.4%), and *blog post* ($n = 12$, 91.7%) as least costly and (b) *transit advertising* ($n = 6$, 100%), *paid radio advertising* ($n = 12$, 63.1%), *on-site registration/saliva-collection events* ($n = 16$, 36.3%) and *paid website advertising* ($n = 3$, 30.0%) as most costly (Table 2). In terms of recruitment methods that study personnel had tried and believed were most effective overall, participants indicated the following strategies that they were likely to repeat/recommend or rate as one of the best: *speaking with SPARK team members in clinic* ($n = 37$, 86.0%), *on-site registration/saliva-collection events* ($n = 32$, 72.7%), *paid social media* ($n = 26$, 59.0%), *speaking with a medical provider in clinic* ($n = 25$, 52.0%), and *printed materials* ($n = 27$, 51.9%). The following were indicated as strategies that they would not repeat/recommend: a *blog post* ($n = 10$, 71.4%), *radio PSA* ($n = 6$, 66.7%), and a *radio or podcast interview* ($n = 11$, 52.4%) (Table 3). Study personnel most often believed that a family heard about SPARK between two and four times before enrolling ($n = 40$, 76.9%).

If a participant had not used a given recruitment strategy, we asked them to indicate the reasons why. Most reported a lack of knowledge about the strategy as a reason (23.0%), followed by lack of money (17.7%), lack of time (17.3%), lack of personnel (12.5%), lack of institutional support/within-site barriers (11.7%), lack of IRB approval (3.3%), or some other reason (14.5%). When asked how easy or difficult it was to recruit for SPARK compared to other studies, 12 (23.1%) said it was easier, 16 (30.8%) said it was harder, and 12 (23.1%) said it was about the same (12 [23.1%] had no response).

Study 2: SPARK Participants' Perceptions of Effective Recruitment Strategies

Participants. We invited primary account holders enrolled in SPARK (i.e., individuals who initiated participation on behalf of themselves and/or their families) and who had (a) consented to providing genetic samples and (b) were affiliated with one of the top 16 performing sites or had no particular site affiliation to participate in a survey about how they learned about SPARK and their understanding of participation. A total of 26,997 primary account holders received invitations about the study; 4,144 completed the survey, for a response rate of 15.3% (response rate range across sites = 12%—18%). Among these participants, 3,617 (87.3%) were parents/legally authorized representatives of minor children or adult dependents with autism and 527 (12.7%) were self-reporting, independent adults with autism; 3,627 (87.5%) reported as female; 2,516 (60.7%) were between the ages of 30-44 and 1,164 (28.1%) were between 45-60; 3,670 (88.6%) were White, 251 (6.1%) were Black/African American, 132 (3.2%) were Asian, 154 (3.7%) endorsed some other race; and 436 (10.5%) were Hispanic/Latinx. A total of 3,425 (82.6%) had provided a sample for the genetic component of SPARK (return of the saliva kit for the individual with ASD in the family), and 1,156 (27.9%) were not affiliated with one of the clinical sites.

At the time of enrollment, all SPARK participants were asked how they heard about the study. If a participant was referred by a clinical site and used a site-specific URL to join, the response to this question was set at “clinical site / hospital / university” from a range of responses in a drop-down menu. Therefore, caution should be taken when interpreting these results, as individuals who were referred by a clinical site could have heard about SPARK in several different ways. For participants of this study, responses to, “How did you hear about us?” were as follows: 2,577 (62.4%) clinical site/hospital/university; 938 (22.7%), online; 144 (3.5%), invited by a family member; 121 (2.9%), media announcement; 116 (2.8%), a friend; 88 (2.1%), the Interactive Autism Network (IAN); 68 (1.7%), healthcare provider; 51 (1.2%), a community-based organization; and 26 (0.6%), through some other method.

Procedure. We collaborated with the SPARK Research Match team to develop and distribute a questionnaire about how families learned about SPARK, methods they deemed most influential in their decisions to enroll, and their understanding of what participation entailed. Potential participants were invited in six batches from October 2, 2019, to November 6, 2019. The survey was open for two months (from October 2, 2019, to December 2, 2019), with up to

three reminder emails sent to non-responders. Participants were incentivized with a chance to have their names drawn to receive one of 25 portable power banks (i.e., phone/tablet charger), valued at \$40 each. Data were analyzed using SPSS v. 28 analytical software.

Results. The survey contained a list of 15 methods through which participants may have learned about SPARK, with instructions for them to select all the ways they heard about the study prior to enrolling. Table 4 shows results for participants who endorsed hearing about the study through each method, as well as the proportions who rated each method as most influential in their decisions to enroll in the study. *Social media* was the most commonly reported way people learned about the study ($n = 1,946$, 47.0%), followed by *speaking with a medical provider* ($n = 958$, 23.1%), an *online search* (which included the SPARK website and/or hospital/clinic website; $n = 833$, 20.1%), and *flyer/print material* ($n = 529$, 12.8%), while *newspaper* ($n = 20$), *radio* ($n = 38$), and *transit ads* ($n = 14$) were the least frequently endorsed ($\leq 1\%$ each). Most participants said that they heard about SPARK only one time ($n = 1,875$, 45.2%) or between two and four times ($n = 1,801$, 43.5%) prior to enrolling. Strategies that were significantly associated with return of the saliva kit for the individual with ASD included *speaking with a medical provider* (24.4% vs. 17.2%, $\chi^2[1] = 16.874$, $p < .001$), *speaking with a study-team member at the doctor's office* (8.7% vs. 5.1%, $\chi^2[1] = 9.967$, $p = .002$), and *speaking with a study-team member at a community event* (5.9% vs. 3.8%, $\chi^2[1] = 5.105$, $p = .024$); hearing about the study via *social media* was associated with a significantly lower rate of saliva-kit return (45.5% vs. 54.0%, $\chi^2[1] = 17.135$, $p < .001$).

Participants who endorsed learning about SPARK from a *medical provider* rated that method as most influential in their decisions to enroll (72.1%). Other methods commonly rated as most influential were *speaking with the study team at a community event* (68.4%), *social media* (63.0%), *friend or family member* (54.8%), and *speaking with a study-team member at the doctor's office* (54.8%). When a participant said that they had heard about SPARK through a given outlet but did not rate that outlet as most influential in their decision to enroll, the alternate strategies commonly endorsed as most influential were *social media*, an *online search*, *speaking with a medical provider*, and hearing about the study from a *friend or family member*. Additional results can be seen in Figure 1.

With respect to demographic influences on recruitment strategies, males more often than females rated hearing about the study from a *friend or family member* (12.1% vs. 7.4%) or via an

online search (17.1 % vs. 12.5%) as most influential in their decision to enroll, whereas females more often than males endorsed *social media* (31.5% vs. 20.8%) as most influential. Younger individuals rated hearing about the study via *social media* as most influential in their decision to enroll (42.4% of those 18-29 vs. 13.0% of those 60+), while older individuals endorsed an *online search* as most influential (16.9% of those 60+ vs. 12.3% of those 30-44). Both the youngest (age 18-29, 13.2%) and oldest (age 60+, 13.0%) individuals rated hearing about the study from a *friend or family member* as most influential in their decision to enroll compared to those in the two middle-aged groups (age 30-44, 7.2%; age 45-60, 7.6%). Differences were also observed across racial groups, as American Indian/Alaska Native participants (41.4%) and Native Hawaiian/Other Pacific Islander participants (42.9%) more often endorsed hearing about the study via *social media* as most influential in their decision to enroll, compared to Asian participants (15.2%), who least frequently indicated *social media* as most influential. Asian participants more often rated an *online search* (22.0%) and *speaking with a medical provider* (18.2%) as most influencing their decision to enroll, whereas Black/African American participants most often endorsed *social media* (30.3%), *speaking with a medical provider* (15.9%), and receiving a *personal email* from the study team (8.4%) as most influential. Hispanic/Latinx participants most often rated hearing about the study via *social media* (29.1%), their *medical provider* (17.4%), and an *online search* (15.1%) as most influential in their decision to enroll. Complete results can be seen in Tables 5 and 6.

Differences were also observed in how groups of primary account holders rated the influence of recruitment strategies. Compared to parents/legally authorized representatives, independent adults with autism were more likely to endorse *social media* (43.9% vs. 28.9%), hearing about the study from a *friend or family member* (15.9% vs. 6.9%), and an *online search* (19.1% vs. 12.1%) as most influential in their decisions to enroll in SPARK. Conversely, parents/legally authorized representatives were more likely than independent adults to endorse *speaking with a medical provider* (20.0% vs. 2.5%), *speaking with a study-team member at the doctor's office* (5.9% vs. 0.6%), and *speaking with the study team at a community event* (4.8% vs. 1.2%) as most influential in their decisions to enroll. Complete results can be seen in Table 7.

Discussion

We assessed study personnel's and research participants' perspectives about effective recruitment strategies for the national SPARK study on individuals with autism and their families. Queries of study personnel focused on strategies that they had tried, perceived costs, and those they would or would not repeat/recommend, whereas queries of SPARK participants focused on the various ways they learned about SPARK and, among those, which they deemed most influential in their decisions to enroll. Study personnel reported using a wide variety of recruitment strategies, which is consistent with recommendations in the literature [e.g., 15]. Relatively large proportions cited collaborations with medical providers and paid social media as effective strategies, which have also been supported in prior reports [e.g., 14]. Yet the recruitment methods that a majority of study personnel endorsed as most effective overall (i.e., would repeat/recommend) were those that involved direct engagement with the study team (i.e., *speaking with SPARK team members in clinic, on-site registration/saliva-collection events*). This is consistent with findings in Denhoff et al.[19], as well as those in Daniels et al.[21] in that clinical sites had higher numbers of participants who returned saliva kits compared to other forms of recruitment, suggesting that personal connections made between study staff and potential participants can positively influence engagement and retention of families in research. Because one of the primary objectives of SPARK is to identify genes associated with autism, ensuring saliva-kit return for DNA analysis is a critical feature of the study (although not a requirement for participation).

Regarding strategies that study personnel reportedly would *not* repeat or recommend to others (i.e., *blog post, radio PSA, radio or podcast interview*), relatively few participants had tried these approaches. Although radio advertising has been used for decades, reports of its success vary; and it can be expensive[14]—a sentiment also reported by personnel in Study 1. Blogs and podcasts, however, are newer communication platforms without much evidence for or against their adoption in participant recruitment. Researchers at institutions with existing podcast stations and/or blog outlets could leverage these tools for research recruitment and evaluate their reach/impact to contribute to this knowledge base, particularly considering that study personnel rated blog posts and podcast interviews among the less costly recruitment strategies. With respect to perceptions of cost for other strategies, most were rated as “least” or “somewhat” costly. It seems likely that “return on investment” factored into study personnel's efficacy ratings for each

strategy, as some strategies that were rated “more/most costly” were not likely to be repeated/recommended (e.g., *radio advertising*) whereas others were (e.g., *on-site registration/saliva-collection events*).

Comparing results from both studies, it is interesting to note that some of the recruitment strategies that study personnel rated as most effective were those least frequently indicated by SPARK participants as ways that they learned about the study. For example, among study personnel who had used an *online news story* or a *TV news story or interview*, more than 40% were highly likely and approximately a third were somewhat likely to repeat/recommend these strategies, whereas only 3.4% of SPARK participants said they learned about the study through a news story, and more than 60% of this group rated some other strategy as more influential in their decision to enroll. Similarly, few study personnel felt that their hospital/clinic website was an effective platform for recruitment ($n = 10$, 25.0%), yet SPARK participants rated *online search* as the third most common way that they heard about the study. Interestingly, while relatively few participants learned about the study via a member of the study team in clinic ($n = 334$, 8.1%) or at an event ($n = 228$, 5.5%), 54% and 68% of them, respectively, rated those methods as most influencing their decisions to enroll. Taken together, these observations highlight the value of understanding which recruitment strategies are most likely to *reach* and *resonate* with the target audience so that limited resources can be allocated to those activities.

Recommendations

Our findings have important implications for aiding research teams in the selection of recruitment strategies that facilitate timely enrollment of study participants. In line with suggestions to employ a variety of strategies, study personnel should consider developing recruitment plans with strategies that enhance overall awareness and engagement. For example, in the current study, 47% of SPARK participants said that they heard about the study via social media; 63% of this group said this was the most influential factor in their decision to enroll, while 11% of this group said that the online search was the most influential. This suggests that investing in paid social media that directs interested parties to a compelling website may work in concert to increase participant engagement. Similarly, 23% of SPARK participants said that they heard about the study via a medical provider; 72% of this group rated this method as the most influential in their decision to enroll, while 9% said it was speaking with a member of the study team during their clinical visit. At the same time, 13% of participants said they heard about the

study via a flyer or print material; 24% of this group rated this method as most influential, whereas another 24% said speaking with a medical provider was most influential. Taken together, this suggests that (a) engaging medical providers in study referrals, (b) ensuring that families speak with a study-team member during clinical visits, and (c) providing families with study flyers during recruitment conversations with medical providers and/or study staff may collectively enhance participant engagement more so than any of these strategies alone.

The most common and influential ways that participants learned about SPARK were fairly consistent across participant groups (i.e., *speaking with a medical provider, social media, friend/family member*). These likely represent “core” strategies that resonate with the general population, and for that reason, study teams should prioritize these strategies when outreach is broad and inclusive. Yet our results also revealed important differences in how individuals learn about and are influenced by *particular* recruitment strategies based on gender, race, ethnicity, and type of account holder (parent/legally authorized representative, independent adult with autism). Males more often reported hearing about the study through a *friend/family member*, which suggests that snowball sampling methods that encourage males to recruit other males in their social networks may effectively increase male representation in study samples. Similarly, compared to all other racial groups, Asian participants were much less likely to report hearing about SPARK through *social media* and more likely to report hearing about it via online searches. This could indicate that *paid Google advertising* or *paid website ads*—which were infrequently used by research personnel in Study 1—could be an effective strategy to increase study visibility in a way that appeals to diverse audiences, thereby enhancing representativeness. Additionally, independent adults with autism were much more likely to hear about the study and be influenced to join via *social media* and *online searches*, whereas parents/legally authorized representatives were more likely to enroll when they heard about the study in clinic (i.e., via *medical providers, study teams at the doctor’s office*). This suggests that social-media/online platforms as more effective in reaching adults with autism; however, it is important to note that most of the SPARK clinical sites were in pediatric departments/hospitals, so strategies that leveraged medical providers and in-clinic opportunities were targeted toward pediatric patients and their parents/guardians. It is possible that these same strategies could yield similar results when applied in adult-specialty practices and/or primary care. Taken together, the current findings

provide at least some evidence-based guidance for how to tailor research-recruitment activities to efficiently meet enrollment objectives.

Limitations

Our assessment of research personnel's perceptions about effective recruitment strategies did not collect information about participants' rationales for why they would or would not repeat/recommend particular outreach efforts. Although we queried their *perceptions* of cost for various recruitment strategies, this is not a metric of *actual* cost, the latter of which certainly has implications for whether and to what extent study personnel can pursue particular strategies. Likewise, we did not directly assess the efficacy of specific recruitment strategies; rather our objective was to describe and compare (1) perceived efficacy of various recruitment strategies from the perspective of research staff (Study 1) and (2) experiences of SPARK participants, who, by definition, had been successfully recruited. However, in Study 2, we did assess saliva-kit return in relation to how SPARK participants said they were recruited into the study, which provides some indication of strategies associated with participant retention/completion. Our companion paper (Daniels et al.[21]) also reported on factors associated with SPARK-study completion, one of which was affiliation with a clinical site. It follows that these participants were likely recruited into the study via strategies that clinical sites most often reported using (i.e., *speaking with SPARK team members in clinic, on-site registration/saliva-collection events, social media, speaking with a medical provider in clinic, printed materials*); however, further research should systematically evaluate rates of study enrollment/completion for these strategies to better assess their efficacy/success. Our survey of study personnel did not explicitly query use of tools within the EHR to aid recruitment efforts because, at the time of Study 1, only one site was actively piloting use of a Best Practice Advisory (BPA) to aid recruitment at the point of care. Finally, study personnel's results are based on their experiences recruiting for the SPARK study, which may not generalize to other types of research projects.

SPARK participants for Study 2 were recruited through Research Match, which is a part of SPARK that allows current participants to be notified of additional study opportunities for which they may be eligible. Those who agree to cooperate in Research Match studies are more motivated to engage in research and may not be representative of the broader SPARK-participant cohort. Likewise, given the criteria used to identify families for inclusion in Study 2, the resulting sample may differ from the overall SPARK sample, as well as that from our companion

paper (Daniels et al.[21]). Although we assessed SPARK participants' experiences with recruitment strategies by race and ethnicity, our efforts did not examine perspectives of anyone who did not speak English. At the time of data collection, SPARK was only available to individuals who were proficient in English; however, in 2022, SPARK became available in Spanish. It will be valuable to gather perceptions about effective recruitment strategies among these Spanish-speaking families in future work to inform efforts to increase diversity within research samples.

At the time of this publication, SPARK was still actively enrolling participants. The central SPARK team routinely evaluates cohort demographic data against national data to gauge representativeness. Additionally, in the last few years, diversity, equity, and inclusion (DEI) initiatives have been implemented to support recruitment/retention of underrepresented groups in the SPARK cohort (<https://sparkforautism.org/portal/page/diversity-equity-inclusion-statement/>).

References

1. Kost RG, Corregano LM, Rainer T, Melendez C, Collier BS. A data-rich recruitment core to support translational clinical research. *Clinical and Translational Science*. 2014; 8(2): 91-99.
2. Kitterman DR, Cheng SK, Dilts DM, Orwoll ES. The prevalence and economic impact of low-enrolling clinical studies at an academic medical center. *Academic Medicine*. 2011; 86(11): 1360–1366. doi.org/10.1097/ACM.0b013e3182306440
3. Nasser N, Grady D, Balke CW. Commentary: Improving participant recruitment in clinical and translational research. *Academic Medicine*. 2011; 86(11): 1334-1335.
4. Vogel A, Comtesse H, Rosner R. Challenges in recruiting and retaining adolescents with abuse-related posttraumatic stress disorder: Lessons learned from a randomized controlled trial. *Child Adolesc Psychiatry Ment Health*. 2020; 14: 14. doi.org/10.1186/s13034-020-00320-y
5. White C, Gilshenan K, Hardy J. A survey of the views of palliative care healthcare professionals towards referring cancer patients to participate in randomized controlled trials in palliative care. *Supportive Care in Cancer*. 2008; 16(12): 1397–1405. doi.org/10.1007/s00520-008-0441-1
6. Canedo JR, Miller ST, Myers HF, Sanderson M. Racial and ethnic differences in knowledge and attitudes about genetic testing in the US: Systematic review. *Journal of Genetic Counseling*. 2019; 28(3): 587–601. doi.org/10.1002/jgc4.1078
7. Corbie-Smith G, Thomas SB, Williams MV, et al. Attitudes and beliefs of African Americans toward participation in medical research. *J Gen Intern Med*. 1999; 14: 537–546. doi.org/10.1046/j.1525-1497.1999.07048.x
8. George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority research participation among African Americans, Latinos, Asian Americans, and Pacific Islanders. *American Journal of Public Health*. 2014;104(2): e16-e31. doi.org/10.2105/ajph.2013.301706

9. Luebbert R, Perez A. Barriers to clinical research participation among african americans. *Journal of Transcultural Nursing*. 2016; 27(5): 456–463. doi.org/10.1177/1043659615575578
10. Lai YS, Afseth JD. A review of the impact of utilizing electronic medical records for clinical research recruitment. *Society for Clinical Trials*. 2019; 16(2): 194-203.
11. Simons AR, Ahmed KL, Limon DL, Duhon GF, Marzano G, Goin-Kochel RP. Utilization of a best practice alert (BPA) at point-of-care for recruitment into a US-based autism research study. *Journal of Autism and Developmental Disorders*. 2023; 53: 359-369.
12. Duhon GF, Simon AR, Limon DL, Ahmed KL, Marzano G, Goin-Kochel RP. Use of a best practice alert (BPA) to increase diversity within a US-based autism research cohort. *Journal of Autism and Developmental Disorders*. 2023; 53, 370-377. doi.org/10.1007/s10803-021-05407-9
13. Chin Feman SP, Nguyen LT, Quilty MT, et al. Effectiveness of recruitment in clinical trials: An analysis of methods used in a trial for irritable bowel syndrome patients. *Contemporary Clinical Trials*. 2008; 29(2): 241-251. doi.org/10.1016/j.cct.2007.08.004
14. Ahmed KL, Simon AR, Dempsey JR, Samaco RC, Goin-Kochel RP. Evaluating two common strategies for research participant recruitment into autism studies: Observational study. *Journal of Medical Internet Research*. 2020; 22(9): e16752. doi: 10.2196/16752.
15. Frandsen M, Thow M, Ferguson SG. The effectiveness of social media (Facebook) compared with more traditional advertising methods for recruiting eligible participants to health research studies: A randomized, controlled clinical trial. *JMIR: Research Protocols*. 2016; 5(3): e161. doi: 10.2196/resprot.5747.
16. Watson JM, Torgerson DJ. Increasing recruitment to randomised trials: A review of randomised controlled trials. *BMC Medical Research Methodology*. 2006; 6, 34. doi.org/10.1186/1471-2288-6-34
17. Treweek S, Pitkethly M, Cook J, et al. Strategies to improve recruitment into randomized trials. *Cochrane Database of Systematic Reviews*. 2018; 2, Art. No.: MR000013.
18. Shneider C, Hilliard ME, Monaghan M, et al. Recruiting and retaining parents in behavioral intervention trials: Strategies to consider. *Contemporary Clinical Trials*. 2021; 108: 106502. doi.org/10.1016/j.cct.2021.106502

19. Denhoff ER, Milliren CE, deFerranti SD, Steltz SK, Osganian SK. Factors associated with clinical research recruitment in a pediatric academic medical center—a web-based survey. *PLoS One*. 2015; 10(10): e0140768.
20. Darden PM, Davis AM, Lee JY, et al. Active vs traditional methods of recruiting children for a clinical trial in rural primary care clinics: A cluster-randomized clinical trial. *JAMA Network Open*. 2022; 5(11): e2244040.
21. Daniels AM, Law JK, Green Snyder L., et al. Effectiveness of multimodal participant recruitment in SPARK, a large, online longitudinal research study of autism. *Journal of Clinical and Translational Science*. (under review).
22. SPARK Consortium. SPARK (Simons Foundation Powering Autism Research for Knowledge): a US cohort of 50,000 families to accelerate autism research. *Neuron*. 2018; 97(3): 488-493.

Acknowledgements

Co-author, Gabrielle Duhon, is now at MD Anderson Cancer Center, Houston, TX. The authors express their thanks to the SPARK team at the Simons Foundation, the SPARK clinical sites, and the SPARK cohort participants. We thank the following members of The SPARK Consortium for their efforts in creating, recruiting, and maintaining the SPARK cohort: Alexandria Aarrestad, Leonard Abbeduto, Gabriella Aberbach, Shelley Aberle, John Acampado, Andrea J. Ace, Adediwura Adegbite, Debbie Adeniji, Maria Aguilar, Kaitlyn Ahlers, Charles Albright, Michael Alessandri, Zach Algaze, Jasem Alkazi, Raquel Amador, David Amaral, Alpha Amatya, Logan Amon, Leonor Amundsen, Alicia Andrus, Claudine Anglo, Robert Annett, Adam Arar, Jonathan Arnold, Ivette Arriaga, Eduardo Arzate, Raven Ashley, Leilemah Aslamy, Irina Astrovsckaya, Kelli Baalman, Melissa Baer, Ethan Bahi, Joshua Bailey, Zachary Baldlock, Elissa Ball, Gabrielle Banks, Gabriele Baraghoshi, Nicole Bardett, Sarah Barns, Mallory Barrett, Yan Bartholomew, Asif Bashar, Heidi Bates, Katie Beard, Juana Becerra, Malia Beckwith, Paige Beechan, Landon Beeson, Josh Beeson, Brandi Bell, Monica Belli, Dawn Bentley, Natalie Berger, Anna Berman, Raphael Bernier, Elizabeth Berry-Kravis, Mary Berwanger, Shelby Birdwell, Elizabeth Blank, Rebecca Bond, Stephanie Booker, Aniela Bordofsky, Erin Bower, Lukas Bowers, Catherine Bradley, Heather Brayer, Stephanie Brewster, Elizabeth Brooks, Jannylyn Brown, Hallie Brown, Alison Brown, Melissa Brown, Catherine Buck, Cate Buescher, Kayleigh Bullon, Joy Buraima, Gabi Burkholz, Martin Butler, Eric Butter, Amalia Caamano, Nicole Cacciato, Wenteng Cai, Katina Calakos, Norma Calderon, Kristen Callahan, Alexies Camba, Claudia Campo-Soria, Giuliana Caprara, Paul Carbone, Laura Carpenter, Sarah Carpenter, Lindsey Cartner, Myriam Casseus, Lucas Casten, Sullivan Catherine, Alyss Cavanagh, Ashley Chappo, Sonny Chauhan, Kimberly Chavez, Randi Cheatham-Johnson, Tia Chen, Wubin Chin, Sharmista Chintalapalli, Daniel Cho, Dave Cho, YB Choi, Nia Clark, Renee Clark, Marika Coffman, Cheryl Cohen, Laura Coleman, Kendra Coleman, Alister Collins, Costanza Columbi, Joaquin Comitre, Stephanie Constant, Arin Contra, Sarah Conyers, Lindsey Cooper, Cameron Cooper, Leigh Coppola, Allison Corlett, Lady Corrales, Dahriana Correa, Hannah Cottrell, Michelle Coughlin, Eric Courchesne, Dan Coury, Deana Crocetti, Carrie Croson, Judith Crowell, Joseph Cubells, Sean Cunningham, Mary Currin, Michele Cutri, Sophia D'Ambrosi, Giancarla David, Ayana Davis, Sabrina Davis, Nickelle Decius, Andrea Deisher, Jennifer Delaporte, Lindsey DeMarco, Brandy Dennis, Alyssa Deronda, Esha Dhawan, Gabriel

Dichter, Christopher Diggins, Emily Dillon James Dixie, Ryan Doan, Kelli Dominick, Leonardo Dominquez Ortega, Erin Doyle, Andrea Drayton, Megan DuBois, Johnny Dudley, Gabrielle Duhon, Gabrielle Duncan, Amie Duncan, Megan Dunlevy, Meaghan Dyer, Rachel Earl, Dominique Earle, Catherine Edmonson, Sara Eldred, Nelita Elliott, Brooke Emery, Malak Enayetallah, Barbara Enright, Sarah Erb, Craig Erickson, Amy Esler, Liza Estevez, Anne Fanta, Carrie Fassler, Ali Fatemi, Faris Fazal, Marilyn Featherston, Jonathan Ferguson, Michael Finlayson, Angela Fish, Kate Fitzgerald, Chris Fleisch, Kathleen Flores, Eric Fombonne, Margaret Foster, Tiffany Fowler, Emma Fox, Emily Fox, Sunday Francis, Margot Frayne, Sierra Froman, Laura Fuller, Virginia Galbraith, Dakota Gallimore, Ariana Gambrell, Swami Ganesan, Vahid Gazestani, Madeleine R. Geisheker, Alicia Geng, Jennifer Gerdts, Daniel Geschwind, Mohammad Ghaziuddin, Haidar Ghina, Jessica Giulietti, Erin Given, Mykayla Goetz, Alexandra Goler, Jared Gong, Kelsey Gonring, Natalia Gonzalez, Antonio Gonzalez, Ellie Goodwill, Rachel Gordon, Carter Graham, Catherine Gray, Tunisia Greene, Ellen Grimes, Anthony Griswold, Luke Grosvenor, Pan Gu, Janna Guilfoyle, Amanda Gulsrud, Jaclyn Gunderson, Chris Gunter, Sanya Gupta, Abha Gupta, Anibal Gutierrez, Frampton Gwynette, Ghina Haidar, Melissa Hale, Monica Haley, Jake Hall, Lauren K. Hall, Jake Hall, Kira Hamer, Piper Hamilton, Bing Han, Nathan Hanna, Antonio Hardan, Christina Harkins, Eldric Harrell, Jill Harris, Nina Harris, Danaizha Harvey, Caitlin Hayes, Braden Hayse, Teryn Heckers, Kathryn Heerwagen, Daniela Hennelly, Lynette Herbert, Luke Hermle, Briana Hernandez, Jessica Hernandez, Clara Herrera, Amy Hess, Michelle Heyman, Lorrin Higgins, Brittani Hilscher Phillips, Kathy Hirst, Theodore Ho, Dabney Hofammann, Emily Hoffman, Margaret Hojlo, Alison Holbrook, Makayla Honaker, Jane Hong, Michael Hong, Gregory Hooks, Susannah Horner, Danielle Horton, Melanie Houchell, Dain Howes, Lark Huang-Storm, Samantha Hunter, Hanna Hutter, Emily Hyde, Teresa Ibanez, Kelly Ingram, Dalia Istephanous, Suma Jacob, Andrea Jarratt, Stanley Jean, Anna Jelinek, Bill Jennsen, Mary Johnson, Erica Jones, Mya Jones, Garland Jones, Mark Jones, Alissa Jorgenson, Roger Jou, Jessyca Judge, Luther Kalb, Taylor Kalmus, Sungeun Kang, Elizabeth Kangas, Stephen Kanne, Oumou Kanoute, Hannah Kaplan, Sara Khan, Sophy Kim, Annes Kim, Alex Kitaygordsky, Cheryl Klaiman, Adam Klever, Hope Koene, Tanner Koomar, Misia Kowanda, Melinda Koza, Sydney Kramer, Meghan Krushena, Eva Kurtz-Nelson, Elena Lamarche, Erica Lampert, Martine Lamy, Rebecca Landa, Alex E. Lash, Noah Lawson, Alexa Lebron-Cruz, Holly Lechniak, Soo Lee, Bruce Leight, Matthew Lerner, Laurie Leshner, Courtney

Lewis, Hai Li, Deana Li, Robin Libove, Natasha Lillie, Danica Limon, Desi Limpoco, Melody Lin, Sandy Littlefield, Nathan Lo, Brandon Lobisi, Laura Locarno, Nancy Long, Bailey Long, Kennadie Long, Marilyn Lopez, Taylor Lovering, Ivana Lozano, Marti Luby, Daniella Lucio, Addie Luo, My-Linh Luu, Audrey Lyon, Julia Ma, Natalie Madi, Malcolm Mallardi, Lacy Malloch, Anup Mankar, Reanna Mankaryous, Lori Mann, Patricia Manning, Julie Manoharan, Alvin Mantey, Olena Marchenko, Richard Marini, Alexandra Marsden, Clarissa Marwali, Gabriela Marzano, Andrew Mason, Sarah Mastel, Sheena Mathai, Emily Matthews, Emma Matusoff, Clara Maxim, Caitlin McCarthy, Lynn McClellen, Nicole Mccoy, James McCracken, Kaylen McCullough, Brooke McDonald, Julie McGalliard, Anne-Marie McIntyre, Brooke McKenna, Alexander McKenzie, Megan McTaggart, Hannah Meinen, Sophia Melnyk, Alexandra Miceli, Sarah Michaels, Jacob Michaelson, Estefania Milan, Melissa Miller, Anna Milliken, Kyla Minton, Terry Mitchell, Amanda Moffitt Gunn, Sarah Mohiuddin, Gina Money, Gabriela Montanez, Jessie Montezuma, Lindsey Mooney, Margo Moore, Amy Morales-Lara, Kelly Morgan, Hadley Morotti, Michael Morrier, Maria Munoz, Ambar Munoz Lavanderos, Shwetha Murali, Karla Murillo, Kailey Murray, Vincent J. Myers, Erin Myhre, Natalie Nagpal, Jason Neely, Emily Neuhaus, Olivia Newman, Darnell Newsum, Ai Nhu Nguyen, Richard Nguyen, Victoria Nguyen, Ai Nhu Nguyen, Evelyn Nichols, Amy Nicholson, Melanie Niederhauser, Megan Norris, Shai Norton, Kerri Nowell, Kaela O'Brien, Eirene O'Connor, Mitchell O'Meara, Molly O'Neil, Brian O'Roak, Edith Ocampo, Cesar Ochoa-Lubinoff, Anna Oft, Jessica Orobio, Elizabeth Orrick, Crissy Ortiz, Opal Ousley, Motunrayo Oyeyemi, Lillian Pacheco, Valeria Palacios, Samiza Palmer, Isabella Palmeri, Katrina Pama, Juhi Pandey, Anna Marie Paolicelli, Jaylaan Parker, Morgan Patterson, Melissa Patxot, Katherine Pawlowski, Ernest Pedapati, Michah Pepper, Esmeralda Perez, Jeremy Perrin, Christine Peura, Diamond Phillips, Karen Pierce, Joseph Piven, Juhi Plate, Jose Polanco, Jibrielle Polite, Natalie Pott-Schmidt, Tiziano Pramparo, Taleen Pratt, Lisa Prock, Stormi Pulver White, Hongjian Qi, Shanping Qiu, Eva Queen, Marcia Questel, Ashley Quinones, Nihal Raman, Desiree Rambeck, Rishiraj Rana, Shelley Randall, Vaikunt Ranganathan, Laurie Raymond, Madelyn Rayos, Kelly Real, Richard Remington, Anna Rhea, Catherine Rice, Harper Richardson, Stacy Riffle, Chris Rigby, Tracy Robertson, Beverly Robertson, Erin Roby, Ana Rocha, Casey Roche, Nicki Rodriguez, Bianca Rodriguez, Katherine Roeder, Laure Rogers, Daniela Rojas, Andrea Rondon, Jacob Rosewater, Hilary Rosselott, Katelyn Rossow, Payton Runyan, Nicole Russo, Tara Rutter,

Elizabeth Ruzzo, Mahfuza Sabiha, Mustafa Sahin, Fatima Salem, Rebecca Sanchez, Muave Sanders, Tayler Sanderson, Sophie Sandhu, Katelyn Sanford, Susan Santangelo, Madeline Santulli, Alisha Sarakki, Marina Sarris, Dustin Sarver, Madeline Savage, Jessica Scherr, Abraham Schneider, Hoa Schneider, Abraham Schneider, Hayley Schools, Gregory Schoonover, Robert Schultz, Brian Schwanwede, Brady Schwind, Tanner Scott, Cheyanne Sebolt, Rebecca Shaffer, Neelay Shah, Swapnil Shah, Sana Shameen, Curry Sherard, Jonah Shifrin, Roman Shikov, Amelle Shillington, Mojeeb Shir, Amanda Shocklee, Clara Shrier, Lisa Shulman, Matt Siegel, Andrea Simon, Laura Simon, Emily Singer, Arushi Singh, Vini Singh, Devin Smalley, Kaitlin Smith, Chris Smith, Ashlyn Smith, Latha Soorya, Julia Soscia, Aubrie Soucy Diksha Srishyla Danielle Stamps, Laura Stchur, Morgan Steele, Alexandra Stephens, Dominique Stilletti Camilla Strathearn, Nicole Sussman, Amy Swanson, Megan Sweeney, Anthony Sziklay, Maira Tafolla, Jabeen Taiba, Nicole Takahashi, Sydney Terroso, Priyam Thind, Taylor Thomas, Hanna Thomas, Samantha Thompson, Jennifer Tjernagel, Jaimie Toroney, Ellyn Touchette, Laina Townsend, Madison Trog, Katherine Tsai, Angela Tseng, Paullani Tshering, Ivy Tso, Laura Tung. Samuel Turecki, Maria Valicenti-Mcdermott, Bonnie VanMetre, Candace VanWade, Alfonso Vargas-Cortes, Kerrigan Vargo, Dennis Vasquez Montes, Cristiana Vattuone, Jeremy Veenstra-Vanderweele, Alison Vehorn, Alan Jesus Benitez Velazquez, Mary Verdi, Brianna Vernoia, Michele Villalobos, Natalia Volfovsky, Jasmine Vongdara, Lakshmi Vrittamani, Allison Wainer, Jermel Wallace, Corrie Walston, Jiayaho Wang, Audrey Ward, Zachary Warren, Katherine Washington, Lucy Wasserberg, Grace Westerkamp, Casey White, Sabrina White, Casey White, Logan Wink, Fiona Winoto, Sarah Winters, Ericka Wodka, Aaron Wong, JessicaWright, Samantha Xavier, Sabrina Xiao, Simon Xu, Sidi Xu, Yi Yang, WhaJames Yang, Amy Yang, Meredith Yinger, Timothy Yu, Christopher Zaro, Hana Zaydens, Cindy Zha, Haicang Zhang, Haoquan Zhao, Allyson Zick, and Lauren Ziegelmayr Salmon.


Funding Statement

Although the authors contribute to SPARK as a clinical site (Simons Foundation Autism Research Initiative grant #385052, RPG-K), this project was initiated by the authors without specific or supplemental funding. A portion of RPG-K's effort was partially supported by the Intellectual and Developmental Disabilities Research Center (1U54 HD083092) at Baylor College of Medicine.

Author Contributions

Authors contributed to the manuscript in the following ways: Conception and design of the work (Goin-Kochel, Duhon, Marzano), collection or contribution of data (Goin-Kochel, Duhon, Marzano), contributions of analysis tools or expertise (Goin-Kochel, Duhon, Daniels), conduct and interpretation of analysis (Goin-Kochel, Daniels, Law, Diehl, Green Snyder, Feliciano, Chung), drafting of the manuscript (Goin-Kochel, Lozano, Daniels, Chung), critical review/feedback on manuscript (Goin-Kochel, Lozano, Duhon, Marzano, Daniels, Law, Diehl, Green Snyder, Feliciano, Chung). Robin Goin-Kochel takes responsibility for the manuscript, as a whole.

Figure 1. Participant Ratings of Most Influential Strategies Among Ways Heard About SPARK*



*Note. Among participants who indicated that they learned about SPARK through a given strategy, we calculated the proportion who then rated that strategy as most influential in their decision to enroll. For example, among participants who said they learned about SPARK through a flyer/print material, 24% rated this outlet as most influential in their decision to enroll, while 76% rated an alternate outlet as most influential.

Table 1. Recruitment Strategies Employed for SPARK and Study Personnel’s Frequency of Use

Recruitment Strategy	Description/Examples	Use Frequency ^a (%)
Print materials	SPARK brochures, post cards, flyers, posters, etc.	52 (100%)
Paid social media	Facebook, Twitter, Instagram sponsored ads	44 (84.6%)
Non-paid social media	Organically sharing/posting on hospital or clinic social media pages, autism-related groups, or via community organizations who share on your behalf	45 (86.5%)
Paid radio advertising	Paid radio campaign/advertising with a pre-recorded script that runs for a specified number of weeks at specific times	20 (38.5%)
Radio or podcast interview	Invitation by a radio/podcast host to interview you about SPARK on air/for a pre-recorded podcast, at no cost to you	20 (38.5%)
Radio PSA ^b	A script or pre-recorded audio file submitted to various radio stations to share if they so choose; PSAs are not guaranteed slots and must be “accepted,” but are free-of-charge because they benefit the community	10 (19.2%)
Mass letter mailing (500+)	Letters mailed to a large number of families/stakeholders inviting them to join SPARK. This may include patient registries, clinic lists, data pulls from the electronic health record, newsletter distribution list, etc.	32 (61.5%)
Medical providers in clinic	Medical providers sharing print materials and information with potentially eligible families during previously scheduled clinic appointments at your site	48 (92.3%)
Blog post	Current participant, study team member, clinical provider, or other stakeholder writes a blog that directly relates to the SPARK study/directs families to a SPARK-related website	13 (25.0%)
Mass email blasts (500+, including newsletters)	Emails sent to a large number of families/stakeholders inviting them to join SPARK. This may include patient registries, clinic lists, data pulls from the electronic health record, newsletter distribution list, etc.	41 (78.8%)
Online news story	Interview/story published online by a news station/group, or a news station sharing a press release about SPARK from your institution	34 (65.4%)
TV news story or interview	Invitation by a TV news or talk-show host to interview you/study personnel about SPARK, to be played on air	35 (67.3%)
TV PSA ^b	Submitting a video file to various TV stations to share freely, if they so choose. PSAs are not guaranteed slots and must be “accepted,” but are free-of-charge because they benefit the community	8 (15.4%)
Paid Google advertising	Paid advertising campaigns for Google that result in your information showing up “first” in search results for specified key words	4 (7.7%)
Paid website	Paying a website to place your information as a banner	11 (21.2%)

Recruitment Strategy	Description/Examples	Use Frequency ^a (%)
advertising	ad, side-bar ad, etc., for their viewers to see (e.g., news websites, community-partner sites)	
Hospital/clinic official website	Listing SPARK on any webpage owned by your institution (e.g., www.bcm.edu , www.texaschildrens.org) This could include a faculty lab page, a larger list of research opportunities at your institution, a specific center's page, etc.	40 (76.9%)
SPARK coordinators in clinic	SPARK coordinators meeting with families as they come through clinic for previously scheduled appointments, or inviting families into clinic specifically to meet with a SPARK coordinator	43 (82.7%)
Walks, informational events, conferences	Attending local or distant autism-related events/conferences as an exhibitor or speaker to share information about SPARK	49 (94.2%)
On-site registration/saliva-collection events	Inviting families to register/collect saliva in person with the study team as part of an outreach event	45 (86.5%)
Transit advertising	Placing banner ads on city transit systems (e.g., buses, subways, trains)	7 (13.5%)

Note: ^aN = 52. ^bPSA = Public service announcement.

Table 2. Recruitment Strategies Employed for SPARK and Study Personnel’s Ratings of Cost*

Recruitment Strategy	Least Somewhat Costly <i>n</i> (%)	or Moderately Costly <i>n</i> (%)	More or Most Costly <i>n</i> (%)
Print materials (<i>n</i> = 51)	23 (45.1%)	21 (41.2%)	7 (13.7%)
Paid social media (<i>n</i> = 43)	19 (44.2%)	13 (30.2%)	11 (25.6%)
Non-paid social media (<i>n</i> = 45)	44 (97.8%)	1 (2.2%)	0 (0%)
Paid radio advertising (<i>n</i> = 19)	2 (10.5%)	5 (26.3%)	12 (63.1%)
Radio or podcast interview (<i>n</i> = 19)	15 (78.9%)	2 (10.5%)	2 (10.5%)
Radio PSA ^b (<i>n</i> = 10)	8 (80.0%)	10 (10.0%)	10 (10.0%)
Mass letter mailing (500+) (<i>n</i> = 32)	12 (37.5%)	12 (37.5%)	8 (25.0%)
Medical providers in clinic (<i>n</i> = 47)	38 (80.8%)	6 (12.8%)	3 (6.4%)
Blog post (<i>n</i> = 12)	11 (91.7%)	1 (8.3%)	0 (0%)
Mass email blasts (500+, including newsletters) (<i>n</i> = 40)	39 (97.5%)	1 (2.5%)	0 (0%)
Online news story (<i>n</i> = 33)	28 (84.8%)	2 (6.1%)	3 (9.1%)
TV news story or interview (<i>n</i> = 34)	29 (85.3%)	4 (11.8%)	1 (2.9%)
TV PSA ^b (<i>n</i> = 7)	6 (85.7%)	1 (14.3%)	0 (0%)
Paid Google advertising (<i>n</i> = 4)	2 (50.0%)	2 (50.0%)	0 (0%)
Paid website advertising (<i>n</i> = 10)	3 (30.0%)	4 (40.0%)	3 (30.0%)
Hospital/clinic official website (<i>n</i> = 39)	38 (97.4%)	1 (2.6%)	0 (0%)
SPARK coordinators in clinic (<i>n</i> = 42)	28 (66.7%)	9 (21.4%)	5 (11.9%)
Walks, informational events, conferences (<i>n</i> = 48)	24 (50.0%)	14 (29.2%)	10 (20.8%)
On-site registration/saliva-collection events (<i>n</i> = 44)	14 (31.8%)	14 (31.8%)	16 (36.3%)
Transit advertising (<i>n</i> = 6)	0 (0%)	0 (0%)	6 (100%)

Note: *Study personnel were asked: “Besides personnel and basic office supplies, how costly is it to execute this recruitment strategy at your site?”

^a*N* = 52. ^bPSA = Public service announcement.

Table 3. Recruitment Strategies Employed for SPARK and Study Personnel's Ratings of Efficacy

Recruitment Strategy	Unlikely to Repeat/ Recommend <i>n</i> (%)	Somewhat Likely to Repeat/ Recommend <i>n</i> (%)	Highly Likely to Repeat/ Recommend <i>n</i> (%)
Print materials (<i>n</i> = 52)	10 (19.2%)	15 (28.9%)	27 (51.9%)
Paid social media (<i>n</i> = 35)	9 (20.5%)	9 (20.5%)	26 (59.0%)
Non-paid social media (<i>n</i> = 46)	14 (30.4%)	14 (30.4%)	18 (39.1%)
Paid radio advertising (<i>n</i> = 20)	9 (45.0%)	8 (40.0%)	3 (15.0%)
Radio or podcast interview (<i>n</i> = 21)	11 (52.4%)	7 (33.3%)	3 (14.3%)
Radio PSA ^b (<i>n</i> = 9)	6 (66.7%)	2 (22.2%)	1 (11.1%)
Mass letter mailing (500+) (<i>n</i> = 33)	12 (36.4%)	7 (21.2%)	14 (42.4%)
Medical providers in clinic (<i>n</i> = 48)	14 (29.2%)	9 (18.8%)	25 (52.0%)
Blog post (<i>n</i> = 14)	10 (71.4%)	3 (21.4%)	1 (7.2%)
Mass email blasts (500+, including newsletters) (<i>n</i> = 41)	12 (29.3%)	10 (24.4%)	19 (46.3%)
Online news story (<i>n</i> = 34)	10 (29.4%)	10 (29.4%)	14 (41.2%)
TV news story or interview (<i>n</i> = 35)	8 (22.8%)	12 (34.3%)	15 (42.9%)
TV PSA ^b (<i>n</i> = 8)	3 (37.5%)	2 (25.0%)	3 (37.5%)
Paid Google advertising (<i>n</i> = 4)	2 (50.0%)	2 (50.0%)	0 (0%)
Paid website advertising (<i>n</i> = 9)	3 (33.3%)	3 (33.3%)	3 (33.3%)
Hospital/clinic official website (<i>n</i> = 40)	17 (42.5%)	13 (32.5%)	10 (25.0%)
SPARK coordinators in clinic (<i>n</i> = 43)	2 (4.7%)	4 (9.3%)	37 (86.0%)
Walks, informational events, conferences (<i>n</i> = 49)	12 (24.5%)	13 (26.5%)	24 (49.0%)
On-site registration/saliva-collection events (<i>n</i> = 44)	9 (20.5%)	3 (6.8%)	32 (72.7%)
Transit advertising (<i>n</i> = 6)	2 (33.3%)	2 (33.3%)	2 (33.3%)

Note: ^a*N* = 52. ^bPSA = Public service announcement.

Table 4. Frequencies of Participants Who Endorsed Learning About SPARK Through Various Recruitment Outlets and Ratings of Those Most Influential in Their Decisions to Enroll

Recruitment Strategy	Heard About SPARK This Way ^a	Most Influential in Decision to Enroll
Flyer or print material	529 (12.8%)	150 (3.6%)
Radio advertisement	38 (0.9%)	11 (0.3%)
Mailed letter	219 (5.3%)	108 (2.6%)
Friend/family member ^b	549 (13.2%)	330 (8.0%)
Transit advertisement ^c	14 (0.3%)	6 (0.1%)
Newspaper	20 (0.5%)	8 (0.2%)
Social media ^d	1,946 (47.0%)	1,270 (30.6%)
Online search ^e	833 (20.1%)	537 (13.0%)
News story ^f	140 (3.4%)	67 (1.6%)
Spoke with medical provider ^g	958 (23.1%)	733 (17.7%)
Spoke with team in clinic ^h	334 (8.1%)	215 (5.2%)
Spoke with team at event ⁱ	228 (5.5%)	180 (4.3%)
Newsletter/bulk email	170 (4.1%)	76 (1.8%)
Personal email from SPARK team	377 (9.1%)	235 (5.7%)
Other	295 (7.1%)	202 (4.9%)

Note: ^aSpecifically refers to ways heard about the study prior to enrollment. Participants were instructed to select all that apply, so proportions exceed 100%. ^bIncluded any person not directly involved in the study. ^cOn a bus, train, etc. ^dFacebook, Twitter, etc. ^eIncluded the SPARK website or hospital/clinic website. ^fOnline or TV. ^gDoctor, nurse, therapist, etc. ^hSpoke with SPARK staff in clinic before or after doctor's appointment. ⁱSpoke with SPARK staff at a community event (resource fairs, walks, conferences, etc.).

Table 5. Most Influential Recruitment Strategies by Gender and Age

Recruitment Strategy	Gender ^a		Age (in years)			
	Male	Female	18-29	30-44	45-60	60+
Flyer or print material	15 (3.4%)	132 (3.6%)	6 (1.6%)	93 (3.7%)	49 (4.2%)	2 (2.6%)
Radio advertisement	0 (0%)	11 (0.3%)	0 (0%)	10 (0.4%)	1 (0.1%)	(0%)
Mailed letter	8 (1.8%)	97 (2.7%)	1 (0.3%)	64 (2.5%)	40 (3.4%)	2 (2.6%)
Friend/family member ^b	53 (12.1%)	267 (7.4%)	50 (13.2%)	182 (7.2%)	88 (7.6%)	10 (13.0%)
Transit ad ^c	3 (0.7%)	2 (0.1%)	2 (0.5%)	3 (0.1%)	0 (0%)	1 (1.3%)
Newspaper	1 (0.2%)	7 (0.2%)	1 (0.3%)	5 (0.2%)	2 (0.2%)	0 (0%)
Social media ^d	91 (20.8%)	1144 (31.5%)	161 (42.4%)	825 (32.8%)	272 (23.4%)	10 (13.0%)
Online search ^e	75 (17.1%)	454 (12.5%)	50 (13.2%)	309 (12.3%)	165 (14.2%)	13 (16.9%)
News story ^f	13 (3.0%)	53 (1.5%)	2 (0.5%)	34 (1.4%)	26 (2.2%)	5 (6.5%)
Medical provider ^g	76 (17.4%)	648 (17.9%)	53 (13.9%)	473 (18.8%)	192 (16.5%)	13 (16.9%)
Team in clinic ^h	13 (3.0%)	202 (5.6%)	17 (4.5%)	151 (6.0%)	47 (4.0%)	0 (0%)
Team at event ⁱ	21 (4.8%)	158 (4.4%)	5 (1.3%)	115 (4.6%)	58 (5.0%)	2 (2.6%)
Newsletter/bulk email	10 (2.3%)	65 (1.8%)	6 (1.6%)	32 (1.3%)	36 (3.1%)	2 (2.6%)
Personal email ^j	30 (6.8%)	203 (5.6%)	14 (3.7%)	101 (4.0%)	111 (9.5%)	8 (10.4%)
Other	26 (5.9%)	172 (4.7%)	9 (2.4%)	115 (4.6%)	68 (5.8%)	9 (11.7%)

Note: ^a20 participants reported as transgender; 13 (65.0%) rated social media as most influential in their decision to enroll. 38 participants reported as gender queer/gender nonconforming; 16 (42.1%) rated social media as most influential in their decision to enroll, while 12 (31.6%) rated friend/family member or an online search as most influential. 2 participants reported as some different gender identity; 1 (50.0%) rated social media and 1 (50.0%) rated speaking with a medical provider as most influential in their decision to enroll. 10 participants preferred not to answer about gender; 6 (60.0%) rated social media or friend/family member as most influential in their decision to enroll. ^bIncluded any person not directly involved in the study. ^cOn a bus, train, etc. ^dFacebook, Twitter, etc. ^eIncluded the SPARK website or hospital/clinic website. ^fOnline or TV. ^gSpoke with a medical provider (doctor, nurse, therapist, etc.) ^hSpoke with SPARK staff in clinic before or after doctor's appointment. ⁱSpoke with SPARK staff at a community event (resource fairs, walks, conferences, etc.). ^jPersonal email from a study-team member.

Table 6. Most Influential Recruitment Strategies by Hispanic/Latinx Ethnicity and Race

Recruitment Strategy	Ethnicity	Race ^a					
		AS	AA	AI	PI	W	Oth
Flyer or print material	25 (5.7%)	6 (4.5%)	16 (6.4%)	2 (2.0%)	0 (0%)	118 (3.2%)	12 (7.8%)
Radio advertisement	3 (0.7%)	0 (0%)	0 (0%)	1 (1.0%)	0 (0%)	10 (0.3%)	0 (0%)
Mailed letter	9 (2.1%)	4 (3.0%)	4 (1.6%)	2 (2.0%)	0 (0%)	98 (2.7%)	5 (3.2%)
Friend/family member ^b	26 (6.0%)	6 (4.5%)	17 (6.8%)	6 (6.1%)	1 (4.8%)	299 (8.1%)	13 (8.4%)
Transit ad ^c	2 (0.5%)	0 (0%)	1 (0.4%)	0 (0%)	0 (0%)	5 (0.1%)	1 (0.6%)
Newspaper	1 (0.2%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	8 (0.2%)	0 (0%)
Social media ^d	127 (29.1%)	20 (15.2%)	76 (30.3%)	41 (41.4%)	9 (42.9%)	1157 (31.5%)	39 (25.3%)
Online search ^e	66 (15.1%)	29 (22.0%)	28 (11.2%)	16 (16.2%)	3 (14.3%)	465 (12.7%)	26 (16.9%)
News story ^f	1 (0.2%)	3 (2.3%)	6 (2.4%)	1 (1.0%)	0 (0%)	60 (1.6%)	1 (0.6%)
Medical provider ^g	76 (17.4%)	24 (18.2%)	40 (15.9%)	14 (14.1%)	4 (19.0%)	647 (17.6%)	24 (15.6%)
Team in clinic ^h	24 (5.5%)	10 (7.6%)	14 (5.6%)	4 (4.0%)	1 (4.8%)	186 (5.1%)	8 (5.2%)
Team at event ⁱ	19 (4.4%)	7 (5.3%)	12 (4.8%)	4 (4.0%)	1 (4.8%)	157 (4.3%)	5 (3.2%)
Newsletter/bulk email	10 (2.3%)	5 (3.8%)	6 (2.4%)	1 (1.0%)	1 (4.8%)	63 (1.7%)	4 (2.6%)
Personal email ^j	24 (5.5%)	8 (6.1%)	21 (8.4%)	3 (3.0%)	0 (0%)	199 (5.4%)	7 (4.5%)
Other	18 (4.1%)	8 (6.1%)	10 (4.0%)	3 (3.0%)	1 (4.8%)	183 (5.0%)	9 (5.8%)

Note: ^aRacial categories are abbreviated as follows: AS = Asian, AA = Black/African American, AI = American Indian/Alaska Native, PI = Native Hawaiian or Other Pacific Islander, W = White, Oth = Other race. Participants had the option to select any/all racial categories to describe themselves. ^bIncluded any person not directly involved in the study. ^cOn a bus, train, etc. ^dFacebook, Twitter, etc. ^eIncluded the SPARK website or hospital/clinic website. ^fOnline or TV. ^gSpoke with a medical provider (doctor, nurse, therapist, etc.) ^hSpoke with SPARK staff in clinic before or after doctor's appointment. ⁱSpoke with SPARK staff at a community event (resource fairs, walks, conferences, etc.). ^jPersonal email from a study-team member.

Table 7. Most Influential Recruitment Strategies by Participant Role

Recruitment Strategy	Participant Role	
	Independent Adult with Autism	Parent/LAR ^a
Flyer or print material	6 (1.2%)	144 (4.0%)
Radio advertisement	1 (0.2%)	10 (0.3%)
Mailed letter	4 (0.8%)	104 (2.9%)
Friend/family member ^b	83 (15.9%)	247 (6.9%)
Transit advertisement ^c	4 (0.8%)	2 (0.1%)
Newspaper	1 (0.2%)	7 (0.2%)
Social media ^d	229 (43.9%)	1041 (28.9%)
Online search ^e	100 (19.1%)	437 (12.1%)
News story ^f	13 (2.5%)	54 (1.5%)
Medical provider ^g	13 (2.5%)	720 (20.0%)
Team in clinic ^h	3 (0.6%)	212 (5.9%)
Team at event ⁱ	6 (1.2%)	174 (4.8%)
Newsletter/bulk email	6 (1.2%)	70 (1.9%)
Personal email ^j	27 (5.2%)	208 (5.8%)
Other	27 (5.2%)	175 (4.9%)

Note: ^aLAR = legally authorized representative. ^bIncluded any person not directly involved in the study. ^cOn a bus, train, etc. ^dFacebook, Twitter, etc. ^eIncluded the SPARK website or hospital/clinic website. ^fOnline or TV. ^gSpoke with a medical provider (doctor, nurse, therapist, etc.) ^hSpoke with SPARK staff in clinic before or after doctor's appointment. ⁱSpoke with SPARK staff at a community event (resource fairs, walks, conferences, etc.). ^jPersonal email from a study-team member.