



Associations between dietary variety, portion size and body weight: prospective evidence from UK Biobank participants

Rochelle Embling*, Menna J. Price, Michelle D. Lee, Alex Jones† and Laura L. Wilkinson†

School of Psychology, Faculty of Medicine, Health and Life Science, Swansea University, Swansea, SA2 8PP, UK

(Submitted 14 June 2022 – Final revision received 19 December 2022 – Accepted 9 January 2023 – First published online 16 January 2023)

Abstract

‘Dietary variety’ has been identified as a factor associated with food intake. Whilst this relationship may have longer-term benefits for body weight management when eating low-energy, nutrient-dense foods, it may increase the risk of overconsumption (and body adiposity) when foods are high energy density. This study sought to further explore pathways underpinning the relationship between dietary variety and body weight, by considering energy density as a moderating factor and portion size as a mediating factor in this relationship. Using prospective data from the UK Biobank, dietary variety scores (DVS), cumulative portion size and energy density were derived from 24-h dietary recall questionnaires at baseline and follow-up. BMI, whole-body fat percentage and fat-free mass were included as outcomes. Contrary to predictions, linear multiple regression models found some evidence of a negative, direct association between DVS and body weight outcomes at baseline ($b = -0.13$). Though dietary variety was significantly associated with larger portions across time points ($b = 41.86-82.64$), a moderated mediation effect was not supported at baseline or follow-up (Index ≤ 0.035). Taken together, these findings provide population-level evidence to support a positive association between variety and food intake, which in turn has potential implications for body weight management, both in terms of moderating food intake and benefitting diet quality.

Key words: Dietary diversity: Food variety: Portion size: Energy density: Body weight: UK Biobank

Diet quality has been highlighted as a key target for interventions aiming to reduce population disease risk^(1,2). Better diet quality has been associated with having lower risk of CVD⁽³⁾, diabetes^(4,5), and cancer⁽⁶⁾, as well as overweight and obesity⁽⁷⁾. One approach used to indicate diet quality is the calculation of dietary variety/diversity scores (i.e. number of different food items consumed in a recall period⁽⁸⁾), as variety is believed to be an effective indicator of nutritional adequacy^(9,10).

In addition to diet quality, several epidemiological studies have reported positive associations between dietary variety scores (DVS) and food intake⁽¹¹⁾, indicating increased energy or weight of food consumed in relation to variety^(12–21). This association likely benefits the consumer when dietary variety consists of typically low energy density, nutrient-dense foods (reflecting better diet quality) but highlights potential risk of overconsumption in response to highly palatable, high energy density foods that more often characterise dietary patterns today^(22–24). For example, where greater overall dietary diversity was associated with poor diet quality, data from the Multi-Ethnic Study of Atherosclerosis (MESA) cohort showed that waist

circumference measurements more than doubled for US adults with the highest food dissimilarity scores during a 5-year period⁽²⁵⁾.

For these reasons, multiple reviews have highlighted a nuanced relationship between dietary variety and body weight whilst accounting for energy density, whereby consuming a greater variety of low (high) energy density foods is negatively (positively) associated with body adiposity^(11,26–28). In particular, such an effect has been supported when measuring consumption of variety within specific food groups that differ in energy density (e.g. ‘sweets, snacks and carbohydrates’ *v.* ‘vegetables’)⁽²⁹⁾. In a key cross-sectional study of seventy-one ‘healthy’ men and women in the USA, McCrory and colleagues⁽²⁹⁾ found that variety was significantly and positively related to energy intake across food groups but showed that greater variety from within vegetables was significantly associated with lower body adiposity, despite positive associations for all other food groups. Since then, dietary intake studies have specifically highlighted the significant relationship between variety within ‘recommended’ food groups and lower body weight/body

Abbreviation: DVS, dietary variety scores; IMD, index of multiple deprivation; DVS-B, dietary variety between food groups; DVS-W, dietary variety within food groups; BMI, body mass index.

* **Corresponding author:** Rochelle Embling, emails r.j.embling@swansea.ac.uk; rochelle.embling@outlook.com

† These authors should be considered joint last author.





adiposity^(13,16,30–32), though the most consistent evidence has been reported for opposing associations within ‘non-recommended’ food groups⁽¹¹⁾. Similar conclusions have been drawn from intervention studies, where greater variety in response to fruits and vegetables⁽³³⁾, and/or restricted variety for ‘less healthful’ foods^(33–35), have both been highlighted as a means to reduce energy intake and support weight loss⁽³⁶⁾.

Experimental studies in the laboratory also suggest that variety may drive the selection of larger portions. Including a greater variety of foods within and across meals – that differ in their sensory characteristics – has consistently been shown to increase consumption across food types⁽³⁷⁾. Repeated exposure to foods (and less variety) appears to result in faster habituation overtime, as individuals behaviourally and physiologically respond less to foods that are sensorially the same or similar to those previously eaten^(38–40). As such, variety has also been shown to increase the size of planned portions when selecting foods before eating⁽⁴¹⁾, and portion size itself is known to significantly influence food intake⁽⁴²⁾. Over a longer period, such evidence suggests that dietary variety may influence body weight outcomes if significant effects on cumulative portion size are observed across eating occasions (akin to the variety effect in a mealtime context). Roberts and colleagues⁽¹⁹⁾ in particular have highlighted energy intake as a factor that reduces the strength of dietary variety as a predictor of BMI when included in models. However, though DVS have been associated with increased energy intake using self-report measures in observational studies (see also above)⁽¹¹⁾, to our knowledge, no study has directly explored cumulative portion size as a mediating factor by which dietary variety may also influence body weight outcomes at a population level, particularly when considering the moderating role of energy density discussed across these literatures.

Therefore, this study aimed to further explore the relationship between dietary variety and body weight in a large cohort using secondary data from the UK Biobank ([https://www.ukbiobank.](https://www.ukbiobank.ac.uk/)

[ac.uk/](https://www.ukbiobank.ac.uk/)) (see Fig. 1). It was predicted that higher DVS would be associated with increased body weight and body fatness when energy density was high. It was also predicted that portion size would mediate this relationship, meaning higher DVS would be associated with a higher body weight and body fatness when individuals reported greater food intake (when energy density was high), constituting a moderated mediation effect.

Method

UK Biobank procedures

Prospective data from the UK Biobank were used in this study to analyse associations between dietary variety, portion size and body weight outcomes (Project ID: 53 159). The UK Biobank is a large-scale, cohort study including > 500 000 participants aged 40–69 years old, living within the UK⁽⁴³⁾. During baseline assessment centre visits, written informed consent was obtained from all participants. Participants completed questionnaires via a touch-screen and verbal interview, and physical measures were collected by Biobank staff. Demographic information including participant sex, age (in years), age completing full-time education (in years), employment status, ethnicity and Index of Multiple Deprivation (IMD) scores were recorded. The initial recruitment phase occurred between 2006 and 2010, and additional follow-up assessments were completed in-person and online. For an overview of the assessment timeline, see Fig. 2.

UK Biobank data collection and access were conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by the Northwest Multi-centre Research Ethics Committee (MREC). Use of data for secondary data analysis was approved by the UK Biobank and received departmental ethical approval from Swansea University’s School of Psychology Research Ethics Committee (no. 1515). The data




Fig. 1. Overview of proposed mediated moderation model predicting body weight outcomes (Y), including dietary variety (X), cumulative portion size (M), and energy density (W) in direct and mediated pathways. DVS, dietary variety scores.




Fig. 2. Overview of the data collection timeline for the UK Biobank, including measures of interest.

analysis plan was preregistered on the Open Science Framework (OSF) prior to conducting main analyses (<https://osf.io/hfrefj/>).

Participant eligibility

Participants were eligible for this study if they had both dietary recall questionnaire data and BMI recorded at baseline. Participants were included if they had self-reported experiencing no major dietary changes in the preceding 5 years, had no recent experiences of poor appetite or overeating, were not currently pregnant, had not been previously or currently diagnosed with cancer, and had not reported any ongoing behavioural addictions or substance dependencies. Participants were excluded if they self-reported having a current or history of professionally diagnosed eating disorders. Participants were also excluded if credibility of dietary intake was flagged by the Biobank (> 20 MJ/4780 kcal for males, > 18 MJ/4302 kcal for females), or if daily energy intake may otherwise be 'implausible' (< 2 MJ/500 kcal)^(44,45). Where information relevant to eligibility criteria and data credibility had been reassessed during follow-up visits, participant data were removed from additional time points if eligibility criteria were no longer met. For full code and data cleaning steps to identify eligible participants, please see the OSF: <https://osf.io/ktqd6>.

Measures

24-h dietary recall questionnaires. The Oxford WebQ 24-h dietary recall questionnaire⁽⁴⁶⁾ was first administered to participants at baseline, before participants were invited to complete up to four follow-up assessments via email⁽⁴⁷⁾. Participants were asked to report their consumption of > 200 individual foods and

beverages during the previous day. For each item, participants were asked to estimate the number of servings consumed from multiple-choice lists. Where possible, servings related to standard units (e.g. 'a slice of bread'). For an item to be reported, participants consumed ≥ 0.25 serving (self-assessed).

Participants provided information about their general diet and physical activity during the recall period using multiple-choice lists. Participants self-reported dietary restrictions (relating to eggs, dairy products, wheat and sugar), and whether or not they followed any type of special diet (including gluten-free, lactose-free, low-calorie, vegetarian and vegan). Participants self-reported time spent doing vigorous (e.g. fast running/cycling), moderate (e.g. brisk walking/jogging) and light exercise (e.g. leisurely walking) for the recall period. As a data check at the end of the survey, participants were asked 'Would you say that what you ate and drank yesterday was fairly typical for you?'

Dietary variety scores. For each 24-h dietary recall questionnaire, an overall DVS was calculated by summing the number of individual foods and energy-containing beverages consumed. Condiments, breakfast food condiments and beverage condiments (see Table 1) were excluded from this count to avoid repetition of items (e.g. oil for cooking and sugar in tea/coffee). Scores had a possible range between 1 and 203 items (reflecting the number of individual coded items), with higher scores indicating greater variety.

Additional scores were calculated to assess dietary variety between food groups (DVS-B) and dietary variety within food groups (DVS-W). Individual items were categorised into ten food groups⁽²⁹⁾ (see Table 1). To calculate DVS-B, participants were assigned a score from 1 to 10 based on the number of food

Table 1. Foods and beverages included in the Oxford WebQ 24-h dietary recall questionnaire, with assignment to ten food groups*

Food group no.	Food group name	Items included
1	Breakfast foods	Cereals (including porridge, muesli, whole-wheat cereals and sweetened cereals); whole eggs, omelette, other eggs; sausage, bacon
2	Lunch and dinner entrees	Pizza; instant, canned and homemade soups; white and wholegrain pasta; sandwiches with eggs; beef, pork, lamb, poultry, ham, liver, other meat intake; crumbed or deep-fried poultry, breaded fish, battered fish; sushi, tinned tuna, oily fish, white fish, prawns, lobster/crab, other fish; vegetarian sausages/burgers, tofu, Quorn, other vegetarian alternatives
3	Sweets, snacks and carbohydrates	Sliced bread, baguettes, baps, bread rolls, naan bread, garlic bread, crisp bread, oatcakes, other bread; white and brown rice, Couscous, other grains; boiled/baked potatoes, fried potatoes, mashed potatoes, sweet potatoes; Yorkshire pudding, Indian snacks; pastry, crumble, pancakes, croissants, scones, cakes, doughnuts, sponge pudding; yogurt, ice cream, milk-based pudding, other milk-based pudding, cheesecake; soya dessert, other dessert; chocolate, sweets, diet sweets, sweet biscuits, cereal bars, other sweets; salted nuts, unsalted nuts, seeds, olives, crisps, savoury biscuits, scotch egg, other savoury snacks, snack pot
4	Condiments	Butter/margarine on breads, butter/margarine added to potatoes; added salt to food, spreads/sauces consumed, use of cooking fat
5	Fruits	Stewed fruit, prune, dried fruit; berry, cherry, grape; grapefruit, mango, melon, orange, satsuma, peach/nectarine, plum, pineapple; apple, banana, pear; mixed fruit, other fruit
6	Vegetables	Baked beans, pulses, broad beans, green beans, peas; coleslaw, cabbage/kale, turnip/swede, butternut squash, parsnip; side salad, watercress, cucumber, celery, lettuce; garlic, onion; avocado, beetroot, broccoli, carrot, cauliflower, courgette, leek, mushroom, sweet pepper, spinach, sprouts, sweetcorn, fresh and tinned tomatoes; mixed vegetables, vegetable pieces, other vegetables
7	Energy-containing beverages	Low-calorie drinks, fizzy drinks, squash; fruit and vegetable juices, smoothies; coffees, teas, decaffeinated coffee, decaffeinated tea, hot chocolate; milk, flavoured milk; other non-alcoholic drinks; beer, cider, wine, fortified wine, and spirits
8	Other dairy products	Low-fat cheeses, hard cheese, soft cheese, blue cheese, mozzarella, goat's cheese, feta; cheese spreads, cottage cheese, and other cheeses
9	Breakfast food condiments	Dried fruit added to cereals, milk added to cereals, sugar added to cereals, artificial sweetener added to cereals
10	Beverage condiments	Added milk, added sugar, added artificial sweetener

* In line with the approach of McCrory *et al.*⁽²⁹⁾, individual foods and beverages were categorised into ten food groups.

groups from which they had consumed at least one item, with higher scores indicating consumption from across a greater variety of food groups. To calculate DVS-W, the percentage of individual items consumed from within each defined food group was determined (excluding condiment groups due to limited data for serving size). For each food group, a higher percentage indicated a greater variety of items consumed.

Portion size and energy density. Overall intake values were available in the dataset. Total weight of foods/beverages consumed (g) was used to indicate cumulative portion size, and energy density (kJ/g) across items was calculated. To estimate intake of individual items, standard portion sizes were used to calculate weight consumed⁽⁴⁸⁾, and standard food composition tables were used to calculate energy intake⁽⁴⁹⁾. Total intake was then derived for each individual food group. Values > 16 kJ/g (4 kcal/g) indicate 'high' energy density (see ref. 50, as reported in ref. 51).

Anthropometric outcome measures. Anthropometric measurements were collected from participants during assessment centre visits (for further information, see ref. 52). Standing height and waist circumference were measured using a Seca 240 cm height measure and Seca 200 cm tape measure, respectively. Body weight (kg), whole-body fat (%) and fat-free mass (kg) were collected using a Tanita BC418MA body composition analyser or manually recorded using standard weighing scales

where required. Height and weight measurements were used to calculate BMI (kg/m²).

Data analyses

At baseline, 35 449 participants were eligible for inclusion in the study, had completed the 24-h dietary recall questionnaire and had BMI recorded. Of these participants, 34 974 and 34 992 had body fatness and fat-free mass measured, respectively. Initial data cleaning was conducted in Python; this included identifying eligible participants for the study, removing unnecessary columns from the dataset, creating variable scores for multiple time points (including energy density and DVS) and reshaping data for analyses. Moderated mediation models were conducted using PROCESS v3.5.3⁽⁵³⁾, and remaining data analyses were conducted in Stata 16.0⁽⁵⁴⁾. For access to data analysis logs, see the OSF (<https://osf.io/hfrej/>).

A series of bivariate correlation analyses (Spearman's) were used to identify associations between potential covariates and outcome variables (online Supplementary Tables A.1 and A.2). Where relevant, categorical covariates were dummy-coded to produce binary variables, and all significant covariates were entered at baseline (T0). If covariates were significant but had missing data, models were repeated with these variables included as part of sensitivity analyses to check for differences in effects (online Supplementary Tables A.3 and A.4).

To test hypothesised relationships, a series of second-stage moderated mediation models were conducted at baseline (T0). DVS were included as the predictor (X), cumulative portion size was entered as the mediator (M) and overall energy density was entered as the moderator (W) of both the direct and mediated pathways. BMI, body fat percentage and fat-free mass were included as the outcome (Y). Participant sex was included as a significant covariate across models ($r_{pb} = 0.185\text{--}0.842$), and effects of controlling for age completing full-time education ($n = 21\,068$, $r_s = -0.100$) were explored in sensitivity analyses due to missing data. Across models, predictors were mean-centred, CI were adjusted using bias-corrected bootstrap approaches (5000 samples) and HC3 was used to adjust standard errors for heteroscedasticity⁽⁵⁵⁾.

To explore longitudinal changes in body weight outcomes, a series of Wilcoxon signed-rank tests were used to check for differences in dietary intake and body weight outcomes between time points. Second-stage moderated mediation models conducted at baseline were repeated with DVS entered at T0, and composite scores for cumulative portion size and energy density entered at T1 (averaged across completed follow-up questionnaires). BMI was included as the outcome at T2 and T3, and body fat percentage and fat-free mass at T2. As < 50 participants had both dietary data and BMI collected at T4, this time point was dropped from analyses. Participant sex ($r_{pb} = 0.177\text{--}0.848$), age ($r_s = 0.143$), IMD score for England ($r_s = 0.105$) and time spent doing vigorous physical activity ($r_{pb} = -0.158$) were included as significant covariates. Baseline measures of BMI, body fat percentage and fat-free mass were controlled in the appropriate model. Effects of controlling for age completing full-time education ($r_s = -0.160$) were explored in sensitivity analyses.

Unless otherwise stated, effects were significant if CI did not cross zero, and $P < 0.05$. However, given the large sample size and increased risk of finding significant but 'trivial' effects across multiple models^(56,57), overall model and coefficient effects were considered meaningful if equivalent to a 'small effect size' (observed $R^2 \geq 0.02$; $r \geq 0.10$)⁽⁵⁸⁾. To align with reported longitudinal trends in weight gain and development of overweight⁽⁵⁹⁾, change in dietary intake was identified as significant if it indicated ≥ 13 kcal increase in daily food consumption per unit (≈ 2.7 g)⁽⁴⁹⁾. Change in BMI was identified as significant if it increased by at least 0.10 kg/m²⁽⁶⁰⁻⁶²⁾, and body fat if it increased by at least 0.3% ⁽⁶²⁾.

As part of exploratory sensitivity checks (online Supplementary Tables A.5 and A.6), estimated values for energy intake and weight of food consumed within food groups were used to calculate energy density after removing estimated beverage scores. Overall energy density scores (kJ/g) were lower than estimated energy density scores (without beverages) at baseline (mean difference = -3.38 , $SD = 1.81$) and follow-up (mean difference = -3.12 , $SD = 1.33$). However, mean scores across measures were < 6.6 kJ/g, indicating generally 'low' energy density⁽⁵¹⁾. See Table 2.

Results

Cross-Sectional analyses at baseline

Participant characteristics. Table 3 displays sample characteristics. Of those included in the sample, 52.3% were female. Most

participants identified themselves as British (87.9%) or Irish (2.7%), or other White background (3.8%), and were either in paid employment (59.6%) or retired (32.7%). Specific dietary restrictions were reported by 16.7% of participants, 13.6% followed a special diet and 39.1% were users of vitamin supplements. Most participants reported doing 0–60 min of vigorous (92.5%) or moderate (83.7%) physical activity, and 0–5 h of light physical activity (93.1%). 83.2% reported that recall of their diet was representative of their typical eating habits. See Table 2 for dietary intake.

Dietary variety scores as predictors of body weight outcomes.

After controlling for effects of participant sex, models conducted at baseline (Fig. 3) showed that contrary to predictions, higher DVS was a significant and negative predictor of BMI and fat-free mass. Though greater dietary variety also significantly predicted increased daily food intake and the consumption of larger portions overall, cumulative portion size did not directly predict BMI, body fatness or fat-free mass. Energy density did not significantly moderate the influence of DVS – or portion size – on BMI, body fatness or fat-free mass. Moderated mediation effects of DVS on BMI (Index = 0.007, bootstrap $se = 0.003$, bootstrap 95% CI = 0.001, 0.013) and fat-free mass (Index = 0.035, bootstrap $se = 0.005$, bootstrap 95% CI = 0.026, 0.044) were significant but relatively small, as the conditional indirect effect of DVS increased at higher levels of energy density for both BMI ($-1SD$ Effect = 0.037 *v.* $+1SD$ Effect = 0.048, bootstrap $se = 0.005$, bootstrap 95% CI = 0.002, 0.021) and fat-free mass ($-1SD$ Effect = 0.098 *v.* $+1SD$ Effect = 0.156, bootstrap $se = 0.008$, bootstrap 95% CI = 0.042, 0.073). The moderated mediation effect of DVS on body fatness was not supported (Index < 0.001 , bootstrap $se = 0.004$, bootstrap 95% CI = -0.009 , 0.009).

A similar pattern of results was observed when DVS-B and DVS-W were included as predictors (online Supplementary Table A.7), and when 'age completing full-time education' was included as a covariate (online Supplementary Table A.3), with little evidence to support associated pathways between dietary variety and body weight outcomes.

Longitudinal analyses at follow-up

Data availability and participant characteristics. Of 17 272 participants who had completed at least one 24-h dietary recall questionnaire at follow-up (T1), up to 738 participants had body weight outcomes recorded at T2 (46.7% female) and 1563 at T3 (47.3% female).

Changes in dietary intake and body weight outcomes across time points were small. Compared with baseline, participants were consuming a smaller cumulative portion size at T1 (T0 $M = 3.3$ kg, $SD = 0.8$ *v.* T1 $M = 3.2$ kg, $SD = 0.7$; $Z = 5.690$, $P < 0.001$, $r = 0.13$), but energy density was significantly higher overall (T0 $M = 2.8$ kJ/g, $SD = 0.8$ *v.* T1 $M = 2.9$ kJ/g, $SD = 0.7$; $Z = -4.660$, $P < 0.001$, $r = 0.10$) (see Table 2 for dietary intake at follow-up). BMI (T0 $M = 25.7$ kg/m², $SD = 3.9$ *v.* T2 $M = 26.0$ kg/m², $SD = 4.1$; $Z = -3.639$, $P < 0.001$, $r = 0.13$) and body fatness (T0 $M = 28.6\%$, $SD = 8.0$ *v.* T2 $M = 29.5\%$, $SD = 8.2$; $Z = -5.175$, $P < 0.001$, $r = 0.19$) had increased at T2, whilst fat-free mass



Table 2. Dietary intake at baseline (T0) and follow-up (T1)

Variable	T0									T1								
	Female (n 18 551)			Male (n 16 898)			Overall (n 35 449)			Female (n 940)			Male (n 1056)			Overall (n 1996)		
	M	SD	Range	M	SD	Range	M	SD	Range	M	SD	Range	M	SD	Range	M	SD	Range
Energy intake (MJ/d)	8.1	2.5	2.1–18.0	9.6	3.0	2.1–20.0	8.8	2.8	2.1–20.0	8.3	1.8	3.1–15.7	9.4	2.3	3.8–17.4	8.9	2.1	3.1–17.4
Food weight consumed (kg/d)	3.1	0.8	0.3–9.8	3.4	1.0	0.6–9.7	3.2	0.9	0.3–9.8	3.1	0.6	1.4–6.1	3.3	0.8	1.4–6.4	3.2	0.7	1.4–6.4
Energy density with beverages (kJ/g)*	2.7 (0.6)	0.8	0.7–9.4 (0.2–2.2)	2.9 (0.7)	0.8	0.8–9.5 (0.2–2.3)	2.8 (0.7)	0.8	0.7–9.5 (0.2–2.3)	2.8 (0.7)	0.6	1.2–5.4 (0.3–1.3)	3.0 (0.7)	0.7	1.3–6.3 (0.3–1.5)	2.9 (0.7)	0.7	1.2–6.3 (0.3–1.5)
Energy density without beverages (kJ/g)*	5.8 (1.4)	2.0	1.1–21.3 (0.3–5.1)	6.6 (1.6)	2.1	1.7–22.8 (0.4–5.4)	6.2 (1.5)	2.1	1.1–22.8 (0.3–5.4)	5.7 (1.4)	1.4	2.6–13.1 (0.6–3.1)	6.1 (1.5)	1.4	2.5–13.5 (0.6–3.2)	5.9 (1.4)	1.4	2.5–13.5 (0.6–3.2)
Protein (g/d)	76.3	25.4	7.8–246.7	85.5	30.2	4.6–364.1	80.7	28.2	4.6–364.1	77.4	19.2	25.8–155.1	84.6	22.1	21.3–189.3	81.2	21.1	21.3–189.3
Carbohydrate (g/d)	239.9	83.1	18.0–702.7	273.7	95.6	6.5–783.4	256.0	90.9	6.5–783.4	240.0	63.4	70.0–524.7	268.6	73.3	80.0–578.4	255.2	70.3	70.0–578.4
Total fat (g/d)	71.8	30.7	2.3–248.2	82.9	35.8	1.5–268.2	77.1	33.7	1.5–268.2	74.2	23.8	16.7–159.9	82.1	28.6	17.9–226.9	78.4	26.7	16.7–226.9
Saturated fat (g/d)	27.5	12.9	0.3–103.9	32.2	15.3	0.7–116.3	29.7	14.3	0.3–116.3	28.4	10.5	4.8–73.4	31.5	12.4	5.6–95.5	30.0	11.6	4.8–95.5
Total sugars (g/d)	116.7	51.2	3.6–522.4	126.5	56.0	0.4–548.3	121.4	53.7	0.4–548.3	115.6	39.8	25.3–342.9	124.7	43.6	29.7–294.3	120.4	42.1	25.3–342.9
DVS†	16.9	5.3	2.0–46.0	16.1	5.4	2.0–53.0	16.5	5.4	2.0–53.0	17.8	4.4	7.5–36.0	17.1±4.7	4.7	4.0–36.0	17.5	4.5	4.0–36.0
DVS-B‡	7.9	1.4	1.0–10.0	7.7	1.5	1.0–10.0	7.8	1.4	1.0–10.0	8.1	1.1	3.0–10.0	7.9	1.2	4.0–10.0	8.0	1.1	3.0–10.0
DVS-W (§)																		
Breakfast foods	7.9	6.0	0.0–53.8	8.6	6.6	0.0–46.2	8.2	6.3	0.0–53.8	8.5	4.9	0.0–26.9	9.2±5.6	5.6	0.0–34.6	8.9	5.3	0.0–34.6
Lunch and dinner entrees	4.9	3.1	0.0–33.3	4.9	3.1	0.0–33.3	4.9	3.1	0.0–33.3	5.2	2.3	0.0–16.7	5.1	2.5	0.0–20.0	5.2	2.4	0.0–20.0
Sweets, snacks and carbohydrates	6.6	2.8	0.0–24.2	6.7	3.0	0.0–25.8	6.7	2.9	0.0–25.8	7.0	2.3	0.8–14.5	7.0	2.5	0.0–16.1	7.0	2.4	0.0–16.1
Condiments	25.7	22.3	0.0–100.0	23.6	22.7	0.0–100.0	24.7	22.5	0.0–100.0	27.8	17.6	0.0–100.0	25.7	19.1	0.0–100.0	26.7	18.4	0.0–100.0
Fruits	11.6	8.6	0.0–73.7	9.4	8.0	0.0–73.7	10.5	8.4	0.0–73.7	11.6	7.5	0.0–47.4	10.2	7.4	0.0–59.6	10.9	7.5	0.0–59.6
Vegetables	10.6	8.8	0.0–62.9	8.6	8.2	0.0–68.6	9.6	8.6	0.0–68.6	10.8	6.7	0.0–40.0	9.5	6.7	0.0–42.9	10.1	6.7	0.0–42.9
Energy-containing beverages	10.6	4.7	0.0–37.5	11.3	4.8	0.0–43.8	10.9	4.8	0.0–43.8	11.5	4.3	0.0–28.1	11.8	4.1	0.0–28.1	11.7	4.2	0.0–28.1
Other dairy products	4.9	5.8	0.0–54.5	4.7	5.9	0.0–90.9	4.8	5.8	0.0–90.9	5.0	4.9	0.0–27.3	4.7	4.6	0.0–27.3	4.9	4.7	0.0–27.3
Breakfast food condiments	25.1	22.6	0.0–100.0	26.0	23.3	0.0–100.0	25.5	22.9	0.0–100.0	26.3	19.8	0.0–75.0	27.8	20.2	0.0–75.0	27.1	20.0	0.0–75.0
Beverage condiments	15.6	10.5	0.0–60.0	17.8	11.6	0.0–70.0	16.7	11.1	0.0–70.0	15.2	9.3	0.0–50.0	16.6	10.7	0.0–50.0	16.0	10.1	0.0–50.0

* Energy density displayed as kcal/g in brackets for the mean and range.

† Dietary variety score calculated as the count of individual foods and beverages consumed across food groups.

‡ Dietary variety score calculated as the count of food groups from which items were consumed, scored from 1 to 10.

§ Dietary variety score calculated as the percentage of items consumed from within each food group.

Table 3. Overview of sample characteristics at baseline (T0)*

Demographic	Female (n 18 551)			Male (n 16 898)			Overall (n 35 449)		
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range
Age (years)	55.9	8.2	40.0–70.0	56.6	8.3	40.0–70.0	56.2	8.3	40.0–70.0
Age completing full-time education (years)	17.0	2.1	0.0–35.0	17.1	2.5	0.0–35.0	17.0	2.3	0.0–35.0
Index of Multiple Deprivation (IMD)†,‡									
England (n 34 338)	15.4	11.5	1.44–79.3	15.9	12.1	1.43–82.0	15.7	11.8	1.43–82.0
Wales (n 299)	12.3	10.2	2.6–61.6	13.2	11.4	2.6–57.7	12.7	10.8	2.6–61.6
BMI (kg/m ²)	26.0	4.7	15.2–59.5	27.2	3.9	14.9–63.4	26.5	4.4	14.9–63.4
Whole-body fat (%)	35.4	6.8	9.4–63.1	24.5	5.7	5.0–52.6	30.2	8.3	5.0–63.1
Fat-free mass (kg)	44.0	4.7	29.5–78.0	63.4	7.5	35.7–100.0	53.2	11.5	29.5–100.0
Waist circumference (cm)	82.1	11.7	48.0–157.0	95.2	10.8	63.0–163.0	88.3	13.1	48.0–163.0

* Mean ± SD (range).

† Higher scores indicate residential areas have greater levels of deprivation.

‡ Statistics not reported for Scotland as IMD score was available for < 50 participants.




Fig. 3. Baseline models of overall dietary variety (DVS) as a predictor of BMI (R² = 0.035), body fatness (BF) (R² = 0.434) and fat-free mass (FFM) (R² = 0.718). Unstandardised regression coefficients (b), adjusted standard error in brackets (se) and bootstrap CI are displayed (LLCI and ULCI). Pathways including the moderator are indicated with a dashed line. Significant coefficients are indicated in bold (*P* < 0.001). DVS, dietary variety scores.

had decreased (T0 M = 53.6 kg, SD = 11.1 v. T2 M = 52.8 kg, SD = 10.9; Z = 7.165, *P* < 0.001, *r* = 0.27). All other differences between time points for dietary intake and body weight outcomes were either not significant or trivial effects.

Time-Lagged associations between dietary variety and weight-related outcomes. After controlling for effects of participant sex, age, IMD score for England and time spent doing vigorous physical activity, time-lagged moderated mediation models (Fig. 4) showed similar results as baseline models. DVS (T0) remained a significant and positive predictor of cumulative portion size (T1) but was no longer directly associated with body weight outcomes (T2 and T3). Portion

size did not significantly predict BMI (T2 and T3), body fatness (T2) or fat-free mass (T2). Energy density (T1) did not significantly interact with portion size or DVS to predict body weight outcomes at follow-up, and the moderated mediation effects of DVS on BMI (T2 Index < 0.001, bootstrap se = 0.004, bootstrap 95 % CI = -0.007, .008; T3 Index < 0.001, bootstrap se = 0.003, bootstrap 95 % CI = -0.006, .006), body fatness (Index = 0.007, bootstrap se = 0.010, bootstrap 95 % CI = -0.012, .027) and fat-free mass (Index = -0.001, bootstrap se = 0.007, bootstrap 95 % CI = -0.014, .012) were not supported.

A similar pattern of results was found when DVS-B and DVS-W were included as the predictor at T0 (online




Fig. 4. Time-lagged models of overall dietary variety (DVS) as a predictor of outcomes at follow-up, including BMI (T2 R2 = 0.926; T3 R2 = 0.872), body fatness (BF) (R2 = 0.899) and fat-free mass (FFM) (R2 = 0.971). Unstandardised regression coefficients (b), adjusted standard error in brackets (se) and Bootstrap CI are displayed (LLCI and ULCI). Pathways including the moderator are indicated with a dashed line. Significant coefficients are indicated in bold ($P < 0.001$). All models included baseline measures of body weight outcomes as predictors. DVS, dietary variety scores.

Supplementary Table A.8) and when ‘age completing full-time education’ was included as a covariate (online Supplementary Table A.4).

Discussion

This study sought to further explore the relationship between dietary variety and body weight outcomes, whilst considering the intermediate roles of cumulative portion size and energy density. First, contrary to predictions, DVS was directly associated with a lower BMI and fat-free mass at baseline, providing further population-level evidence to support a potentially beneficial role of variety in the consumer diet. In this study sample, energy density overall was noticeably low⁽⁵¹⁾. Though there was little evidence to support significant associations between DVS within food groups and body weight outcomes in supplementary analyses (with the exception of the largest group containing ‘sweets, snacks, and carbohydrates’), participants were consuming foods from across a wide range of food groups (including fruits and vegetables). This appears to reflect prior research that has associated dietary variety with higher diet quality^(8–10,27). Indeed, evidence suggests that moderating variety appropriately across foods can be helpful to consumers when utilised in dietary interventions^(33,34,39).

Second, as predicted, consuming a greater variety of foods (including scores across and within food groups) was

significantly associated with increased weight of food consumed. In an important step towards triangulation⁽⁶³⁾, these results further corroborate findings from short-term experimental studies conducted in a mealtime context⁽³⁷⁾, supporting variety as a factor that may be significantly associated with food intake at a broader level. Past studies exploring relationships between dietary variety and consumption have typically focused on cross-sectional effects at a single time point, in relation to energy intake⁽¹¹⁾. By adopting a time-lagged modelling approach, this study specifically highlights consistent effects of variety on weight of food consumed over a longer period of time, outside of a single mealtime context, and across several potential eating sessions. Results also highlight a level of consistency between dietary variety scores in relation to food intake, as scores based on the overall count of individual items, number of food groups from which items were consumed and percentage of items consumed within food groups were all significantly associated with cumulative portion size.

However, energy density scores had less variability across participants in this study, potentially accounting for the non-significant moderated mediation effect of DVS on body weight outcomes (via cumulative portion size). One explanation is that dietary data may have been susceptible to ‘underreporting’, though data credibility was checked to reduce this bias. Previous research also suggests that including beverages within calculations of energy density can reduce overall estimates^(64–66). Though sensitivity analyses indicated that energy density

calculated from estimates of energy intake and weight consumed – when all beverages were excluded – was still low across participants, justification for removing beverage intake from analyses may be of interest particularly when energy density is high for foods alone. It is also important to acknowledge that cumulative portion size within a single recall period may not be expected to significantly influence body weight stability, particularly if energy density is low⁽⁵¹⁾. Indeed, research suggests that cumulative dietary recall measures, that are collected over several days or weeks, may provide more comprehensive self-report data⁽⁶⁷⁾.

Some additional limitations of data analyses in this study should be addressed. Missing data across time points meant that models were cross-sectional at baseline, and only a single time point was included for dietary data at follow-up (averaged across available reports). Dietary data were also derived from self-report 24-h recall questionnaires. Though this specific measure has been validated within the Biobank^(46,68), there was a lack of specificity when reporting serving sizes (as this was defined by multiple-choice lists). Participants were instructed to use an ingredient-based approach to reporting food intake, whereby mixed meals/dishes were disaggregated into component parts (e.g. 'Spaghetti bolognese' was reported as 'pasta', 'beef' and 'tomato-based sauce'⁽⁴⁷⁾). Recent evidence has highlighted that this can improve sensitivity when estimating nutritional intake^(69,70), and consideration of multiple components included within mixed dishes/foods may be particularly important when considering effects of 'sensory' variety on food intake^(28,71). However, information about specific flavours and brands may be relevant to calculations of dietary variety when measuring food intake across several days⁽⁶⁷⁾.

Food groups were also constructed in line with previous food categories used for FFQ data⁽²⁹⁾, meaning the number of items included to indicate variety differed between groups. DVS-W was included as an exploratory measure in addition to overall DVS (and energy density) in models, as previous studies have reported negative *v.* positive associations with body adiposity when scoring variety separately for groups of 'healthful' foods (typically fruits, vegetables and low energy density foods) and 'energy-dense' foods, respectively (for a review, see ref. 11). As McCrory and colleagues highlighted a key difference in the direction of associations between vegetables and other food groups specifically⁽²⁹⁾, this food group approach was adopted in the current study. However, 'sweets, snacks, and carbohydrates' included a comparatively greater number of items than others, potentially allowing for more scope to indicate 'variety' within a limited recall period. It is also acknowledged that the categories used for the purpose of this study were generally broad and collapsed across foods that may be further distinguished based on nutritional content, particularly where diet quality specifically is of interest (for a more recent example with UK Biobank dietary data, see ref. 72).

Considering the findings of this study, two key issues are acknowledged. First, it is important to consider the possibility of a bidirectional relationship between dietary variety and body weight outcomes, whereby having a lower body weight (and body fatness) can predict healthier eating patterns.

Bidirectionality has previously been discussed in relation to other predictors of body weight outcomes and food intake, such as weight stigma⁽⁷³⁾ and social effects on eating⁽⁷⁴⁾. Though this was not directly explored within models presented here, results do support relevant negative associations between intake of dietary variety for low energy density food groups and body weight outcomes. In light of these associations, future research should explore the direction of possible 'causality', whereby potential reciprocal predictive pathways between body weight and dietary variety are observed in cross-panel, time-lagged models of effects over a longer period with more frequent time points (e.g. a similar approach has been adopted to explore longitudinal associations between BMI and internalising symptoms in a cohort study⁽⁷⁵⁾).

Second, results of this study further highlight potential sample bias. Though educational attainment generally aligns with trends in the UK population for this age group⁽⁷⁶⁾, and we note the variability of the eligible sample for BMI (including participants with potential underweight and obesity), UK Biobank participants in particular have been found to be more likely to live in areas with less deprivation, have a lower average BMI, be less likely to have obesity and have lower incidence of self-reported health conditions, such as CVD, diabetes and cancers⁽⁷⁷⁾. Therefore, it is important to acknowledge potential limitations of generalisability for such population samples that may be susceptible to 'healthy' volunteer bias (particularly when using a subset of the sample).

Conclusions

Contrary to predictions, findings of this study highlight a direct association between overall dietary variety and body weight, that indicates lower BMI and body adiposity in relation to greater variety, and may have particular implications for assisting body weight management and diet quality. Though there was little evidence to support the predicted moderated mediation effect of dietary variety on body weight outcomes when considering the role of cumulative portion size and energy density in this relationship, this study also further supports dietary variety as a factor associated with increased food intake. Further research is needed to explore the influence of these associations over a longer period, particularly as this relates to variability in energy density across the consumer diet.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 53159.

This work was supported by the ESRC Wales Doctoral Training Partnership (Project Reference: ES/P00069X/1, Studentship 1 947 139). For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

R. E.: Conceptualisation, Methodology, Formal analysis, Writing – original draft, Writing – review and editing, and Project administration. M. P.: Conceptualisation, Methodology,

and Writing – review and editing. M. D. L.: Conceptualisation, Methodology, and Writing – review and editing. A. J.: Conceptualisation, Methodology, Formal analysis, and Writing – review and editing. L. L. W.: Conceptualisation, Methodology, Writing – original draft, Writing – review and editing, Supervision, and Project administration.

The authors declare that they have no competing interests.

Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114523000156>

References

- Verghese A, Raber M & Sharma S (2019) Interventions targeting diet quality of Supplemental Nutrition Assistance Program (SNAP) participants: a scoping review. *Prev Med* **119**, 77–86.
- Burden S, Jones DJ, Sremanakova J, *et al.* (2019) Dietary interventions for adult cancer survivors. *Cochrane Database Syst Rev* **2019**, CD011287.
- Xu Z, Steffen LM, Selvin E, *et al.* (2020) Diet quality, change in diet quality and risk of incident CVD and diabetes. *Public Health Nutr* **23**, 329–338.
- Conklin AI, Monsivais P, Khaw K-T, *et al.* (2016) Dietary diversity, diet cost, and incidence of type 2 diabetes in the united kingdom: a prospective cohort study. *PLoS Med* **13**, e1002085.
- De Koning L, Chiuve SE, Fung TT, *et al.* (2011) Diet-quality scores and the risk of type 2 diabetes in men. *Diabetes Care* **34**, 1150–1156.
- Park SY, Boushey CJ, Wilkens LR, *et al.* (2017) High-quality diets associate with reduced risk of colorectal cancer: analyses of diet quality indexes in the multiethnic cohort. *Gastroenterology* **153**, 386–394.e2.
- Asghari G, Mirmiran P, Yuzbashian E, *et al.* (2017) A systematic review of diet quality indices in relation to obesity. *Br J Nutr* **117**, 1055–1065.
- Drewnowski A, Henderson SA, Driscoll A, *et al.* (1997) The dietary variety score: assessing diet quality in healthy young and older adults. *J Am Diet Assoc* **97**, 266–271.
- Nair MK, Augustine LF & Konapur A (2016) Food-based interventions to modify diet quality and diversity to address multiple micronutrient deficiency. *Front Public Heal* **3**, 277.
- Meng L, Wang Y, Li T, *et al.* (2018) Dietary diversity and food variety in chinese children aged 3–17 years: are they negatively associated with dietary micronutrient inadequacy? *Nutrients* **10**, 1674.
- Vadiveloo MK, Dixon LB & Parekh N (2013) Associations between dietary variety and measures of body adiposity: a systematic review of epidemiological studies. *Br J Nutr* **109**, 1557–1572.
- Gregory CO, McCullough ML, Ramirez-Zea M, *et al.* (2009) Diet scores and cardio-metabolic risk factors among Guatemalan young adults. *Br J Nutr* **101**, 1805.
- Azadbakht L & Esmailzadeh A (2011) Dietary diversity score is related to obesity and abdominal adiposity among Iranian female youth. *Public Health Nutr* **14**, 62–69.
- Heidari-Beni M, Riahi R, Massoudi S, *et al.* (2021) Association between dietary diversity score and anthropometric indices among children and adolescents: the weight disorders survey in the CASPIAN-IV study. *J Sci Food Agric* **101**, 5075–5081.
- Zhang Q, Chen X, Liu Z, *et al.* (2017) Diet diversity and nutritional status among adults in southwest China. *PLoS One* **12**, e0172406.
- Vadiveloo MK, Dixon LB, Mijanovich T, *et al.* (2015) Dietary variety is inversely associated with body adiposity among US adults using a novel food diversity index. *J Nutr* **145**, 555–563.
- Moraes L, Lindroos AK, Warensjö Lemming E, *et al.* (2020) Diet diversity score and healthy eating index in relation to diet quality and socio-demographic factors: results from a cross-sectional national dietary survey of Swedish adolescents. *Public Health Nutr* **23**, 1754–1765.
- Bernstein MA, Tucker KL, Ryan ND, *et al.* (2002) Higher dietary variety is associated with better nutritional status in frail elderly people. *J Am Diet Assoc* **102**, 1096–1104.
- Roberts SB, Hajduk CL, Howarth NC, *et al.* (2005) Dietary variety predicts low body mass index and inadequate macronutrient and micronutrient intakes in community-dwelling older adults. *J Gerontol Ser A Biol Sci Med Sci* **60**, 613–621.
- Royo-Bordonada MA, Gorgojo L, Ortega H, *et al.* (2003) Greater dietary variety is associated with better biochemical nutritional status in Spanish children: the Four Provinces Study. *Nutr Metab Cardiovasc Dis* **13**, 357–364.
- Marshall TA, Stumbo PJ, Warren JJ, *et al.* (2001) Inadequate nutrient intakes are common and are associated with low diet variety in rural, community-dwelling elderly. *J Nutr* **131**, 2192–2196.
- Popkin BM, Adair LS & Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* **70**, 3–21.
- Kearney J (2010) Food consumption trends and drivers. *Philos Trans R Soc B Biol Sci* **365**, 2793–2807.
- Johnson F & Wardle J (2014) Variety, palatability, and obesity. *Adv Nutr* **5**, 851–859.
- De Oliveira Otto MC, Padhye NS, Bertoni AG, *et al.* (2015) Everything in moderation – dietary diversity and quality, central obesity and risk of diabetes. *PLoS One* **10**, e0141341.
- McCrorry MA, Burke A & Roberts SB (2012) Dietary (sensory) variety and energy balance. *Physiol Behav* **107**, 576–583.
- Kennedy E (2004) Dietary diversity, diet quality, and body weight regulation. *Nutr Rev* **62**, S78–S81.
- Raynor HA & Vadiveloo MK (2018) Understanding the relationship between food variety, food intake, and energy balance. *Curr Obes Rep* **7**, 68–75.
- McCrorry MA, Fuss PJ, McCallum JE, *et al.* (1999) Dietary variety within food groups: association with energy intake and body fatness in men and women. *Am J Clin Nutr* **69**, 440–447.
- Tande DL, Magel R & Strand BN (2010) Healthy Eating Index and abdominal obesity. *Public Health Nutr* **13**, 208–214.
- Vadiveloo MK, Parkeh N & Mattei J (2015) Greater healthful food variety as measured by the US healthy food diversity index is associated with lower odds of metabolic syndrome and its components in US adults. *J Nutr* **145**, 564–571.
- Vadiveloo MK, Sacks FM, Champagne CM, *et al.* (2016) Greater healthful dietary variety is associated with greater 2-year changes in weight and adiposity in the preventing overweight using novel dietary strategies (POUNDS Lost) trial. *J Nutr* **146**, 1552–1559.
- Epstein LH, Kilanowski C, Paluch RA, *et al.* (2015) Reducing variety enhances effectiveness of family-based treatment for pediatric obesity. *Eat Behav* **17**, 140–143.
- Raynor HA, Steeves EA, Hecht J, *et al.* (2012) Limiting variety in non-nutrient-dense, energy-dense foods during a lifestyle intervention: a randomized controlled trial. *Am J Clin Nutr* **95**, 1305–1314.

35. Raynor HA, Niemeier HM & Wing RR (2006) Effect of limiting snack food variety on long-term sensory-specific satiety and monotony during obesity treatment. *Eat Behav* **7**, 1–14.
36. Haws KL, Liu PJ, Redden JP, *et al.* (2017) Exploring the relationship between varieties of variety and weight loss: when more variety can help people lose weight. *J Mark Res* **54**, 619–635.
37. Embling R, Pink AE, Gatzemeier J, *et al.* (2021) Effect of food variety on intake of a meal: a systematic review and meta-analysis. *Am J Clin Nutr* **113**, 716–741.
38. Epstein LH, Carr KA, Cavanaugh MD, *et al.* (2011) Long-term habituation to food in obese and nonobese women. *Am J Clin Nutr* **94**, 371–376.
39. Epstein LH, Fletcher KD, O'Neill J, *et al.* (2013) Food characteristics, long-term habituation and energy intake. Laboratory and field studies. *Appetite* **60**, 40–50.
40. Wilkinson LL & Brunstrom JM (2016) Sensory specific satiety: more than 'just' habituation? *Appetite* **103**, 221–228.
41. Wilkinson LL, Hinton EC, Fay SH, *et al.* (2013) The 'variety effect' is anticipated in meal planning. *Appetite* **60**, 175–179.
42. Zlatevska N, Dubelaar C & Holden SS (2014) Sizing up the effect of portion size on consumption: a meta-analytic review. *J Mark Res* **78**, 140–154.
43. Allen N, Sudlow C, Downey P, *et al.* (2012) UK Biobank: current status and what it means for epidemiology. *Health Policy Technol* **1**, 123–126.
44. Banna JC, McCrory MA, Fialkowski MK, *et al.* (2017) Examining plausibility of self-reported energy intake data: considerations for method selection. *Front Nutr* **4**, 25.
45. Rhee JJ, Sampson L, Cho E, *et al.* (2015) Comparison of methods to account for implausible reporting of energy intake in epidemiologic studies. *Am J Epidemiol* **181**, 225–233.
46. Liu B, Young H, Crowe FL, *et al.* (2011) Development and evaluation of the Oxford WebQ, a low-cost, web-based method for assessment of previous 24 h dietary intakes in large-scale prospective studies. *Public Health Nutr* **14**, 1998–2005.
47. The UK Biobank (2012) 24-Hour Dietary Recall Questionnaire: Version 1.1. <https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/DietWebQ.pdf> (accessed April 2020).
48. Ministry of Agriculture Fisheries and Food (1993) *Food Portion Sizes*, 3rd ed. London: HMSO.
49. Holland B, Welch AA, Unwin ID, *et al.* (1991) *McCance and Widdowson's The Composition of Foods. McCance and Widdowson's Composition of Foods*. Cambridge: Royal Society of Chemistry.
50. Rolls B (2012) *The Ultimate Volumetrics Diet*. New York, NY: Harper Collins.
51. Rolls BJ (2017) Dietary energy density: applying behavioural science to weight management. *Nutr Bull* **42**, 246–253.
52. The UK Biobank (2014) Anthropometry: Version 1.0. <https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/Anthropometry.pdf> (accessed April 2020).
53. Hayes AF (2018) *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-Based Approach*, 2nd ed. New York, NY: Guildford Press.
54. StataCorp (2019) *Stata Statistical Software: Release 16*. College Station, TX: StataCorp LLC.
55. Hayes AF & Cai LI (2007) Using heteroskedasticity-consistent standard error estimators in OLS regression: an introduction and software implementation. *Behav Res Methods* **39**, 709–722.
56. Lantz B (2013) The large sample size fallacy. *Scand J Caring Sci* **27**, 487–492.
57. Kaplan RM, Chambers DA & Glasgow RE (2014) Big data and large sample size: a cautionary note on the potential for bias. *Clin Transl Sci* **7**, 342–346.
58. Cohen J (1992) Quantitative methods in psychology. A power primer. *Psychol Bull* **112**, 155–159.
59. Katan MB & Ludwig DS (2010) Extra calories cause weight gain— but how much? *JAMA* **303**, 65–66.
60. Rosell M, Appleby P, Spencer E, *et al.* (2006) Weight gain over 5 years in 21 966 meat-eating, fish-eating, vegetarian, and vegan men and women in EPIC-Oxford. *Int J Obes* **30**, 1389–1396.
61. Hughes VA, Frontera WR, Roubenoff R, *et al.* (2002) Longitudinal changes in body composition in older men and women: role of body weight change and physical activity. *Am J Clin Nutr* **76**, 473–481.
62. Guo SS, Zeller C, Chumlea WC, *et al.* (1999) Aging, body composition, and lifestyle: the Fels Longitudinal Study. *Am J Clin Nutr* **70**, 405–411.
63. Munafò MR & Davey Smith G (2018) Robust research needs many lines of evidence. *Nat* **553**, 399–401.
64. Wrieden WL & Barton KL (2011) The Scottish Diet: Estimations of Energy Density and Expenditure. https://www.foodstandards.gov.scot/downloads/Energy_Density_Report.pdf (accessed March 2022).
65. Ledikwe JH, Blanck HM, Khan LK, *et al.* (2005) Dietary energy density determined by eight calculation methods in a nationally representative United States population. *J Nutr* **135**, 273–278.
66. Cox DN & Mela DJ (2000) Determination of energy density of freely selected diets: methodological issues and implications. *Int J Obes* **24**, 49–54.
67. Raynor HA, Thomas JG, Cardoso CC, *et al.* (2019) Examining the pattern of new foods and beverages consumed during obesity treatment to inform strategies for self-monitoring intake. *Appetite* **132**, 147–153.
68. Galante J, Adamska L, Young A, *et al.* (2015) The acceptability of repeat Internet-based hybrid diet assessment of previous 24 h dietary intake: administration of the Oxford WebQ in UK Biobank. *Br J Nutr* **115**, 681–686.
69. Carter MC, Hancock N, Albar SA, *et al.* (2016) Development of a new branded UK food composition database for an online dietary assessment tool. *Nutrients* **8**, 480.
70. Marconi S, Durazzo A, Camilli E, *et al.* (2018) Food composition databases: considerations about complex food matrices. *Foods* **7**, 2.
71. Wilkinson LL, Embling R, Raynor HA, *et al.* (2022) Multi-component food-items and eating behaviour: what do we know and what do we need to know? *Appetite* **168**, 105718.
72. Piernas C, Perez-Cornago A, Gao M, *et al.* (2021) Describing a new food group classification system for UK biobank: analysis of food groups and sources of macro- and micronutrients in 208,200 participants. *Eur J Nutr* **60**, 2879–2890.
73. Major B, Tomiyama AJ & Hunger JM (2018) *The Negative and Bidirectional Effects of Weight Stigma on Health*. Oxford Handbook of Stigma, Discrimination, and Health. New York, NY: Oxford University Press. pp. 499–519.
74. Robinson E, Benwell H & Higgs S (2013) Food intake norms increase and decrease snack food intake in a remote confederate study. *Appetite* **65**, 20–24.
75. Patalay P & Hardman CA (2019) Comorbidity, codevelopment, and temporal associations between Body Mass Index and internalizing symptoms from early childhood to adolescence. *JAMA Psychiatr* **76**, 721–729.
76. Davies NM, Dickson M, Davey Smith G, *et al.* (2018) The causal effects of education on health outcomes in the UK Biobank. *Nat Hum Behav* **2**, 117–125.
77. Fry A, Littlejohns TJ, Sudlow C, *et al.* (2017) Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. *Am J Epidemiol* **186**, 1026.