

# The growth of the outflows from evolved stars: A live poster

Tiina Liimets<sup>1,2</sup>, Romano L. M. Corradi<sup>3,4</sup>, Bruce Balick<sup>5</sup>  
and Miguel Santander-García<sup>6,7,3,4</sup>

<sup>1</sup>Tartu Observatoorium, Tõravere, 61602, Estonia, email: [tiina@not.iac.es](mailto:tiina@not.iac.es)

<sup>2</sup>Nordic Optical Telescope, Apartado 474, E-38700 Santa Cruz de La Palma, Spain

<sup>3</sup>Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

<sup>4</sup>Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

<sup>5</sup>Astronomy Department, University of Washington, Seattle, WA 98195, USA

<sup>6</sup>Observatorio Astronómico Nacional, Madrid, Spain

<sup>7</sup>Centro de Astrobiología, INTA-CSIC, Madrid, Spain

**Abstract.** We present our long-term monitoring, using both ground-based telescopes and the HST, of the expansion in the plane of the sky of the outflows from evolved stars. They include the PN M2–9 (suspected of hosting a symbiotic nucleus), the classical nova remnant GK Per, the symbiotic Miras R Aqr, Hen 2–147 and Hen 2–104, and the proto-PNe CRL 618, CRL 2688, and Hen 3–1475.

**Keywords.** binaries: symbiotic, stars: individual (M2-9, R Aqr, Hen 2-147, Hen 2-104 GK Per), novae, cataclysmic variables, planetary nebula: individual (CRL 618, CRL 2688, Hen 3-1475), interstellar medium: kinematics and dynamics


---

The image quality achievable with modern telescopes allows the apparent growth of relatively close/fast stellar outflows to be revealed and measured in timescales of few years. The information about the tangential component of the gas and shocks motions, coupled with line-of-sight velocities from Doppler shifts, allows a detailed view of the kinematics of the outflows. Combining this information from two or more epochs provides us with a tool to test the forces acting on the ejecta (i.e its dynamical evolution). With this goal, we have obtained multi-epoch images of several sources over several years. Some results are briefly summarised below (those on pre-PNe can be found in the contribution of B. Balick to these proceedings and in Balick *et al.* 2011, *ApJ*, submitted). Outflows are illustrated in Fig. 1.

*M2-9.* The unique property of this object is its 90-yr rotating pattern in the inner lobes. Given its corkscrew geometry and associated travelling times, excitation structure, we conclude that the phenomenon is caused by a particle jet from the central source hitting and exciting the walls of the inner lobes of M2-9 (Corradi *et al.* 2011). The speed of the jet – between 11000 and 16000 km s<sup>-1</sup> – would be unusually high among PNe and symbiotic stars.

*GK Per.* Kinematical modeling of the nova remnant shows that the nebula is a thick clumpy shell, with knots expanding with a range of speeds between 300 and 1000 km s<sup>-1</sup>. Modest deceleration occurred since their ejection on 1901. The nebula is overall fading, but individual knots show significant brightness changes (Liimets *et al.*, in preparation).

*Hen 2-147, Hen2-104, R Aqr.* Symbiotic novae illustrate the mass-loss geometry from long-period interacting binaries. The expansion pattern of Hen 2-147 and Hen2-104 (the “Southern Crab”), their expansion parallaxes, the role of shocks, are discussed in



**Figure 1.** Images of the studied outflows.

Santander *et al.* (2007) and Santander *et al.* (2008). The study of the expansion of the jet of R Aqr is in progress.

### References

- Corradi, R. L. M., Balick, B., & Santander-García, M. 2011, *A&A*, 529, 43  
 Santander-García, Corradi, R. L. M., Whitelock, P. A., *et al.* 2007, *A&A*, 465, 481  
 Santander-García, Mampaso, A., Morisset, C., *et al.* 2008, *A&A*, 485, 117