

Vegetarian diets and cardiovascular risk factors in black members of the Adventist Health Study-2

Gary Fraser^{1,*}, Sozina Katuli¹, Ramtin Anousheh², Synnove Knutsen¹, Patti Herring³ and Jing Fan⁴

¹Department of Biostatistics, Loma Linda University, 24951 North Circle Drive, NH 2005, Loma Linda, CA 92350, USA: ²Internal Medicine, VA Loma Linda Healthcare System, Loma Linda, CA, USA: ³Health Promotion and Education, Loma Linda University, Loma Linda, CA, USA: ⁴Adventist Health Study, Loma Linda University, Loma Linda, CA, USA

Submitted 8 April 2013: Final revision received 18 November 2013: Accepted 21 November 2013: First published online 17 March 2014

Abstract

Objective: To compare cardiovascular risk factors between vegetarians and non-vegetarians in black individuals living in the USA.

Design: A cross-sectional analysis of a sub-set of 592 black women and men enrolled in the Adventist Health Study-2 (AHS-2) cohort of Seventh-day Adventists.

Setting: Members of the AHS-2 cohort, who lived in all states of the USA and provinces of Canada.

Subjects: Black/African-American members of two sub-studies of AHS-2 where blood and physiological measurements were obtained.

Results: Of these women and men, 25% were either vegan or lacto-ovo-vegetarians (labelled ‘vegetarian/vegans’), 13% were pesco-vegetarian and 62% were non-vegetarian. Compared with non-vegetarians, the vegetarian/vegans had odds ratios for hypertension, diabetes, high blood total cholesterol and high blood LDL-cholesterol of 0.56 (95% CI 0.36, 0.87), 0.48 (95% CI 0.24, 0.98), 0.42 (95% CI 0.27, 0.65) and 0.54 (95% CI 0.33, 0.89), respectively, when adjusted for age, gender, education, physical activity and sub-study. Corresponding odds ratios for obesity in vegetarian/vegans and pesco-vegetarians, compared with non-vegetarians, were 0.43 (95% CI 0.28, 0.67) and 0.47 (95% CI 0.27, 0.81), respectively; and for abdominal obesity 0.54 (95% CI 0.36, 0.82) and 0.50 (95% CI 0.29, 0.84), respectively. Results for pesco-vegetarians did not differ significantly from those of non-vegetarians for other variables. Further adjustment for BMI suggested that BMI acts as an intermediary variable between diet and both hypertension and diabetes.

Conclusions: As with non-blacks, these results suggest that there are sizeable advantages to a vegetarian diet in black individuals also, although a cross-sectional analysis cannot conclusively establish cause.

Keywords
Vegetarian diet
Cardiovascular risks
Blacks
Adventists

CVD is a leading cause of death^(1,2) and this is also so among African Americans⁽³⁾. Among non-Hispanic blacks aged 20 years and older, 46% suffer from CVD⁽⁴⁾, and CVD caused the deaths of nearly 100 000 African Americans in 2012. CVD risk factors such as age and gender are non-modifiable, while smoking, blood pressure (BP), diabetes, blood lipids and obesity, particularly waist size, can be modified. Of these, hypertension may be most important⁽⁵⁾, particularly considering its high prevalence in this group⁽⁶⁾.

A substantial body of evidence has shown that diet plays a role in the pathogenesis of cardiovascular risk factors. Hypertension, for example, can sometimes be controlled by diet alone as verified by the DASH (Dietary Approaches to Stop Hypertension) diet⁽⁷⁾, which is rich in

fruits, vegetables and low-fat dairy foods, and significantly lowers BP. The mechanism of diet's effect on CVD risk factors is complex. Evidence available from a number of studies links several nutrients, minerals, food groups and dietary patterns with an increased or decreased risk of CVD. Dietary *trans* and saturated fats, for example, are associated with increased risk of CVD most likely because they increase blood cholesterol⁽⁸⁾, while polyunsaturated fats reduce blood cholesterol and appear to be protective^(9–11). Dietary Na is associated with higher BP, while dietary K, well represented in fruits and vegetables, has been associated with reduced risk of hypertension, CHD and stroke⁽¹²⁾.

Americans subscribe to a variety of diets aside from the standard American diet. These include the Mediterranean

*Corresponding author: Email gfraser@llu.edu

diet and various vegetarian diets such as lacto-ovo-vegetarian, pesco-vegetarian and less commonly the vegan diet which excludes all animal products. Research suggests that a Mediterranean diet is protective for CVD⁽¹³⁾. Other studies have shown that vegetarians also have lower BP values and lower risk of CHD^(14,15). In addition, vegetarian diets probably lower rates of diabetes mellitus⁽¹⁶⁾ and the prevalence of obesity⁽¹⁷⁾. Researchers attribute these benefits to less cholesterol and saturated fat in the vegetarian diet, also to the higher content of unsaturated fats, fibre and K^(18,19) and that these diets contain more foods with a lower glycaemic index.

However, much less evidence exists for the health effects of vegetarian diets among African Americans. Studies by Melby *et al.* have suggested that African-American vegetarians exhibit a more favourable serum lipid profile and also lower BP values than non-vegetarians^(20,21). Studies conducted in Nigeria among native African Seventh-day Adventists demonstrated that the vegetarian diet in this setting also is associated with lower levels of serum total cholesterol (TC), TAG and perhaps BP^(22,23).

The African-American diet is typically rich in both fats and complex carbohydrates⁽²⁴⁾. Obesity is highly prevalent, as 44% of non-Hispanic blacks have BMI above 30 kg/m²⁽²⁵⁾. In addition, according to estimates in 2011, the prevalence of diabetes is 53% above the national average in black individuals⁽²⁶⁾. We therefore conducted a study in blacks (75% African American and 25% West Indian) attending Adventist churches in the USA and Canada. Clinics were conducted to measure BP, blood lipids, blood glucose, waist circumference (WC) and BMI. The objective of the present report is to test the hypotheses that there are associations between the identified cardiovascular/metabolic risk factors and different dietary patterns among black individuals.

Experimental methods

Four hundred and seven black women and 185 black men were selected for the present study, which is nested within the Adventist Health Study-2 (AHS-2) cohort. Specifically, these black individuals were those enrolled in the AHS-2 calibration sub-study (n 427) and the Bio-physical Manifestations of Religion sub-study (BioMRS, n 165). All individuals were also part of AHS-2. The sub-studies were joined to increase numbers of black participants. The detailed recruitment methods of the AHS-2, the calibration study, the particular methods used to recruit black participants and the BioMRS are described elsewhere^(27–30). Briefly, the calibration sub-study participants were representative of all black individuals in AHS-2, and thus were aged 30 years and above, and lived throughout the USA and Canada. Participants from the BioMRS clinics however, lived in Southern California and were at least 50 years of age.

At the clinics, an overnight fasting blood sample, an overnight urine collection and a subcutaneous adipose sample were collected. Biometrics, percentage of body fat (bio-impedance), BP, lipid profiles and selected metabolic parameters were measured using defined protocols. BP was measured while seated and after resting for at least 10 min, using an Omron automated sphygmomanometer⁽³¹⁾. Systolic BP and diastolic BP were estimated three times 1–2 min apart and the second and third readings were averaged. Lipid profiles – TC (mg/dl), HDL-cholesterol (HDL-C; mg/dl), TAG (mg/dl) and the ratio TC:HDL-C – were measured from fasting blood samples collected in heparin tubes, using a Cholesterol LDX Analyzer (Cholestech Corp., Hayward, CA, USA). LDL-cholesterol (LDL-C) was calculated using the Friedewald equation (TC minus HDL-C minus very-low-density lipoprotein-cholesterol (estimated as TAG divided by 5)) when TAG was <400 mg/dl. The two participants with TAG >400 mg/dl were excluded from this analysis. Weight was determined using the scale function of the Tanita BF-350 Body Fat Analyzer (Tanita UK Ltd, Middlesex, UK). Height was measured using a portable stadiometer, and then BMI was calculated as [weight (kg)]/[height (m)]². The circumferences of waist and hip were measured using a tape, and their ratio was calculated.

Participants in the BioMRS completed the baseline AHS-2 FFQ within 2–3 years before their clinic visit. Those in the calibration sub-study completed an identical FFQ within 2–3 months of the clinic visit. These FFQ were used to define the dietary patterns. Details of these questionnaires and their use to define the dietary patterns are given elsewhere^(17,32). Briefly, the dietary patterns are: (i) vegans, who eat no animal products; (ii) lacto-ovo-vegetarians, who eat dairy and/or eggs but no meat; (iii) pesco-vegetarians, who eat fish but no other meats; and (iv) non-vegetarians who eat meats at least once per week.

Physical activity was assessed as hours spent per day in moderate, vigorous and extremely vigorous activities. These were based on questions that gave examples of such activities, including work and leisure exercise, and separated weekdays from weekend days.

Data analysis

The odds ratios of prevalent hypertension, diabetes, larger WC and high lipid levels, comparing dietary patterns, were estimated by the use of logistic regression. Hypertension was defined as any participant with systolic BP ≥ 140 mmHg or diastolic BP ≥ 90 mmHg or taking BP-lowering medications; elevated TC (or LDL-C) was defined as values ≥ 200 mg/dl (or ≥ 130 mg/dl for LDL-C) or taking lipid-lowering agents; low HDL-C was defined as values <40 mg/ml (males) or <50 mg/dl (females); elevated TAG was defined as values ≥ 150 mg/dl; diabetes was defined as fasting plasma glucose ≥ 126 mg/dl or taking diabetic medications. Covariates always included an indicator variable specifying which sub-study an individual

came from. The identity of medications used by study participants and which are commonly used to treat hypertension, hyperlipidaemia or diabetes was determined by a cardiologist and internist (G.F.).

Adjusted mean levels of each risk factor by dietary pattern among participants not taking medications for the medical condition or associated risk factor under consideration were calculated from ANCOVA, adjusting for age, gender, education, physical activity and an indicator variable for sub-study. In separate analyses adjustment was also made for BMI. Small percentages of missing data (3–7%) for individual dietary variables used to establish dietary patterns were imputed as described elsewhere^(14,33).

Statistical analyses were performed using the SAS statistical software package version 9.3.

Results

The baseline demographics of the 592 participants, stratified by dietary pattern, are summarized in Table 1. In the combined data, fifty-one participants were vegans, ninety-five lacto-ovo-vegetarians, eighty pesco-vegetarians and 366 were non-vegetarians. We further combined vegans and lacto-ovo-vegetarians because of the small vegan population and because in nearly all dietary respects Adventist lacto-ovo-vegetarians fall between vegans and non-vegetarians⁽³⁴⁾. The new category is referred to as

'vegetarian/vegans'. Non-vegetarians tended to be less educated and to have higher BMI than participants with other dietary patterns (Table 1).

Apart from age ($P < 0.0001$), which is a consequence of the design of the two sub-studies (as is probably true also for educational status, $P < 0.0001$), there were no statistically significant differences in covariate values between the two sub-study groups. Therefore, all subsequent analyses were adjusted for age and education, as well as gender and sub-study indicator variable. The BioMRS participants were, however, more likely to be vegan and less likely to be pesco-vegetarian than the calibration sub-study participants.

Hypertension and diet

Of the 592 participants, 305 were classified as hypertensive and 212 of these were taking anti-hypertensive medications. In multiple logistic regression analysis (Table 2), the vegetarian/vegan group had a significantly lower odds of hypertension compared with non-vegetarians (OR = 0.56, $P = 0.01$). Results for pesco-vegetarians did not differ significantly from either of the other groups. When restricting the analysis to all participants not taking anti-hypertensive medications (Table 3), adjusted mean systolic and diastolic BP values were on average 1.3 and 0.6 mmHg ($P = 0.57$ and 0.67) lower, respectively, in vegetarian/vegans than in non-vegetarians. Values for pesco-vegetarians not taking anti-hypertensive medications were higher than those for vegetarian/vegans ($P = 0.25$) and non-vegetarians ($P = 0.43$).

Table 1 Relationship of selected demographic factors to dietary patterns among black participants in the Adventist Health Study-2 (AHS-2)

	Vegetarian/vegan (<i>n</i> 146; 25%)		Pesco-vegetarian (<i>n</i> 80; 13%)		Non-vegetarian (<i>n</i> 366; 62%)		All (<i>n</i> 592) <i>P</i> value*
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Age (years)							
<50	26	18	19	24	97	27	0.40
50–59	49	33	22	27	107	29	
60–69	39	27	24	30	83	23	
≥70	32	22	15	19	79	21	
Mean		61		60		59	0.17
sd		12		12		13	
Gender							
Women	95	65	54	68	258	70	0.53
Education							
High school or below	19	13	13	16	71	20	0.01
Some college	60	41	32	40	181	49	
Bachelors and above	67	46	35	44	114	31	
BMI (kg/m ²)							
<25.0	51	35	21	26	71	19	<0.0001
25.0–29.9	56	38	37	46	124	34	
≥30.0	39	27	22	28	171	47	
Mean		27.6		28.2		30.8	
sd		5.6		5.7		7.3	
Physical activity (h/d)							
Mean		2.4		2.6		2.8	0.54
sd		2.9		3.5		3.4	
Sub-study							
Calibration	89	61	64	80	274	75	0.0016
BioMRS	57	39	16	20	92	25	

BioMRS, Biophysical Manifestations of Religion sub-study.

**P* values test differences between means of dietary patterns or the null hypotheses of no association between the named variable and dietary pattern.

Table 2 Multiple logistic regression analyses to test associations between major cardiovascular risk factors and dietary patterns* among black participants in the Adventist Health Study-2 (AHS-2)

	Vegetarian/vegan (reference: Non-vegetarian)		Pesco-vegetarian (reference: Non-vegetarian)		Vegetarian/vegan (reference: Pesco-vegetarian)	
	OR	95% CI	OR	95% CI	OR	95% CI
A. Not adjusted for BMI						
Hypertension†	0.56	0.36, 0.87	0.94	0.54, 1.63	0.60	0.32, 1.11
Diabetes	0.48	0.24, 0.98	1.11	0.56, 2.22	0.44	0.23, 0.79
High TC‡	0.42	0.27, 0.65	0.99	0.56, 1.75	0.42	0.23, 0.79
High LDL-C‡	0.54	0.33, 0.89	1.31	0.77, 2.22	0.41	0.23, 0.75
Low HDL-C‡	0.69	0.45, 1.07	0.99	0.59, 1.66	0.70	0.38, 1.27
High TAG‡	1.05	0.56, 2.00	0.82	0.37, 1.83	1.28	0.51, 3.21
Obesity§, females	0.38	0.22, 0.65	0.59	0.32, 1.12	0.64	0.31, 1.36
Obesity, males	0.52	0.24, 1.13	0.22	0.06, 0.78	2.39	0.61, 9.34
Obesity, both sexes	0.43	0.28, 0.67	0.47	0.27, 0.81	0.92	0.49, 1.72
Abdominal adiposity§, females	0.55	0.33, 0.92	0.53	0.29, 1.00	1.04	0.50, 2.14
Abdominal adiposity, males	0.48	0.22, 1.04	0.43	0.15, 1.21	1.10	0.35, 3.53
Abdominal adiposity, both sexes	0.54	0.36, 0.82	0.50	0.29, 0.84	1.09	0.59, 1.99
B. Adjusted for BMI						
Hypertension	0.73	0.46, 1.16	1.16	0.65, 2.07	0.63	0.33, 1.19
Diabetes	0.60	0.29, 1.25	1.29	0.63, 2.63	0.47	0.19, 1.12
'High cholesterol'	0.44	0.28, 0.70	1.03	0.58, 1.84	0.43	0.23, 0.80
High LDL-C	0.61	0.36, 1.03	1.44	0.83, 2.48	0.42	0.23, 0.77
Low HDL-C	0.88	0.47, 1.66	1.20	0.62, 2.31	0.73	0.33, 1.66
High TAG	1.27	0.66, 2.45	0.93	0.41, 2.09	1.38	0.55, 3.45

TC, total cholesterol; LDL-C, LDL-cholesterol; HDL-C, HDL-cholesterol; BP, blood pressure; WC, waist circumference.

*A separate logistic regression model was run for each risk factor, in each case adjusted for age, gender (as appropriate), education, physical activity, sub-study indicator and BMI as indicated.

†Hypertension is systolic BP \geq 140 mmHg and/or diastolic BP \geq 90 mmHg or taking medications for high BP.

‡High TC is \geq 200 mg/dl and high LDL-C is \geq 130 mg/dl or taking cholesterol-lowering medications in either case; low HDL-C is $<$ 40 mg/dl (males) or $<$ 50 mg/dl (females); high TAG is $>$ 150 mg/dl.

§Obesity is BMI \geq 30.0 kg/m²; abdominal adiposity is WC $>$ 88 cm in females and $>$ 102 in males.

Diabetes and diet

Of 592 participants, eighty-five were classified as diabetic and fifty-six of these were taking diabetic medications. The prevalence of diabetes was 8.9% in vegetarian/vegans, 18.8% in pesco-vegetarians and 15.6% in non-vegetarians. Vegetarian/vegan diets were associated with a lower risk of diabetes compared with non-vegetarians (OR = 0.48, P = 0.04; Table 2). Of all vegetarian/vegans who were not taking anti-diabetic medications, the average adjusted fasting plasma glucose level was 7.3 mg/dl lower (P = 0.0003) than that of non-vegetarians (Table 3). Pesco-vegetarians were not significantly different in odds ratio of prevalent diabetes when compared with either vegetarian/vegan or non-vegetarian groups, but those not on medication had higher glucose levels than vegetarian/vegans (P = 0.004).

Serum total cholesterol and diet

Of vegetarian/vegan participants, 30% were classified as having 'high cholesterol' compared with 48% in both the pesco- and non-vegetarian groups. The odds ratio for high cholesterol in the vegetarian/vegan group, as compared with the non-vegetarians, was 0.42 (P = 0.0001), and a similar value was observed when comparing to pesco-vegetarians (OR = 0.42, P = 0.006; Table 2). Vegetarian/vegans who were not taking cholesterol-lowering medications had an average adjusted fasting plasma TC level that was 10.0 mg/dl lower (P = 0.01) than that in equivalent

non-vegetarians (Table 3). Levels in pesco-vegetarians did not differ significantly from those in non-vegetarians.

Serum LDL-cholesterol and diet

In the vegetarian/vegan group 28% of participants met the criteria for high LDL-C, but 48% of pesco-vegetarians and 43% of non-vegetarians were classified in the high LDL-C category. Thus, vegetarian/vegans had a significantly lower odds of high LDL-C compared with non-vegetarians (OR = 0.54, P = 0.015), and a significantly lower odds was also observed when comparing them to pesco-vegetarians (OR = 0.41, P = 0.003; Table 2). Vegetarian/vegans who were not taking cholesterol-lowering medications had on average a 9.6 mg/dl (P = 0.014) lower adjusted mean fasting plasma LDL-C level compared with equivalent non-vegetarians (Table 3). Levels in pesco-vegetarians did not differ significantly from those in non-vegetarians.

Serum HDL-cholesterol and TAG

In these analyses we did not observe any significant differences in fasting plasma levels of HDL-C or TAG among dietary patterns (Tables 2 and 3).

BMI, waist circumference and diet

Results are computed separately for males and females as we found that the average BMI in females was significantly higher than that of males. The odds ratios for obesity (BMI \geq 30.0 kg/m²) and abdominal adiposity

Table 3 Adjusted mean levels of risk factors* among black participants in the Adventist Health Study-2 (AHS-2)

	Vegetarian/vegan	Pesco-vegetarian	Non-vegetarian	<i>P</i> value, vegetarian/vegan v. pesco-vegetarian	<i>P</i> value, vegetarian/vegan v. non-vegetarian	<i>P</i> value, pesco-vegetarian v. non-vegetarian
A. Not adjusted for BMI						
Systolic BP (mmHg)	119.8	123.5	121.1	0.25	0.57	0.43
Diastolic BP (mmHg)	74.3	75.8	74.9	0.44	0.67	0.62
Fasting blood glucose (mg/dl)	82.6	90.7	89.9	0.004	0.0003	0.75
TC (mg/dl)	177.6	183.7	187.6	0.29	0.013	0.46
LDL-C (mg/dl)	110.9	119.1	120.5	0.14	0.014	0.77
BMI (kg/m ²), females	27.8	28.3	31.1	0.72	0.0004	0.018
BMI (kg/m ²), males	26.9	27.2	30.7	0.84	0.0007	0.012
BMI (kg/m ²), both sexes	27.3	27.6	30.7	0.74	<0.0001	0.0006
WC (cm), females	88.1	89.0	95.6	0.75	0.0004	0.017
WC (cm), males	95.5	94.7	103.5	0.82	0.0035	0.0093
WC (cm), both sexes	91.6	91.6	99.2	0.99	<0.0001	0.0003
HDL-C (mg/dl)	51.5	49.9	49.4	0.47	0.18	0.80
TAG (mg/dl)	101.8	100.6	104.7	0.89	0.64	0.59
B. Adjusted for BMI						
Systolic BP (mmHg)	121.1	124.6	120.8	0.29	0.88	0.21
Diastolic BP (mmHg)	75.3	76.6	74.7	0.51	0.66	0.29
Fasting blood glucose (mg/dl)	84.8	91.9	90.4	0.019	0.0090	0.60
TC (mg/dl)	178.0	184.2	187.5	0.29	0.021	0.53
LDL-C (mg/dl)	111.6	119.8	120.3	0.14	0.027	0.92
HDL-C (mg/dl)	50.2	48.6	49.9	0.47	0.86	0.51
TAG (mg/dl)	105.9	104.7	103.3	0.89	0.68	0.85

BP, blood pressure; TC, total cholesterol; LDL-C, LDL-cholesterol; WC, waist circumference; HDL-C, HDL-cholesterol.

*Among participants not on medication for that particular risk factor (applies to hypertension, diabetes and lipids), adjusted for age, gender (as appropriate), education, sub-study indicator, physical activity and BMI as indicated.

†*P* values test the null hypotheses of no difference between the pairs of dietary groups indicated.

(WC >88 cm in females and >102 cm in males) are shown in Table 2. Those for obesity are 0.38 (females, $P=0.0005$) and 0.52 (males, $P=0.10$) comparing vegetarian/vegans with non-vegetarians. Corresponding odds for abdominal adiposity are 0.55 ($P=0.02$) and 0.48 ($P=0.06$). Thus similar trends in both sexes were noted by dietary pattern except that the point estimates for the pesco-vegetarian males were non-significantly lower than those for vegetarian/vegans.

Adjusted mean BMI and WC data for each dietary pattern are shown in Table 3. Both mean BMI and WC were significantly associated with the vegetarian dietary patterns. A trend of increasing BMI was noted passing from vegetarian/vegan, to pesco-vegetarian, to non-vegetarian patterns, the difference between vegetarian/vegan and non-vegetarian being highly significant. This was true for both genders. Values of mean WC were significantly lower in both vegetarian/vegans and pesco-vegetarians compared with non-vegetarian participants. Pesco-vegetarians had significantly lower mean BMI ($P=0.0006$) and WC ($P=0.0003$) than non-vegetarians, but did not differ significantly from vegetarian/vegans on either the odds ratios or the adjusted means.

Tables 2 and 3 also show results with additional adjustment for BMI. In Table 2 there is usually some moderate movement of the effect estimate towards the null, indicating that differences in BMI according to diet partially mediate the dietary effects. However, dietary effects on TC and LDL-C still retain statistical significance (or very close), suggesting that here diet also acts independently of BMI. In Table 3, among those not taking diabetic or lipid-lowering medications, significant differences remain between dietary patterns for TC, LDL-C and fasting glucose.

Discussion

We found that black vegans and lacto-ovo-vegetarians (as a combined category) had significantly lower prevalences of major CVD risk factors compared with non-vegetarian black Adventists. These results are from a population that includes both genders, a wide age range and a broad span of socio-economic status and location of residence. There is a growing body of evidence about the beneficial effects of vegetarian dietary habits on cardiovascular health among non-black populations, but similar effects have not been extensively studied among black populations. As we have previously reported, black Adventists may have lower rates of CVD than other blacks⁽³⁵⁾. If so, this could be due in part to the sizeable percentage of vegetarians among the Adventists.

As compared with non-vegetarians, vegetarian/vegans had a 44% lower odds of hypertension, an important finding for black individuals where hypertension is so prevalent. However, there was no significant difference in risk between pesco-vegetarians and non-vegetarians.

These results generally agree with those of another study of black Adventists in Nigeria where omnivores had higher systolic BP values than vegetarians, although this was not statistically significant⁽²²⁾. Two studies of African Americans^(20,21) demonstrated significantly lower BP in vegetarians, with the first showing lower systolic pressures (by 10.2 mmHg)⁽²¹⁾ compared with omnivores and the second study finding similar but non-significant trends⁽²⁰⁾. Among blacks in Barbados who had been vegetarian for at least 5 years there was a lower percentage with self-reported hypertension (although not statistically significant)⁽³⁶⁾. The CARDIA (Coronary Artery Risk Development in Young Adults) study⁽³⁷⁾ investigated a cohort with approximately equal numbers of black and white participants. Consistent across race–sex groups the intake of plant foods was inversely associated, while meat was positively associated, with elevated BP. Although analyses were adjusted for race and other factors, it was not stated if this was true separately in black individuals. The DASH diet also lowered BP significantly by 6.9/3.7 mmHg in African Americans⁽³⁸⁾. This diet has a reduced amount of meat and more fruit, vegetables and whole grains than a typical US diet.

Protective associations between vegetarian diets and high BP are consistent with studies of non-black populations. Several small experimental studies^(39–42) and larger observational studies⁽⁴³⁾, including white participants in AHS-2⁽¹⁴⁾, have generally found similar results to those that we report here in black North Americans. Further, studies of non-vegetarians have often, in multivariate analyses, associated plant foods (such as whole grains, fruits, etc.) with lower BP and meat consumption with higher BP, as recently reviewed by Steffen *et al.*⁽³⁷⁾.

The mechanisms underlying the effect of vegetarian diets on hypertension are unclear. However, the greater intakes of K, other minerals and fibre in vegetarian diets have been postulated to reduce BP⁽⁴⁴⁾, as have higher intakes of the minerals, Ca, K, Mg and P⁽⁴⁵⁾. It is possible that meat appears to increase the risk of hypertension because of its association with fewer vegetables and fruits. As found by others⁽⁴³⁾, it also seems clear in our data that a large part of this effect is related to the lower BMI among vegetarians. Plant foods are also known to protect against insulin resistance which in turn is related to hypertension and other risk factors for heart disease⁽⁴⁶⁾.

Our study also found that vegetarian/vegans have less than half the odds of diabetes of non-vegetarian black participants, a potentially important finding for black women particularly, as they had higher glycaemic index and glycaemic load than Caucasians in the SWAN (Study of Women's Health Across the Nation) cohort⁽⁴⁷⁾. In black participants from the parent AHS-2 cohort, Tonstad *et al.*⁽¹⁶⁾ reported in covariate-adjusted analyses that vegans (OR = 0.381; 95% CI 0.236, 0.617), lacto-ovo-vegetarians (OR = 0.618; 95% CI 0.503, 0.760) and semi-vegetarians (OR = 0.486, 95% CI 0.312, 0.755) had a lower self-reported

incidence of diabetes than non-vegetarians. A few other cohorts have found that certain foods, often those preferentially consumed by vegetarians, were associated with lower risk of diabetes in black individuals. Analysis of a small cohort of ethnic minority women (half were black) from the Nurses' Health Study, but with long follow-up, suggested that a healthy diet high in cereal fibre but low in *trans* fat and glycaemic load was perhaps even more strongly associated with reduced diabetes risk than in white individuals⁽⁴⁸⁾. In the 59 000 black women enrolled in the Black Women's Health Study, higher intake of cereal fibre was associated with lower risk, and higher glycaemic index with higher risk of diabetes⁽⁴⁹⁾. The ARIC (Atherosclerosis Risk in Communities) study found a protective association between cereal fibre and diabetes risk in both white and black participants, but this was not statistically significant among the blacks⁽⁵⁰⁾. However, in the CARDIA study (where half of the subjects were black) those following the Dietary Guidelines for Americans had increased insulin resistance compared with others in this population⁽⁵¹⁾.

Studies of vegetarian diets from other populations, and risk of diabetes in black or non-black individuals, are rare, although much is known of the use of such diets in the treatment and control of diabetes⁽⁵²⁾. A previous cohort of US Adventists has provided strong positive associations between intake of meat and incidence of diabetes⁽⁵³⁾.

Several mechanisms are postulated for the beneficial effects of the vegetarian diet on diabetes risk. Undoubtedly the most important is the lower rates of overweight associated with these diets, and in our data, the magnitude of the effect is diminished after adjustment for BMI. Vegetarian diets usually also have lower glycaemic load, which is known to decrease risk of diabetes⁽⁵⁴⁾. Red meats increase insulin resistance, an effect possibly mediated by saturated fats⁽⁵⁵⁾. In addition, fruit and vegetable intake is associated with reduced oxidative stress and inflammation, possible precursors of diabetes⁽⁵⁶⁾.

The vegetarian/vegan diet was associated with 9.6 mg/dl lower LDL-C, 10.0 mg/dl lower TC and just over half the risk of elevated LDL-C as found in non-vegetarians. Other studies of effects of vegetarian diets on lipid levels in black populations are rare. Toohey *et al.*⁽⁵⁷⁾ found that lipid levels in black vegans were more favourable than those of lacto-ovo-vegetarians in the USA. In our study, because of the small number of vegans, we combined these two dietary patterns. Melby *et al.*⁽²⁰⁾ found lower TC, LDL-C and TAG among sixty-six black vegetarians compared with forty-five non-vegetarians in the USA. Very similar results were demonstrated by Famodu *et al.*⁽²²⁾ among black Adventists (vegetarian and non-vegetarian) in Nigeria.

Vegetables are known to decrease TC and LDL-C values⁽⁵⁸⁾, while saturated fats significantly increase and polyunsaturated fats lower both serum LDL-C and TC⁽⁵⁹⁾. Dietary fibre and phytosterols, found only in plant foods, also have a well-established but modest effect to lower

serum cholesterol⁽⁶⁰⁾, whereas red meat raises it⁽³⁹⁾. Thus there is every reason to expect a more favourable lipid profile among vegetarians, which has usually been found^(61–63).

Our results showed that vegetarian/vegans have a risk of obesity that is only 43% that of non-vegetarian black participants. The risk of abdominal adiposity was similarly much reduced in both vegetarian dietary patterns. These were the only cardiovascular risk factors where pesco-vegetarians seemed to benefit at least as much as the vegetarian/vegans. Brathwaite *et al.*⁽³⁶⁾ demonstrated less obesity among black participants in Barbados who had been vegetarian for at least 5 years. Other studies in black individuals have also found lower BMI and/or waist and hip sizes among vegetarians^(20–22). The CARDIA study, a cohort consisting of about half black participants, showed strong inverse associations between intake of plant foods and both BMI and WC⁽³⁷⁾. That vegetarians are less overweight has been shown in several non-black research populations^(53,64,65). The mechanisms are unclear but probably include the higher energy density of meats and characteristics of plant foods that replace them in a vegetarian diet. We adjusted for differences in physical activity in the present study. Differences in energy intake by dietary pattern appear to be small⁽³⁴⁾.

Although broadly similar trends are found in Tables 2 and 3, an important difference in Table 3 is that for BP, blood glucose and blood lipids, the results here are limited to those participants who either do not have abnormal values or if they do, are untreated. This is because drug treatment lowers values to an unpredictable extent, perhaps to below normal values in some cases, thus potentially obscuring any dietary effect within the normal range. Even with this exclusion, differences between dietary patterns may be diminished as individuals with more severe abnormality are often excluded. Smaller numbers of participants may also impact statistical significance.

The study has the weakness of being cross-sectional, which does not establish causality. However, most Adventists with health problems will most likely change towards vegetarianism rather than in the opposite direction because of church recommendations on lifestyle. This is indicated by data from AHS-2 (M Martins and GE Fraser, unpublished results) that show strong within-subject trends towards lower animal food intake with ageing. Such dietary changes due to poor health would bias our findings towards the null. A strength of the study is the relatively large sample size of black participants with high-quality dietary data and the careful risk factor assessment.

Conclusion

In summary, we provide evidence that black vegetarians show less hazardous values of several established

cardiovascular risk factors when compared with non-vegetarians. This would be expected to translate into reduced risk of heart attack, stroke and renal failure. Being cross-sectional, our results do not prove a causal association with vegetarianism, but provide some evidence for this.

Acknowledgements

Sources of funding: The authors gratefully acknowledge support from the National Institutes of Health (NIH)/National Cancer Institute (NCI; grant # U01CA152939). The NIH/NCI had no role in the design, analysis or writing of this article. *Conflict of interest:* None of the authors has a conflict of interest to report. *Ethics:* The study was approved by the Loma Linda University Research Protection Program. *Authors' contributions:* G.F. obtained funding, contributed to the study design and wrote part of the manuscript. R.A. performed part of the analysis and reviewed the manuscript. P.H. promoted study enrolment among black individuals, assisted with data collection and reviewed the manuscript. S. Knutsen contributed to the study design, assisted with data collection and reviewed the manuscript. S. Katuli wrote part of the manuscript and reviewed the manuscript. Jing Fan contributed to the data analysis and reviewed the manuscript.

References

- Centers for Disease Control and Prevention, Division for Heart Disease and Stroke Prevention (2013) Heart Disease Fact Sheet. http://www.cdc.gov/dhdsp/data_statistics/fact-sheets/fs_heart_disease.htm (accessed March 2014).
- Miniño AM, Murphy SL, Xu J *et al.* (2011) Deaths: final data for 2008. *Natl Vital Stat Rep* **59**, issue 10, 1–126.
- Yusuf S, Reddy S, Öunpuu S *et al.* (2001) Global burden of cardiovascular diseases. *Circulation* **104**, 2855–2864.
- National Association for the Advancement of Colored People (2012) Health care fact sheet. <http://www.naacp.org/pages/health-care-fact-sheet> (accessed February 2014).
- Kannel WB (1996) Blood pressure as a cardiovascular risk factor: prevention and treatment. *JAMA* **275**, 1571–1576.
- Watson KE (2008) Cardiovascular risk reduction among African Americans: a call to action. *J Natl Med Assoc* **100**, 18–26.
- Moore TJ, Conlin PR, Ard J *et al.* (2001) DASH (Dietary Approaches to Stop Hypertension) diet is effective treatment for stage 1 isolated systolic hypertension. *Hypertension* **38**, 155–158.
- Willett WC (2012) Dietary fats and coronary heart disease. *J Intern Med* **272**, 13–24.
- Iacono JM, Marshall MW, Dougherty RM *et al.* (1975) Reduction in blood pressure associated with high polyunsaturated fat diets that reduce blood cholesterol in man. *Prev Med* **4**, 426–443.
- Anderson JT, Jacobs Jr DR, Foster N *et al.* (1979) Scoring systems for evaluating dietary pattern effect on serum cholesterol. *Prev Med* **8**, 525–537.
- Oliver MF (1997) It is more important to increase the intake of unsaturated fats than to decrease the intake of saturated fats: evidence from clinical trials relating to ischemic heart disease. *Am J Clin Nutr* **66**, 4 Suppl., 980S–986S.
- Srinath Reddy K & Katan MB (2004) Diet, nutrition and the prevention of hypertension and cardiovascular diseases. *Public Health Nutr* **7**, 167–186.
- Williams CM, Lovegrove JA & Griffin BA (2013) Dietary patterns and cardiovascular disease. *Proc Nutr Soc* **72**, 407–411.
- Pettersen BJ, Anousheh R, Fan J *et al.* (2012) Vegetarian diets and blood pressure among white subjects: results from the Adventist Health Study-2 (AHS-2). *Public Health Nutr* **15**, 1909–1916.
- Fraser GE (2003) *Diet, Life Expectancy, and Chronic Disease: Studies of Seventh-Day Adventists and Other Vegetarians*. Oxford/New York: Oxford University Press.
- Tonstad S, Stewart K, Oda K *et al.* (2013) Vegetarian diets and incidence of diabetes in the Adventist Health Study-2. *Nutr Metab Cardiovasc Dis* **23**, 292–299.
- Fraser GE (2009) Vegetarian diets: what do we know of their effects on common chronic diseases? *Am J Clin Nutr* **89**, issue 5, 1607S–1612S.
- Craig WJ (2010) Nutrition concerns and health effects of vegetarian diets. *Nutr Clin Pract* **25**, 613–620.
- Armstrong B, van Merwyk AJ & Coates H (1977) Blood pressure in Seventh-day Adventist vegetarians. *Am J Epidemiol* **105**, 444–449.
- Melby CL, Toohey ML & Cebrick J (1994) Blood pressure and blood lipids among vegetarian, semivegetarian, and nonvegetarian African Americans. *Am J Clin Nutr* **59**, 103–109.
- Melby CL, Goldflies DG & Toohey ML (1993) Blood pressure differences in older black and white long-term vegetarians and nonvegetarians. *J Am Coll Nutr* **12**, 262–269.
- Famodu AA, Osilesi O, Makinde YO *et al.* (1998) Blood pressure and blood lipid levels among vegetarian, semi-vegetarian, and non-vegetarian native Africans. *Clin Biochem* **31**, 545–549.
- Famodu AA, Osilesi O, Makinde YO *et al.* (1999) The influence of a vegetarian diet on haemostatic risk factors for cardiovascular disease in Africans. *Thromb Res* **95**, 31–36.
- Kulkarni KD (2004) Food, culture, and diabetes in the United States. *Clin Diabetes* **22**, 190–192.
- Flegal KM, Carroll MD, Ogden CL *et al.* (2010) Prevalence and trends in obesity among US adults, 1999–2008. *JAMA* **303**, 235–241.
- Centers for Disease Control and Prevention, Diabetes Public Health Resource (2013) 2011 National Diabetes Fact Sheet. http://www.cdc.gov/diabetes/pubs/factsheet11/tables_2.htm (accessed March 2014).
- Butler TL, Fraser GE, Beeson WL *et al.* (2008) Cohort profile: the Adventist Health Study-2 (AHS-2). *Int J Epidemiol* **37**, 260–265.
- Jaceldo-Siegl K, Knutsen SF, Sabate J *et al.* (2010) Validation of nutrient intake using an FFQ and repeated 24 h recalls in black and white subjects of the Adventist Health Study-2 (AHS-2). *Public Health Nutr* **13**, 812–819.
- Herring RP, Butler T, Hall S *et al.* (2010) Recruiting black Americans in a large cohort study: the Adventist Health Study-2 (AHS-2) design, methods and participant characteristics. *Ethn Dis* **20**, 437–443.
- Lee JW, Morton KR, Walters J *et al.* (2009) Cohort profile: the biopsychosocial religion and health study (BRHS). *Int J Epidemiol* **38**, 1470–1478.
- O'Brien E, Mee F, Atkins N *et al.* (1996) Evaluation of three devices for self-measurement of blood pressure according to the revised British Hypertension Society Protocol: the Omron HEM-705CP, Philips HP5332, and Nissei DS-175. *Blood Press Monit* **1**, 55–61.

32. Jaceldo-Siegl K, Sharma S, Knutsen S *et al.* (2009) Validation of nutrient intake by questionnaire (FFQ) against 24-hour recalls in adult Adventists in US and Canada. *Am J Epidemiol* **169**, Suppl. 11, S120.
33. Fraser G & Yan R (2007) Guided multiple imputation of missing data: using a subsample to strengthen the missing-at-random assumption. *Epidemiology* **18**, 246–252.
34. Rizzo NS, Jaceldo-Siegl K, Sabate J *et al.* (2013) Nutrient profiles of vegetarian and nonvegetarian dietary patterns. *J Acad Nutr Diet* **113**, 1610–1619.
35. Montgomery S, Herring P, Yancey A *et al.* (2007) Comparing self-reported disease outcomes, diet, and lifestyles in a national cohort of black and white Seventh-day Adventists. *Prev Chronic Dis* **4**, A62.
36. Brathwaite N, Fraser HS, Modeste N *et al.* (2003) Obesity, diabetes, hypertension, and vegetarian status among Seventh-Day Adventists in Barbados: preliminary results. *Ethn Dis* **13**, 34–39.
37. Steffen LM, Kroenke CH, Yu X *et al.* (2005) Associations of plant food, dairy product, and meat intakes with 15-y incidence of elevated blood pressure in young black and white adults: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Clin Nutr* **82**, 1169–1177.
38. Sacks FM, Appel LJ, Moore TJ *et al.* (1999) A dietary approach to prevent hypertension: a review of the Dietary Approaches to Stop Hypertension (DASH) Study. *Clin Cardiol* **22**, III6–III10.
39. Sacks FM, Donner A, Castelli WP *et al.* (1981) Effect of ingestion of meat on plasma cholesterol of vegetarians. *JAMA* **246**, 640–644.
40. Rouse IL, Beilin LJ, Mahoney DP *et al.* (1983) Vegetarian diet and blood pressure. *Lancet* **2**, 742–743.
41. Sciarone SE, Strahan MT, Beilin LJ *et al.* (1993) Biochemical and neurohormonal responses to the introduction of a lacto-ovo-vegetarian diet. *J Hypertens* **11**, 849–860.
42. Margetts BM, Beilin LJ, Vandongen R *et al.* (1986) Vegetarian diet in mild hypertension: a randomised controlled trial. *Br Med J (Clin Res Ed)* **293**, 1468–1471.
43. Appleby PN, Davey GK & Key TJ (2002) Hypertension and blood pressure among meat eaters, fish eaters, vegetarians and vegans in EPIC-Oxford. *Public Health Nutr* **5**, 645–654.
44. He J & Bazzano LA (2000) Effects of lifestyle modification on treatment and prevention of hypertension. *Curr Opin Nephrol Hypertens* **9**, 267–271.
45. Reusser ME, DiRienzo DB, Miller GD *et al.* (2003) Adequate nutrient intake can reduce cardiovascular disease risk in African Americans. *J Natl Med Assoc* **95**, 188–195.
46. Lee CC, Lorenzo C, Haffner SM *et al.* (2013) Components of metabolic syndrome and 5-year change in insulin clearance – The Insulin Resistance Atherosclerosis Study (IRAS). *Diabetes Obes Metab* **15**, 441–447.
47. Hu Y, Block G, Sternfeld B *et al.* (2009) Dietary glycemic load, glycemic index, and associated factors in a multiethnic cohort of midlife women. *J Am Coll Nutr* **28**, 636–647.
48. Shai I, Jiang R, Manson JE *et al.* (2006) Ethnicity, obesity, and risk of type 2 diabetes in women: a 20-year follow-up study. *Diabetes Care* **29**, 1585–1590.
49. Krishnan S, Rosenberg L, Singer M *et al.* (2007) Glycemic index, glycemic load, and cereal fiber intake and risk of type 2 diabetes in US black women. *Arch Intern Med* **167**, 2304–2309.
50. Stevens J, Ahn K, Juhaeri *et al.* (2002) Dietary fiber intake and glycemic index and incidence of diabetes in African-American and white adults: the ARIC study. *Diabetes Care* **25**, 1715–1721.
51. Zamora D, Gordon-Larsen P, He K *et al.* (2011) Are the 2005 Dietary Guidelines for Americans associated with reduced risk of type 2 diabetes and cardiometabolic risk factors? Twenty-year findings from the CARDIA study. *Diabetes Care* **34**, 1183–1185.
52. Barnard ND, Katcher HI, Jenkins DJ *et al.* (2009) Vegetarian and vegan diets in type 2 diabetes management. *Nutr Rev* **67**, 255–263.
53. Vang A, Singh PN, Lee JW *et al.* (2008) Meats, processed meats, obesity, weight gain and occurrence of diabetes among adults: findings from Adventist Health Studies. *Ann Nutr Metab* **52**, 96–104.
54. Barclay AW, Petocz P, McMillan-Price J *et al.* (2008) Glycemic index, glycemic load, and chronic disease risk – a meta-analysis of observational studies. *Am J Clin Nutr* **87**, 627–637.
55. Bruce WR, Wolever TM & Giacca A (2000) Mechanisms linking diet and colorectal cancer: the possible role of insulin resistance. *Nutr Cancer* **37**, 19–26.
56. Romeo GR, Lee J & Shoelson SE (2012) Metabolic syndrome, insulin resistance, and roles of inflammation – mechanisms and therapeutic targets. *Arterioscler Thromb Vasc Biol* **32**, 1771–1776.
57. Toohey ML, Harris MA, DeWitt W *et al.* (1998) Cardiovascular disease risk factors are lower in African-American vegans compared to lacto-ovo-vegetarians. *J Am Coll Nutr* **17**, 425–434.
58. Fraser GE, Jacobs DR, Anderson JT *et al.* (1981) The effect of various vegetable supplements on serum-cholesterol. *Am J Clin Nutr* **34**, 1272–1277.
59. Jacobs DR Jr, Anderson JT & Blackburn H (1979) Diet and serum cholesterol: do zero correlations negate the relationship? *Am J Epidemiol* **110**, 77–87.
60. Bazzano LA (2008) Effects of soluble dietary fiber on low-density lipoprotein cholesterol and coronary heart disease risk. *Curr Atheroscler Rep* **10**, 473–477.
61. Ferdowsian HR & Barnard ND (2009) Effects of plant-based diets on plasma lipids. *Am J Cardiol* **104**, 947–956.
62. Appleby PN, Thorogood M, McPherson K *et al.* (1995) Associations between plasma lipid concentrations and dietary, lifestyle and physical factors in the Oxford Vegetarian Study. *J Hum Nutr Diet* **8**, 305–314.
63. West RO & Hayes OB (1968) Diet and serum cholesterol levels. A comparison between vegetarians and nonvegetarians in a Seventh-Day Adventist Group. *Am J Clin Nutr* **21**, 853–862.
64. Berkow SE & Barnard N (2006) Vegetarian diets and weight status. *Nutr Rev* **64**, 175–188.
65. Key T & Davey G (1996) Prevalence of obesity is low in people who do not eat meat. *BMJ* **313**, 816–817.