



COMPOSITIO MATHEMATICA

Enough vector bundles on orbispaces

John Pardon

Compositio Math. **158** (2022), 2046–2081.

[doi:10.1112/S0010437X22007783](https://doi.org/10.1112/S0010437X22007783)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865





Enough vector bundles on orbispaces

John Pardon

ABSTRACT

We show that every orbispace satisfying certain mild hypotheses has ‘enough’ vector bundles. It follows that the K -theory of finite rank vector bundles on such orbispaces is a cohomology theory. Global presentation results for smooth orbifolds and derived smooth orbifolds also follow.

1. Introduction

A (separated) orbispace [Hae90, Beh14] is a topological stack X which admits a cover by open substacks of the form Y/Γ (where $\Gamma \curvearrowright Y$ is a continuous action of a finite group on a topological space) and whose diagonal $X \rightarrow X \times X$ is proper (see §3 for background on topological stacks). Familiar examples of orbispaces include orbifolds, graphs of groups, complexes of groups, and (the analytifications of) separated Deligne–Mumford algebraic stacks over \mathbb{C} .

An interesting question to ask about a given orbispace X is whether there exists a global presentation $X = Y/G$ for G a compact Lie group; let us call such an orbispace a *global quotient*. There are a number of known conditions which imply an orbispace is a global quotient. If $X = Y/\Gamma$ for a (possibly infinite) discrete group Γ , then X is a global quotient by Lück and Oliver [LO01, Corollary 2.7]. Every (paracompact) smooth *effective* n -dimensional orbifold is a global quotient (of its orthonormal frame bundle by $O(n)$), and it is an old question whether every (not necessarily effective) smooth orbifold is a global quotient. A sufficient criterion for an orbifold to be a global quotient was given by Henriques and Metzler [HM04]. Henriques [Hen05] conjectured that every compact orbispace is a global quotient, however other experts have expressed skepticism that such a general result would be true [Sie98, §6.4]. The analogous question for algebraic stacks has been studied by Edidin *et al.* [EHKV01] and Totaro [Tot04]. It is a result of Kresch and Vistoli [KV04, Theorem 2], [Kre09, Theorem 4.4] (using a result of Gabber [dJ03], [CTS21, Chapter 4]) that smooth separated Deligne–Mumford stacks over \mathbb{C} with quasi-projective coarse moduli space are global quotients.

Our main result implies that all orbispaces satisfying very mild hypotheses are global quotients. In particular, all compact orbispaces are global quotients.

Received 4 August 2020, accepted in final form 29 August 2022, published online 28 November 2022.

2020 Mathematics Subject Classification 55R91, 57R18 (primary), 55P91, 55S91, 57R91, 19L47, 19L10, 55N32 (secondary).

Keywords: orbispaces, orbibundles, orbifolds, derived orbifolds, K -theory, global homotopy theory, Chern character, orbifold cohomology.

This research was conducted during the period the author served as a Clay Research Fellow and was partially supported by a Packard Fellowship and by the National Science Foundation under the Alan T. Waterman Award, Grant No. 1747553.

© 2022 The Author(s). This is an Open Access article, distributed under the terms of the Creative Commons Attribution-ShareAlike licence (<http://creativecommons.org/licenses/by-sa/4.0>), which permits re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is used to distribute the re-used or adapted article and the original article is properly cited. *Compositio Mathematica* is © Foundation Compositio Mathematica.

THEOREM 1.1. *Let X be an orbispace, with isotropy groups of bounded order, whose coarse space $|X|$ is coarsely finite-dimensional (every open cover has a locally finite refinement with finite-dimensional nerve). Then there exists a complex vector bundle V of rank $n > 0$ over X , whose fiber over every $x \in X$ is isomorphic to a direct sum of copies of the regular representation of the isotropy group G_x . We may take $n = n(d, m)$ if $|X|$ is d -dimensional (every open cover has a locally finite refinement with nerve of dimension $\leq d$) and has isotropy groups of order $\leq m$.*

(Note that the ‘real’ and ‘complex’ versions of Theorem 1.1 are equivalent, by tensoring from \mathbb{R} to \mathbb{C} and by forgetting from \mathbb{C} to \mathbb{R} .)

The proof of Theorem 1.1 is split into two parts. In §2, we prove Theorem 1.1 for any orbispace presented by a simplicial complex of groups; this special case carries the essential topological content of the result. Due to one particular step in this proof, we have no explicit bound on the rank $n(d, m)$ of the vector bundles proven to exist. In §4, we deduce Theorem 1.1 in general by showing that every paracompact orbispace admits a representable map to a simplicial complex of groups. This result (Proposition 4.9) is likely of independent interest; it is the analogue of mapping a paracompact Hausdorff space to the nerve of an open covering using a partition of unity (it is, thus, worthy of note that its proof is not trivial).

The following immediate corollaries of Theorem 1.1 are derived in §5.

COROLLARY 1.2. *For X as in Theorem 1.1, we have $X = P/U(n)$ for a space P .*

COROLLARY 1.3. *Every paracompact smooth orbifold X of dimension $\leq d$ with isotropy groups of order $\leq m$ is the quotient $X = P/U(n)$ of a smooth manifold P by a smooth action of the compact Lie group $U(n)$, where $n = n(d, m)$.*

It seems that Theorem 1.1 does not help resolve the question of whether every separated Deligne–Mumford stack of finite type over \mathbb{C} is a global quotient; the question of whether the vector bundles produced by Theorem 1.1 on its analytification are algebraic (or even analytic) seems difficult.

In §6, we prove the following corollary of Theorem 1.1, to which the titular phrase ‘having enough vector bundles’ refers.

COROLLARY 1.4. *Let $X \rightarrow Y$ be a representable map of orbispaces satisfying the hypothesis of Theorem 1.1. Every vector bundle on X of bounded rank embeds into the pullback of a vector bundle of bounded rank on Y .*

It is well known that Corollary 1.4 is the key statement needed to show that the K -theory of finite rank vector bundles on orbispaces satisfies excision and exactness and is, thus, a cohomology theory (see [LO01, §3] and [Hen05, §6.3]). We elaborate on this assertion in §6, using the suggestive reformulation of Corollary 1.4 as the statement that pullback of vector bundles is *cofinal*. The K -theory of finite-rank vector bundles should agree (for reasonable orbispaces) with the other standard models of K -theory for orbispaces, such as using bundles of Fredholm operators [Seg70, Mat71, Ati69, AS69] or using orthogonal spectra [Sch18, §§6.3–6.4] (compare with Remark 1.6).

Remark 1.5. Another known (to experts) consequence of Corollary 1.4 (which we do not explain in detail) is that every paracompact quasi-smooth derived smooth orbifold with tangent and obstruction spaces of dimension $\leq d$ and isotropy groups of order $\leq m$ is the derived zero set of a smooth section of a vector bundle of rank $\leq n$ over a smooth orbifold of dimension $\leq n = n(d, m)$. (‘Quasi-smooth’ means locally isomorphic to the derived zero set of a smooth section of a smooth vector bundle over a smooth orbifold.)

Remark 1.6. Combined work of Schwede [Sch20] and Gepner and Henriques [HG07] established an equivalence between orthogonal spaces up to global equivalence (with respect to the ‘global family’ of all finite groups) and certain categories of ‘cellular’ topological stacks (which include, but are more general than, what we call orbispaces here). The vector bundles produced by Theorem 1.1 allow for a concrete description of the functor from orbispaces to orthogonal spaces (compare [Sch18, Definition 1.1.27]). Let X be an orbispace, and let E be any *faithful* vector bundle over X (meaning the fibers of E are faithful representations of the isotropy groups of X). The orthogonal space corresponding to X is given by

$$V \mapsto \text{Emb}_X(E, \underline{V}),$$

where $\text{Emb}_X(E, \underline{V})$ denotes the total space of the bundle of embeddings of E into V (note that $\text{Emb}_X(E, \underline{V})$ is a space because E is faithful).

Schwede [Sch18] also associates to every orthogonal spectrum a cohomology theory on orthogonal spaces, hence on orbispaces. Given an orthogonal spectrum A and an orbispace X which admits faithful vector bundles, the degree-zero A -cohomology of X is (in view of the above) given by the direct limit over vector bundles E over X of the set of homotopy classes of sections of the fibration $\Omega^E A(E) \rightarrow X$. More generally, we may consider the mapping spectrum $F(X, A)$ defined by

$$n \mapsto \varinjlim_{E/X} \Gamma(X, \Omega^E A(E \oplus \mathbb{R}^n))$$

whose stable homotopy groups are the A -cohomology groups of X . If A is a global Ω -spectrum [Sch18, Definition 4.3.8], then this direct limit is achieved at any faithful E , and the above definition of $F(X, A)$ is an Ω -spectrum. Let us also propose a possible definition of the A -homology groups of X as the stable homotopy groups of the spectrum

$$n \mapsto \varinjlim_{E/X} |\Omega^E A(E \oplus \mathbb{R}^n)|$$

where $|\cdot|$ indicates taking the coarse space of the total space of $\Omega^E A(E \oplus \mathbb{R}^n)$ over X .

Remark 1.7. It is natural to ask to what extent Theorem 1.1 may be generalized to the case of ‘Lie orbispaces’ (topological stacks locally modelled on Y/G for G a compact Lie group). The naive generalization is simply false: there are purely ineffective Lie orbispaces with isotropy group S^1 and coarse space S^3 which have no finite rank faithful vector bundles [Tot04, §2]. It is, however, reasonable to conjecture that the proof of Theorem 1.1 could be generalized to prove that the existence of enough vector bundles on a Lie orbispace is a purely cohomological question.

2. The main construction

This section is devoted to proving Theorem 1.1 for orbispaces which are presented by a simplicial complex of groups. This special case (stated as Theorem 2.13) carries the essential topological content of Theorem 1.1. For general background on orbispaces and topological stacks, we refer the reader to §3.

We begin by describing the basic idea of the proof, which we then implement in detail. Our orbispace X comes with a filtration by skeleta

$$\emptyset = X_{-1} \subseteq X_0 \subseteq X_1 \subseteq \dots,$$

where X_k is obtained from X_{k-1} by attaching cells of the form $(D^k, \partial D^k) \times \mathbb{B}G$ for finite groups G (here $\mathbb{B}G = */G$ is the orbispace quotient). We construct the desired vector bundle by induction on skeleta.

A direct implementation of this strategy runs immediately into the following obstruction. A (complex) vector bundle V on $\partial D^k \times \mathbb{B}G$ decomposes canonically as $V = \bigoplus_{\rho \in \hat{G}} V_\rho \otimes \rho$ for vector bundles V_ρ on ∂D^k indexed by the complex irreducible representations ρ of G . Thus, V is classified by an element of $\prod_{\rho \in \hat{G}} \pi_{k-1}(\prod_n BU(n))$, which vanishes if and only if V extends to $D^k \times \mathbb{B}G$. We seek to detect these obstructions using Chern characters. According to Bott periodicity, the homotopy groups of BU are given by

$$\pi_i(BU) = \begin{cases} \mathbb{Z} & i \text{ even,} \\ 0 & i \text{ odd,} \end{cases}$$

and the Chern character of a generator of $\pi_{2i}(BU)$ is nonzero. Now the Chern character of $V = \bigoplus_{\rho \in \hat{G}} V_\rho \otimes \rho$ is given by $\text{ch}(V) = \sum_{\rho \in \hat{G}} \text{ch}(V_\rho) \dim \rho$, so triviality of $\text{ch}(V)$ does not imply triviality of each $\text{ch}(V_\rho)$ or of the aforementioned obstructions to extending V .

To capture the information we need about V , we consider the more refined characteristic class which we call the *inertial Chern character*¹ ch^I (studied previously by Adem and Ruan [AR03]), which is a cohomology class on the *inertia stack*

$$IX := \text{Eq}(X \rightrightarrows X) = X \times_{X \times X} X.$$

In local coordinates $X = Y/G$, we have $IX = (\bigsqcup_{g \in G} Y^g)/G$. In such coordinates, the inertial Chern character is defined by recalling the Chern–Weil description of the usual Chern character $\text{ch}(V) = \text{tr} \exp(\Omega/(-2\pi i))$, for Ω the curvature of a hermitian connection on V , and writing

$$\text{ch}^I(V) = \text{tr} \left[g \exp \left(\frac{\Omega}{-2\pi i} \right) \right].$$

Now the inertia stack of $\partial D^k \times \mathbb{B}G$ is $\partial D^k \times G/G$ (quotient by the conjugation action), and the inertial Chern character of $V = \bigoplus_{\rho \in \hat{G}} V_\rho \otimes \rho$ is given by $\text{ch}^I(V) = \sum_{\rho \in \hat{G}} \text{ch}(V_\rho) \chi_\rho$. Thus, if $\text{ch}^I(V)$ is trivial, then so is each $\text{ch}(V_\rho)$ by linear independence of characters, and hence V extends to $D^k \times \mathbb{B}G$ as desired.

We are thus led to consider the modified problem of constructing a vector bundle on X with the desired fibers and with trivial inertial Chern character. It suffices to show that if X_{k-1} admits such a vector bundle, then so does X_k . The vanishing of $\text{ch}^I(V) \in H^*(IX_{k-1})$ guarantees that the obstructions to extending V to X_k vanish, by the above discussion. It thus remains to show that this extension can be taken to have trivial inertial Chern character. The inertial Chern character of any extension is an element of $\ker(H^*(IX_k) \rightarrow H^*(IX_{k-1}))$. By the long exact sequence

$$\dots \rightarrow H^*(IX_k, IX_{k-1}) \rightarrow H^*(IX_k) \rightarrow H^*(IX_{k-1}) \rightarrow \dots$$

it is thus in the image of $H^*(IX_k, IX_{k-1})$. The inertial Chern character is an even degree class, so we may assume that k is even. By modifying how we extend V from X_{k-1} to X_k by an element of $\prod_{\rho \in \hat{G}} \pi_k(BU)$ over a given k -simplex, we can shift its inertial Chern character by any integral linear combination of characters in $\text{Hom}(G/G, \mathbb{C}) \subseteq H^*(IX_k, IX_{k-1})$ (for G the isotropy group of that given k -simplex). By replacing V with $V^{\oplus a}$, we can multiply its inertial Chern character by any positive integer a . A combination of these two operations suffice to kill the inertial Chern character, provided that it is χ -rational (rational with respect to a certain \mathbb{Q} -structure on $H^*(IX)$ differing from the usual one). This rationality is not at all obvious given the transcendental definition we give of the inertial Chern character, but it is true, and thus the proof is complete.

¹ It is tempting to call it the ‘Chern character character’.

The remainder of this section is devoted to making the above outline precise. We begin by recalling the definition of a simplicial complex of groups (see also Haefliger [Hae91], Corson [Cor92], or Bridson and Haefliger [BH99]).

DEFINITION 2.1. A *simplicial complex of groups* is a pair (Z, G) consisting of a simplicial complex Z together with the following data.

- For every simplex $\sigma \subseteq Z$, a group G_σ .
- For every pair of simplices $\sigma \subseteq \tau$, an *injective* group homomorphism $G_\tau \hookrightarrow G_\sigma$.
- For every triple of simplices $\rho \subseteq \sigma \subseteq \tau$, an element of G_ρ conjugating the inclusion $G_\tau \hookrightarrow G_\rho$ to the composition of inclusions $G_\tau \hookrightarrow G_\sigma \hookrightarrow G_\rho$.
- For every quadruple of simplices $\pi \subseteq \rho \subseteq \sigma \subseteq \tau$, the resulting product of elements of G_π conjugating $G_\tau \hookrightarrow G_\pi$ to $G_\tau \hookrightarrow G_\rho \hookrightarrow G_\pi$ to $G_\tau \hookrightarrow G_\sigma \hookrightarrow G_\rho \hookrightarrow G_\pi$ and back to $G_\tau \hookrightarrow G_\pi$ must be the identity element of G_π .

Injectivity of each map $G_\tau \hookrightarrow G_\sigma$ will ensure that the geometric realization of (Z, G) is an orbispace, rather than some sort of more exotic topological stack.

A simplicial complex of groups (Z, G) presents an orbispace $\|(Z, G)\|$ called its *geometric realization*. A precise definition of this geometric realization is given in §4. For now, it will suffice to know that the coarse space of $\|(Z, G)\|$ is (the geometric realization of) Z itself, and that over the open star $\text{st}(\sigma) \subseteq Z$ of a simplex $\sigma \subseteq Z$, the geometric realization $\|(Z, G)\|$ is given by the orbispace quotient

$$\left(\bigsqcup_{\tau \supseteq \sigma} \tau \times (G_\tau \setminus G_\sigma) \right) / G_\sigma,$$

where the pieces for $\tau \supseteq \tau' \supseteq \sigma$ are glued together via the map $G_\tau \setminus G_\sigma \rightarrow G_{\tau'} \setminus G_\sigma$ induced by the map $G_\tau \rightarrow G_{\tau'}$ and the element of G_σ conjugating the composition $G_\tau \rightarrow G_{\tau'} \rightarrow G_\sigma$ to the map $G_\tau \rightarrow G_\sigma$.

We often abuse terminology and say ‘simplicial complex of groups’ when we really mean its geometric realization. Thus, we refer to $X = \|(Z, G)\|$ as a simplicial complex of groups. Its coarse space $|X|$ is the geometric realization of Z .

To apply the methods of differential topology to a given simplicial complex of groups, we fix a family of (germs of) smooth retractions $\tau \rightarrow \sigma$ for every pair of simplices $\sigma \subseteq \tau$ such that the maps $\tau \rightarrow \sigma \rightarrow \rho$ and $\tau \rightarrow \rho$ agree for $\rho \subseteq \sigma \subseteq \tau$ (such a family of smooth retractions may be constructed by induction). Given this data, objects of differential topology on τ (functions, differential forms, bundles, connections, etc.) are required to be pulled back under $\tau \rightarrow \sigma$ in a (neighborhood of every $\sigma \subseteq \tau$). For bundles, this requirement consists of the *data* of a compatible family of isomorphisms with the pullback bundles (one could equivalently consider only vector bundles built out of transition functions which satisfy the given pullback conditions).

In particular, the de Rham complex $\Omega^*(X)$ of a simplicial complex of groups X is defined, and it coincides with the de Rham complex of its coarse space $\Omega^*(|X|)$. There is a natural map $\Omega^*(|X|) \rightarrow C^*(|X|; \mathbb{R}) = C^*(X; \mathbb{R})$ from the de Rham complex to the simplicial cochain complex over \mathbb{R} , given by integrating differential forms over oriented simplices. It is a standard fact that this map is a quasi-isomorphism (proof: filter X by skeleta and invoke the five lemma to reduce to showing that $\Omega^*(\Delta^k, \partial\Delta^k) \rightarrow C^*(\Delta^k, \partial\Delta^k; \mathbb{R})$ is a quasi-isomorphism, which follows from the Poincaré lemma).

DEFINITION 2.2 (Chern character). Let X be a simplicial complex of groups, and let $V \rightarrow X$ be a complex vector bundle. Given a hermitian metric μ and hermitian connection θ on V , the

curvature of θ is a 2-form $\Omega(V, \theta)$ valued in $\mathfrak{u}(V, \mu) \subseteq \mathfrak{gl}(V) = \text{End}(V)$, and the Chern character form

$$\text{ch}(V, \theta) := \text{tr} \exp\left(\frac{\Omega(V, \theta)}{-2\pi i}\right) \in \Omega^{\text{even}}(X)$$

is closed. Its class in cohomology $\text{ch}(V) \in H^{\text{even}}(X; \mathbb{R})$ is called the Chern character of V . This class is independent of μ and θ by an interpolation argument (interpolate on $X \times [0, 1]$ between any two metric/connection pairs on $X \times 0$ and $X \times 1$, and use the fact that $H^*(X \times [0, 1]) = H^*(X)$).

For any stack X , its *inertia stack* is the stack

$$IX := \text{Eq}(X \rightrightarrows X) = X \times_{X \times X} X.$$

When $X = Y/G$, we have $IX = (\bigsqcup_{g \in G} Y^g)/G$. In particular, the local description of the geometric realization of a simplicial complex of groups provides a local description of its inertia stack as well. In fact, if X is a simplicial complex of groups then so is IX : a simplex of IX is a pair $(\sigma, [g])$ where $\sigma \subseteq X$ is a simplex and $[g] \subseteq G_\sigma$ is a conjugacy class, and $G_{(\sigma, [g])}$ is the centralizer of any element of the conjugacy class $[g] \subseteq G_\sigma$, etc.

DEFINITION 2.3 (Inertial Chern character). Let X be a simplicial complex of groups, and let $V \rightarrow X$ be a complex vector bundle. The inertial Chern character $\text{ch}^I(V) \in H^{\text{even}}(IX; \mathbb{C})$ is represented by the closed form

$$\text{ch}^I(V, \theta) := \text{tr} \left[g \exp\left(\frac{\Omega}{-2\pi i}\right) \right] \in \Omega^{\text{even}}(IX; \mathbb{C})$$

for any choice of hermitian metric and connection θ on V . Closedness can be seen by splitting the pullback of V to IX into isotypic pieces for the action of the cyclic group generated by g and observing that the contribution of each such piece is a (usual) Chern character form. Independence of the metric and connection follows from the same interpolation argument as before.

Example 2.4. If $X = \mathbb{B}G$, a vector bundle over X is simply a representation of G , the inertia stack $IX = G/G$ is the stack quotient of the conjugation action of G on itself, and the inertial Chern character $\text{ch}^I(V) : G/G \rightarrow \mathbb{C}$ is the character $g \mapsto \text{tr}(g|V)$ of V regarded as a representation of G .

Example 2.5. If $X = Y \times \mathbb{B}G$ for a smooth manifold Y and $V = \bigoplus_{\rho \in \hat{G}} V_\rho \otimes \rho$, then $IX = Y \times (G/G)$ and the inertial Chern character $\text{ch}^I(V) : G/G \rightarrow H^*(Y; \mathbb{C})$ is given by $\text{ch}^I(V) = \sum_\rho \text{ch}(V_\rho)\chi_\rho$. As the characters χ_ρ of the irreducible representations ρ of G form a basis for the space of maps $G/G \rightarrow \mathbb{C}$, we see that for $X = Y \times \mathbb{B}G$, the inertial Chern character determines (and is determined by) the Chern characters of each of the associated bundles V_ρ .

Example 2.6. The degree-zero part of the inertial Chern character $\text{ch}_0^I(V) \in H^0(IX; \mathbb{C})$ records the characters of the fibers of V , regarded as representations of the isotropy groups G_x of the points of X .

Despite admitting the aforementioned transcendental definition in terms of differential forms, the (usual) Chern character $\text{ch}(V) \in H^*(X; \mathbb{R})$ is well known to be rational, i.e. it lies in the subspace $H^*(X; \mathbb{Q}) \subseteq H^*(X; \mathbb{R})$. The inertial Chern character does not always lie in the subspace $H^*(IX; \mathbb{Q}) \subseteq H^*(IX; \mathbb{C})$, rather it is rational with respect to a different \mathbb{Q} -structure, which we now introduce.

DEFINITION 2.7 (χ -integral cohomology of IX). Regard the simplicial cochain group $C^*(IX; \mathbb{C})$ as the group of simplicial cochains on X with coefficients in $\text{Hom}(G_x/G_x, \mathbb{C})$. In this description, let us replace $\text{Hom}(G_x/G_x, \mathbb{C})$ with its subspace of integral (respectively, rational) linear combinations of characters of G_x . Equivalently, we consider the complex of simplicial cochains on IX which act on $\sigma \times (G_\sigma/G_\sigma) \subseteq IX$ (for any simplex $\sigma \subseteq X$) by an integral (respectively, rational) linear combination of characters of G_σ . We denote the resulting cohomology groups by $H^*(IX; \mathbb{Z}_\chi)$ (respectively, $H^*(IX; \mathbb{Q}_\chi)$), which are evidently functorial in X . We say that an element of $H^*(IX; \mathbb{C})$ is χ -integral (respectively, χ -rational) to mean that it lies in the image of $H^*(IX; \mathbb{Z}_\chi)$ (respectively, in the subspace $H^*(IX; \mathbb{Q}_\chi)$).

Example 2.8. If $X = Y \times \mathbb{B}G$, then $IX = Y \times (G/G)$ and $H^*(IX; \mathbb{Z}_\chi) = H^*(X; \mathbb{Z}[\hat{G}])$. Thus, $\text{ch}^I(V) = \sum_\rho \text{ch}(V_\rho)\chi_\rho$ is χ -rational because each $\text{ch}(V_\rho)$ is rational.

As the coefficient systems on X appearing in Definition 2.7 are finite free modules over \mathbb{Z} , we may dualize to define homology groups $H_*(IX; \mathbb{Z}_\chi)$ as well.

LEMMA 2.9. *There are canonical isomorphisms*

$$\begin{aligned} H^*(IX; \mathbb{C}) &= \text{Hom}(H_*(IX; \mathbb{Z}_\chi), \mathbb{C}), \\ H^*(IX; \mathbb{Q}_\chi) &= \text{Hom}(H_*(IX; \mathbb{Z}_\chi), \mathbb{Q}), \end{aligned}$$

and a canonical short exact sequence

$$0 \rightarrow \text{Ext}^1(H_{*-1}(IX; \mathbb{Z}_\chi), \mathbb{Z}) \rightarrow H^*(IX; \mathbb{Z}_\chi) \rightarrow \text{Hom}(H_*(IX; \mathbb{Z}_\chi), \mathbb{Z}) \rightarrow 0.$$

Proof. The complex $C_*(IX; \mathbb{Z}_\chi)$ is degreewise free, and the complexes $C^*(IX; \mathbb{Z}_\chi)$, $C^*(IX; \mathbb{Q}_\chi)$, and $C^*(IX; \mathbb{C})$ are obtained from it by applying the functors $\text{Hom}(-, \mathbb{Z})$, $\text{Hom}(-, \mathbb{Q})$, and $\text{Hom}(-, \mathbb{C})$, respectively. \square

PROPOSITION 2.10 (χ -rationality of ch^I). *Let X be a simplicial complex of groups of order $\leq m$. The inertial Chern character of any vector bundle V/X is χ -rational. In fact, there exists a positive integer $N = N(m, d, \dim V)$ such that $N \cdot \text{ch}_d^I(V)$ is χ -integral.*

It is worth remarking that the proof we give does not provide an explicit expression for $N(m, d, n)$ other than for $n = 1$.

Proof. We first show that the inertial Chern character of any line bundle L/X is χ -rational. We have $\text{tr}(ab) = \text{tr}(a)\text{tr}(b)$ for endomorphisms a, b of \mathbb{C} , so the inertial Chern character of a line bundle L splits as the product

$$\text{ch}^I(L) = \text{ch}_0^I(L) \exp(\text{ch}_1(L)),$$

where $\text{ch}_0^I(L) \in H^0(X; \mathbb{Z}_\chi)$ is the fiberwise character of L and $\text{ch}_1(L) \in H^2(X; \mathbb{C})$ is the Chern character in degree one. It thus suffices to show that $\text{ch}_1(L)$ is rational. We have $k \text{ch}_1(L) = \text{ch}_1(L^{\otimes k})$, so it suffices to show that $\text{ch}_1(L^{\otimes k})$ is integral for some positive integer k . As all isotropy groups of X have order $\leq m$, the tensor power $L^{\otimes m!}$ descends to a line bundle M on the coarse space. As M is an ordinary line bundle over a space (rather than an orbispace), its first Chern class $c_1(M) = \text{ch}_1(M)$ is integral. We have thus shown that $\text{ch}^I(L)$ is χ -rational (in fact, we have shown that $d!(m!)^d \text{ch}_d^I(L)$ is χ -integral).

To treat the case of vector bundles of dimension greater than one, we use the *splitting principle*. Given a vector bundle V/X with hermitian metric, let $\mathbb{F}(V) \rightarrow X$ denote the fibration whose fiber over $x \in X$ is the space of decompositions of V_x into ordered orthogonal one-dimensional subspaces. In other words, $\mathbb{F}(V) = P/U(1)^n$ where $P \rightarrow X$ is the principal

$U(n)$ -bundle associated to V/X with its chosen metric. We claim that the pullback map

$$H^*(IX; \mathbb{C}) \rightarrow H^*(IF(V); \mathbb{C})$$

is injective. The fiber of $IF(V) \rightarrow IX$ over a given point $(x, g) \in IX$ is the space of g -invariant ordered decompositions of V_x into one-dimensional subspaces. As the group generated by g is abelian, its irreducible representations are one-dimensional, and hence there are plenty of such decompositions which are g -invariant, namely when each one-dimensional subspace is contained in some g -isotypic piece of V_x . Thus, $IF(V) \rightarrow IX$ is a disjoint union of iterated projective space bundles, from which the desired injectivity statement follows (for any projective space bundle $\mathbb{P}(W) \rightarrow Z$ over an orbispace Z , the pullback map $H^*(Z) \rightarrow H^*(\mathbb{P}(W))$ is split by the map $\alpha \mapsto \int (c_1(L)^{\dim W - 1} \cup \alpha)$ where $L/\mathbb{P}(W)$ denotes the tautological line bundle and $\int : H^{*+2(\dim W - 1)}(\mathbb{P}(W)) \rightarrow H^*(Z)$ denotes fiberwise integration).

As $H^*(IX; \mathbb{C}) \rightarrow H^*(IF(V); \mathbb{C})$ is injective and $\text{Hom}_{\mathbb{Z}}(-, \mathbb{C})$ is exact, we conclude from Lemma 2.9 that $\text{Hom}(H_*(IX; \mathbb{Z}_{\chi})/H_*(IF(V); \mathbb{Z}_{\chi}), \mathbb{C}) = 0$, and hence that the domain is torsion. From this and Lemma 2.9 again, it follows that a class in $H^*(IX; \mathbb{C})$ is χ -rational if and only if its pullback to $H^*(IF(V); \mathbb{C})$ is χ -rational. Now consider the pullback of V to $\mathbb{F}(V)$. This pullback splits as a direct sum of line bundles, so because the inertial Chern character is additive under direct sum, we conclude that the pullback of $\text{ch}^I(V)$ to $IF(V)$ is χ -rational, and hence that $\text{ch}^I(V)$ is itself χ -rational.

It remains to produce an integer $N = N(m, d, n)$ such that $N \cdot \text{ch}_d^I(V)$ is χ -integral for $\dim V = n$. We claim that there exists a finite orbi-complex $\mathbb{B}_{m,d}U(n)$ carrying a principal $U(n)$ -bundle

$$\mathbb{E}_{m,d}U(n) \rightarrow \mathbb{B}_{m,d}U(n)$$

with the property that every principal $U(n)$ -bundle over a simplicial complex of groups X of dimension $\leq d$ and isotropy $\leq m$ is a pullback of $\mathbb{E}_{m,d}U(n) \rightarrow \mathbb{B}_{m,d}U(n)$. As the inertial Chern character of $\mathbb{E}_{m,d}U(n) \rightarrow \mathbb{B}_{m,d}U(n)$ is χ -rational and $\mathbb{B}_{m,d}U(n)$ is finite, there exists an integer $N = N(m, d, n)$ such that N times this inertial Chern character is χ -integral in cohomological degree d . By pullback, the same integer N works for any vector bundle of rank n over a simplicial complex of groups of order $\leq m$. Note that this argument gives no explicit bound on the integer N .

It remains to construct $\mathbb{E}_{m,d}U(n) \rightarrow \mathbb{B}_{m,d}U(n)$. We seek $U(n)$ -spaces $\mathbb{E}_{m,d}U(n)$ such that

$$\mathbb{E}_{m,-1}U(n) = \emptyset,$$

$$\mathbb{E}_{m,d}U(n) = \mathbb{E}_{m,d-1}U(n) \cup \left(\text{finitely many cells of the form } (D^d, \partial D^d) \times U(n)/G \right),$$

$$\pi_r((\mathbb{E}_{m,d}U(n))^G) = 0 \quad \text{for } r < d,$$

where $G \subseteq U(n)$ ranges over all finite subgroups of $U(n)$ of order $\leq m$. Note that there are finitely many conjugacy classes of such subgroups, and it suffices to consider just one representative of each conjugacy class. To show that $\mathbb{E}_{m,d}U(n)$ exists given $\mathbb{E}_{m,d-1}U(n)$, argue as follows. We have $\pi_{d-1}((\mathbb{E}_{m,d-1}U(n))^G) = H_{d-1}((\mathbb{E}_{m,d-1}U(n))^G)$ by Hurewicz, and the latter group is finitely generated because $\mathbb{E}_{m,d-1}U(n)$ is made up of finitely many cells. We define $\mathbb{E}_{m,d}U(n)$ by attaching cells $(D^d, \partial D^d) \times U(n)/G$ along a choice of finitely many generators of $\pi_{d-1}((\mathbb{E}_{m,d-1}U(n))^G)$, for each of the finitely many subgroups $G \subseteq U(n)$ on our fixed set of representatives. Note that $(D^d \times U(n)/H)^G = D^d \times \{a \in U(n) \mid a^{-1}Ga \subseteq H\}$, so any map $S^r \rightarrow (\mathbb{E}_{m,d}U(n))^G$ for $r < d$ can be homotoped to land inside $(\mathbb{E}_{m,d-1}U(n))^G$. It follows that the homotopy groups of $(\mathbb{E}_{m,d}U(n))^G$ satisfy the desired vanishing property.

Let us show that any principal $U(n)$ -bundle $P \rightarrow X$ is pulled back from $\mathbb{E}_{m,d}U(n) \rightarrow \mathbb{B}_{m,d}U(n)$ when $\dim X \leq d$ with isotropy groups of order $\leq m$. That is, we should construct an equivariant map $P \rightarrow \mathbb{E}_{m,d}U(n)$ (note that since the target is a space, this is the same as an equivariant map from the coarse space $|P|$). By induction on the cells of X , it suffices to solve the extension problem for equivariant maps from $(D^r, \partial D^r) \times U(n)/G$ to $\mathbb{E}_{m,d}U(n)$ for $r \leq d$ and $|G| \leq m$. This is equivalent to the extension problem for maps $(D^r, \partial D^r) \rightarrow (\mathbb{E}_{m,d}U(n))^G$, whose positive solution for $r \leq d$ is one of the defining properties of $\mathbb{E}_{m,d}U(n)$. \square

Example 2.11. A previous version of this text claimed a version of Proposition 2.10 with N independent of m . This stronger result is false; here is a counterexample. Begin with $\mathbb{B}(\mathbb{Z}/m)$ with a line bundle given by multiplication by $e^{2\pi i/m}$ on \mathbb{C} . Glue on a disk D^2 using an attaching map $\partial D^2 = S^1 \rightarrow \mathbb{B}(\mathbb{Z}/m)$ to a generator of \mathbb{Z}/m . The line bundle extends because all complex line bundles on a circle are trivial. The inertial Chern character of this line bundle has denominator m .

LEMMA 2.12 (Bott). *The image of the Chern character map*

$$\text{ch}_n : \pi_{2n}(BU) \rightarrow H^{2n}(S^{2n}; \mathbb{Q})$$

is precisely $H^{2n}(S^{2n}; \mathbb{Z})$.

Proof. This is an immediate corollary of Bott periodicity; for completeness, we include the proof from [Mat11]. Let η/S^2 denote the line bundle with $c_1(\eta) = 1$, and let 1 denote the trivial line bundle. According to Bott periodicity, multiplication with $\eta - 1 \in \tilde{K}^0(S^2)$ defines an isomorphism $\tilde{K}^0(X) \rightarrow \tilde{K}^0(\Sigma^2 X)$. In particular, $(\eta - 1)^{\otimes n} \in \tilde{K}^0(S^{2n})$ is a free generator. Multiplicativity of the Chern character gives $\text{ch}_n((\eta - 1)^{\otimes n}) = \text{ch}_1(\eta - 1)^n = 1^n = 1$. \square

We now have all the ingredients we need to prove the main result of this section, namely Theorem 1.1 for simplicial complexes of groups.

THEOREM 2.13. *Let X be a d -dimensional simplicial complex of groups of order $\leq m$. There exists a complex vector bundle V of rank $n = n(d, m) > 0$ over X , whose fiber over $x \in X$ is isomorphic to a direct sum of copies of the regular representation of G_x .*

Proof. The condition on the fibers of V amounts to the assertion that $\text{ch}_0^I(V) = n\chi_1$ where $\chi_1 \in H^0(IX; \mathbb{Q}_X)$ denotes ‘the characteristic function of the identity’ as a function $G_x/G_x \rightarrow \mathbb{C}$. We construct V satisfying

$$\text{ch}^I(V) = n\chi_1$$

(meaning all higher inertial Chern characters vanish). Certainly such a vector bundle exists over the 0-skeleton X_0 of X , with rank $m!$, because all isotropy groups have order $\leq m$. It therefore suffices to show that any such vector bundle on X_{k-1} extends to X_k .

To extend V as a vector bundle from X_{k-1} to X_k amounts to doing an extension from $\partial D^k \times \mathbb{B}G$ to $D^k \times \mathbb{B}G$ for each k -simplex. The obstruction to doing this lies in $\prod_{\rho \in \hat{G}} \pi_{k-1}(BU(n_\rho))$, where $n_\rho = (\dim V)(\dim \rho)/|G|$. The map $BU(n_\rho) \rightarrow BU$ is an isomorphism on π_{k-1} provided $k - 1 < 2n_\rho + 2$, which is guaranteed by taking say $\dim V > dm$. The Chern character detects $\pi_*(BU)$ by Lemma 2.12, so vanishing of the inertial Chern character and linear independence of characters means that these obstructions all vanish. Thus, V extends to X_k .

It remains to show that the extension of V to X_k can be taken to have trivial inertial Chern character. The space of extensions over a given k -simplex is a torsor for $\prod_{\rho \in \hat{G}} \pi_k(BU(n_\rho))$, which is again $\prod_{\rho \in \hat{G}} \pi_k(BU)$ once we take $\dim V > dm$. We use this freedom to ensure that the inertial

Chern character of V on X_k vanishes. Consider the long exact sequence

$$\cdots \rightarrow H^*(IX_k, IX_{k-1}; \mathbb{Q}_\chi) \rightarrow H^*(IX_k; \mathbb{Q}_\chi) \rightarrow H^*(IX_{k-1}; \mathbb{Q}_\chi) \rightarrow \cdots .$$

As the inertial Chern character in $H^*(IX_k; \mathbb{Q}_\chi)$ maps to zero in $H^*(IX_{k-1}; \mathbb{Q}_\chi)$, it is in the image of $H^*(IX_k, IX_{k-1}; \mathbb{Q}_\chi)$. In particular, it may be nonzero only in degree k . Let us now replace V with $V^{\oplus a}$ (which multiplies its inertial Chern character by a) for an appropriate positive integer a so that by Proposition 2.10 its inertial Chern character is χ -integral. Modifying our bundle by an element of (the product over the k -simplices of) $\prod_{\rho \in \hat{G}} \pi_k(BU)$ shifts its inertial Chern character by anything in $H^k(IX_k, IX_{k-1}; \mathbb{Z}_\chi)$ by Lemma 2.12. We are done because $H^k(IX_k, IX_{k-1}; \mathbb{Z}_\chi) \rightarrow H^k(IX_k; \mathbb{Z}_\chi)$ is surjective by the long exact sequence. \square

Remark 2.14. One may interpret the proof of Theorem 2.13 in homotopy theoretic terms as follows. Vector bundles are classified by maps to a classifying space $\mathbb{B}U(n)$, and because $\mathbb{B}U(n)$ is not contractible, the extension problem for vector bundles has nontrivial obstructions. Vector bundles with rationally trivialized inertial Chern character are classified by maps to the total space of a fibration over $\mathbb{B}U(n)$ whose fiber classifies odd-dimensional rational cohomology classes on the inertia stack. Our observation that the obstructions to this new problem are torsion is essentially the observation that this total space is rationally contractible. The definition of this fibration over $\mathbb{B}U(n)$ depends on the additivity of the Chern character (note that $\mathbb{E}U(n) \rightarrow \mathbb{B}U(n)$ is not suitable for this argument because $\mathbb{E}U(n)$ is a space, hence every bundle pulled back from $\mathbb{E}U(n)$ has trivial isotropy representations). It may prove interesting to interpret this argument within Schwede’s framework of global homotopy theory [Sch18].

3. Topological stacks

We review some basic facts about stacks (on the category of topological spaces), we give a precise definition of what we mean by an ‘orbispace’, and we establish some of their basic properties. For further background, the reader may wish to consult Noohi [Noo05], Gepner and Henriques [HG07], Behrend [Beh14], Metzler [Met03], Behrend and Noohi [BN06], Heinloth [Hei05], or Laumon and Moret-Bailly [LMB00].

Let \mathbf{Top} denote the category of topological spaces and continuous maps, and let \mathbf{Grpd} denote the 2-category of (essentially) small groupoids. A *stack* is a functor $F : \mathbf{Top}^{\text{op}} \rightarrow \mathbf{Grpd}$ which satisfies *descent*, i.e. such that for every topological space U and every open cover $\{U_i \rightarrow U\}_i$, the natural functor

$$F(U) \rightarrow \text{Eq} \left[\prod_i F(U_i) \rightrightarrows \prod_{i,j} F(U_i \cap U_j) \rightrightarrows \prod_{i,j,k} F(U_i \cap U_j \cap U_k) \right]$$

is an equivalence. Stacks form a 2-category, with morphisms given by natural transformations of functors. The 2-category of stacks is complete, meaning all (small) limits exist; furthermore, these limits may be calculated pointwise in the sense that $(\lim_\alpha F_\alpha)(U) = \lim_\alpha (F_\alpha(U))$. Note that, as we are working in a 2-categorical context, all functors are 2-functors, all diagrams are 2-diagrams, all limits are 2-limits, etc. (though we usually omit the prefix ‘2-’).

The Yoneda lemma implies that the Yoneda functor $X \mapsto \text{Hom}(-, X)$ embeds the category of topological spaces fully faithfully into the 2-category of stacks, and moreover that the natural map from $F(X)$ to the groupoid of maps of stacks $\text{Hom}(-, X) \rightarrow F(-)$ is an equivalence. The category of topological spaces is complete, and the Yoneda embedding is continuous (commutes with limits). Hence, we make no distinction between a topological space X and the associated

stack $\text{Hom}(-, X)$ of maps to X , nor between objects of $F(X)$ and maps $\text{Hom}(-, X) \rightarrow F(-)$ (which we simply write as $X \rightarrow F$).

Every stack X has a *coarse space* $|X|$ (a topological space) which is initial in the category of maps from X to topological spaces. Concretely, the points of $|X|$ are the isomorphism classes of maps $* \rightarrow X$, and a subset $U \subseteq |X|$ is open if and only if for every map $Y \rightarrow X$ from a topological space Y , the inverse image of U is an open subset of Y .

A stack is called *representable* if and only if it is in the essential image of the Yoneda embedding (i.e. it is isomorphic to a topological space). A morphism of stacks $F \rightarrow G$ is called representable if and only if for every map $X \rightarrow G$ from a topological space X , the fiber product $F \times_G X$ is representable.

For any property \mathcal{P} of morphisms of topological spaces which is preserved under pullback, a representable morphism of stacks $F \rightarrow G$ is said to have property \mathcal{P} if and only if the pullback $F \times_G X \rightarrow X$ has \mathcal{P} for every map $X \rightarrow G$ from a topological space X . The following are examples of properties \mathcal{P} of morphisms $f : X \rightarrow Y$ which are preserved under pullback:

- f is *injective*;
- f is *surjective*;
- f is *open*, meaning that the image of any open set is open. In contrast, the property of being *closed*, meaning that the image of any closed set is closed, is *not* preserved under pullback;
- f is an *embedding*, meaning that it is a homeomorphism onto its image;
- f is a *closed embedding*;
- f is *étale*, meaning that for every $x \in X$ there exists an open neighborhood $x \in U \subseteq X$ such that $f|_U : U \rightarrow Y$ is an open embedding;
- f is *separated*, meaning that for every distinct pair $x_1, x_2 \in X$ with $f(x_1) = f(x_2)$, there exist open neighborhoods $x_i \in U_i \subseteq X$ which are disjoint $U_1 \cap U_2 = \emptyset$ (this is equivalent to the relative diagonal $X \rightarrow X \times_Y X$ being a closed embedding);
- f is *universally closed*, meaning that $X \times_Y Z \rightarrow Z$ is closed for every $Z \rightarrow Y$; this is equivalent to the assertion that for every $y \in Y$ and every collection of open sets $\{U_i \subseteq X\}_i$ covering $f^{-1}(y)$, there exists a finite subcollection which covers $f^{-1}(V)$ for some open neighborhood $y \in V \subseteq Y$ (one proof of this equivalence goes via yet a third equivalent condition, namely that every net $\{x_\alpha \in X\}_\alpha$ with $f(x_\alpha) \rightarrow y$ has a subnet converging to some $x \in f^{-1}(y)$);
- f is *proper*, meaning that it is separated and universally closed;
- f *admits local sections*, meaning that there is an open cover $\{U_i \subseteq Y\}_i$ such that every restriction $f|_{f^{-1}(U_i)} : f^{-1}(U_i) \rightarrow U_i$ admits a section;
- f is a *finite covering space*, meaning that there is an open cover $\{U_i \subseteq Y\}_i$ such that every restriction $f|_{f^{-1}(U_i)} : f^{-1}(U_i) \rightarrow U_i$ is isomorphic to $U_i^{\sqcup n_i} \rightarrow U_i$ for some integer $n_i \geq 0$.

Each of these properties \mathcal{P} is also closed under composition, and thus also under fiber products, meaning that for maps $X \rightarrow X'$ and $Y \rightarrow Y'$ over Z , if both $X \rightarrow X'$ and $Y \rightarrow Y'$ have \mathcal{P} , then $X \times_Z Y \rightarrow X' \times_Z Y'$ also has \mathcal{P} (indeed, $X \times_Z Y \rightarrow X \times_Z Y'$ is a pullback of $Y \rightarrow Y'$).

For any stack X , open (respectively, closed) embeddings $Y \hookrightarrow X$ are in natural bijection with open (respectively, closed) subsets of $|X|$.

To check that a given map of spaces satisfies one of the properties \mathcal{P} above, it is often helpful to make use of the fact that these \mathcal{P} are all *local on the target*, meaning that for every open cover $\{U_i \subseteq Y\}_i$, if $X \times_Y U_i \rightarrow U_i$ has \mathcal{P} for every i , then so does $X \rightarrow Y$. This leads to the following generalization for maps of stacks: if $F \rightarrow G$ is a representable morphism of stacks and $G' \rightarrow G$ is a representable morphism of stacks admitting local sections, then $F \rightarrow G$ has \mathcal{P} if and only if $F \times_G G' \rightarrow G'$ has \mathcal{P} . In fact, in this statement we need not assume that $G' \rightarrow G$ is

representable, just that it admit local sections in the generalized sense that for every map $X \rightarrow G$ from a topological space X , there exists an open cover $\{U_i \subseteq X\}_i$ such that each $G' \times_G U_i \rightarrow U_i$ admits a section. We thus say that ‘ \mathcal{P} descends along maps admitting local sections’. The same descent property holds for representability itself.

LEMMA 3.1 (Representability descends under maps admitting local sections). *Let $F \rightarrow G$ be a map of stacks, and let $G' \rightarrow G$ be a map of stacks admitting local sections. If $F \times_G G' \rightarrow G'$ is representable, then so is $F \rightarrow G$.*

Proof. By replacing $F \rightarrow G$ and $G' \rightarrow G$ with their pullbacks under $U \rightarrow G$ for a topological space U , we may assume without loss of generality that G is representable. As $G' \rightarrow G$ admits local sections, we may replace $G' \rightarrow G$ with the composition $\bigsqcup_i U_i \rightarrow G' \rightarrow G$ where $\{U_i \subseteq G'\}_i$ is an open cover. Now each $F \times_G U_i$ is representable by assumption, and gluing these spaces together on their common overlaps $F \times_G (U_i \cap U_j)$ gives a topological space representing F . \square

A complex vector bundle over a stack X is a representable map $V \rightarrow X$ together with maps $V \times_X V \rightarrow V$ and $\mathbb{C} \times V \rightarrow V$ (both over X) such that for every map $U \rightarrow X$ from a topological space U , there exists an open cover $\{U_i \subseteq U\}_i$ and integers $n_i \geq 0$ such that $V \times_X U_i \rightarrow U_i$ is isomorphic to $\mathbb{C}^{n_i} \times U_i \rightarrow U_i$ equipped with its fiberwise addition and scaling maps. A pullback of a complex vector bundle is naturally a complex vector bundle.

The class of so called *topological stacks* (those which admit a presentation via a topological groupoid) are somewhat better behaved than general stacks. A *topological groupoid* $M \rightrightarrows O$ consists of a pair of topological spaces O (‘objects’) and M (‘morphisms’), two maps $M \rightrightarrows O$ (‘source’ and ‘target’), a map $O \rightarrow M$ (‘identity’), an involution $M \rightarrow M$ (‘inverse’), and a map $M \times_O M \rightarrow M$ (‘composition’) satisfying the axioms of a groupoid. A topological groupoid $M \rightrightarrows O$ presents a stack $[M \rightrightarrows O]$ defined as follows. An object of $[M \rightrightarrows O](X)$ is an open cover $\{U_i \subseteq X\}_i$ together with maps $U_i \rightarrow O$ and $U_i \cap U_j \rightarrow M$ satisfying a compatibility condition, and an isomorphism in $[M \rightrightarrows O](X)$ consists of maps $U_i \cap U'_j \rightarrow M$ satisfying a compatibility condition. There is a natural map $O \rightarrow [M \rightrightarrows O]$ (take the trivial open cover $\{X \subseteq X\}$) and the fiber product $O \times_{[M \rightrightarrows O]} O$ is naturally identified with M . The morphism $O \rightarrow [M \rightrightarrows O]$ admits local sections (by definition), so because $O \times_{[M \rightrightarrows O]} O \rightarrow O$ is representable, it follows by descent that $O \rightarrow [M \rightrightarrows O]$ is representable. Conversely, a representable map $U \rightarrow X$ admitting local sections from a topological space U to a stack X determines a topological groupoid $U \times_X U \rightrightarrows U$ presenting X . Indeed, the fiber product $U \times_X U$ is representable (because U and $U \rightarrow X$ are), it admits two maps to U (the two projections), an involution (exchanging the two factors), and a composition map $(U \times_X U) \times_U (U \times_X U) = U \times_X U \times_X U \rightarrow U \times_X U$ (forgetting the middle factor), and one can check using the stack property that the natural map $[U \times_X U \rightrightarrows U] \rightarrow X$ is an equivalence. A stack X for which there exists a representable map $U \rightarrow X$ admitting local sections from a space U is called *topological*, and such $U \rightarrow X$ is called an *atlas* for X .

For a topological stack X with atlas $U \rightarrow X$, for any property \mathcal{P} which descends along maps admitting local sections, the map $U \rightarrow X$ has \mathcal{P} if and only if both maps $U \times_X U \rightarrow U$ have \mathcal{P} , and the diagonal $X \rightarrow X \times X$ has \mathcal{P} if and only if the map $U \times_X U \rightarrow U \times U$ has \mathcal{P} . In particular, for any topological stack X , the diagonal $X \rightarrow X \times X$ is representable, and thus every map $Z \rightarrow X$ from a topological space Z is representable. More generally, for any map of topological stacks $X \rightarrow Y$, the relative diagonal $X \rightarrow X \times_Y X$ is representable (by descent from its pullback $V \times_X V \rightarrow V \times_Y V$ for an atlas $V \rightarrow X$). If $X \rightarrow Y$ is representable and Y is topological, then so is X (if $U \rightarrow Y$ is an atlas for Y , then its pullback $V = U \times_Y X \rightarrow X$ is an atlas for X), and the relative diagonal $X \rightarrow X \times_Y X$ has \mathcal{P} if and only if the relative diagonal $V \rightarrow V \times_U V$ of atlases has \mathcal{P} (the latter is the pullback of the former under the map

$V \times_U V \rightarrow X \times_Y X$, which is representable and admits local sections because it is a pullback of $U \rightarrow Y$).

For a topological space V , a topological group G , and a continuous group action $G \curvearrowright V$, we may consider the action groupoid $G \times V \rightrightarrows V$ with source and target maps $(g, x) \mapsto x$ and $(g, x) \mapsto gx$. The stack associated to this groupoid is denoted V/G and is called the stack quotient of the action $G \curvearrowright V$. If G is discrete, the two maps $G \times V \rightarrow V$ are étale, so $V \rightarrow V/G$ is étale. If G is compact and V is Hausdorff, the map $G \times V \rightarrow V \times V$ is universally closed (factor as $G \times V \rightarrow G \times V \times V$ which is a closed embedding because V is Hausdorff and $G \times V \times V \rightarrow V \times V$ which is universally closed because G is compact), so V/G has universally closed diagonal. If G is Hausdorff, then the map $G \times V \rightarrow V \times V$ is separated, so V/G has separated diagonal. Thus, if G is compact Hausdorff and V is Hausdorff, then V/G has proper diagonal.

A groupoid presentation of a topological stack X also gives a description of its coarse space $|X|$ as follows. For an atlas $U \rightarrow X$, consider the equivalence relation \sim_X on U given by the image of $U \times_X U \rightarrow U \times U$. There is a map $X \rightarrow U/\sim_X$ (which is tautological once we regard X as $[U \times_X U \rightrightarrows U]$), inducing a map $|X| \rightarrow U/\sim_X$, which is a bijection, essentially by definition. As an open substack of X pulls back to an open subset of U invariant under \sim_X , it follows that $|X| \xrightarrow{\sim} U/\sim_X$ is open and is, thus, a homeomorphism. In particular, it follows that the coarse space $|V/G|$ of the stack quotient V/G is the usual topological quotient of V by G .

An action $G \curvearrowright V$ is called *locally trivial* if and only if V admits a cover by G -invariant open sets $\{G \times Z_i \subseteq V\}_i$ where $G \curvearrowright G \times Z_i$ acts by left multiplication on G and trivially on Z_i . If $G \curvearrowright V$ is locally trivial, then the natural map from the stack quotient to the topological quotient $V/G \rightarrow |V/G|$ is an equivalence. Indeed, this assertion is local on $|V/G|$, so it suffices to consider the case of $G \curvearrowright G \times Z$, where it holds by inspection.

DEFINITION 3.2. A (*separated*) *orbispace* is a stack X for which there exists a representable étale surjection $U \rightarrow X$ from a topological space U (an ‘étale atlas’), and the diagonal $X \rightarrow X \times X$ is proper.

PROPOSITION 3.3. A stack X is an orbispace if and only if $|X|$ is Hausdorff and there exists an open cover $\{V_i/G_i \subseteq X\}_i$ where G_i are finite discrete groups acting on Hausdorff spaces V_i .

(Similar results include [Beh14, Theorem 1.108] and [Noo05, Proposition 14.10].)

Proof. Let X be an orbispace, and let us show that there is an open cover $\{V_i/G_i \subseteq X\}_i$. Fix an étale atlas $U \rightarrow X$, and let $u \in U$. The automorphism group $G_u := \{u\} \times_X \{u\} \subseteq U \times_X U$ is finite and discrete because $X \rightarrow X \times X$ is proper. As the two projections $U \times_X U \rightrightarrows U$ are étale, for every $g \in G$ there exists an open neighborhood $g \in U_g \subseteq U \times_X U$ such that each projection restricted to U_g is an open embedding (this gives another proof that G has the discrete topology). As G is finite and $U \times_X U \rightarrow U \times U$ is separated, we may take these U_g to be disjoint. Now the complement of $\bigsqcup_g U_g$ is closed and disjoint from G , so it projects to a closed (by properness) subset of $U \times U$ disjoint from (u, u) . Hence, there is an open neighborhood $u \in V \subseteq U$ such that $V \times_X V \subseteq \bigsqcup_g U_g$. Thus, $V \times_X V$ is a disjoint union of pieces indexed by G , and each piece maps homeomorphically to V under each projection. By further shrinking V , we may assume that the map $V \times_X V \rightarrow G$ respects composition (this is possible because composition is continuous). It follows that $V \times_X V \rightrightarrows V$ is an action groupoid $G \times V \rightrightarrows V$ for an action $G \curvearrowright V$. As $V = \{1\} \times V \hookrightarrow G \times V = V \times_X V \rightarrow V \times V$ expresses the diagonal of V as a composition of proper maps, we conclude that V is Hausdorff. Now the map of groupoids $(G \times V \rightrightarrows V) = (V \times_X V \rightrightarrows V) \rightarrow (U \times_X U \rightrightarrows U)$ induces a map of stacks $V/G \rightarrow X$. To see that this is an open embedding, let $V^+ \subseteq U$ denote

the orbit of V under the morphisms $U \times_X U \rightrightarrows U$. As the projections $U \times_X U \rightrightarrows U$ are étale, it follows that $V^+ \subseteq U$ is open, and hence $[V^+ \times_X V^+ \rightrightarrows V^+] \rightarrow [U \times_X U \rightrightarrows U]$ is an open embedding of stacks; denote this open substack by $Z \subseteq X$, so $V \rightarrow Z$ is an étale atlas. Thus, the topological groupoid $V \times_X V = V \times_Z V \rightrightarrows V$ presents Z , so $V/G \rightarrow X$ is an open embedding as desired.

Let us now show that for any orbispace X , its coarse space $|X|$ is Hausdorff. We saw earlier that $|X|$ is the quotient of U by the image of $U \times_X U \rightarrow U \times U$ (which is an equivalence relation). This equivalence relation is closed because $U \times_X U \rightarrow U \times U$ is proper, so $|X|$ is Hausdorff provided the quotient map $U \rightarrow |X|$ is open. Now openness of $U \rightarrow |X|$ does not depend on which atlas $U \rightarrow X$ we take: there are étale maps $U \leftarrow U \times_X U' \rightarrow U'$ over $|X|$, which means that $U \rightarrow |X|$ is open if and only if $U' \rightarrow |X|$ is open. Moreover, openness of $U \rightarrow |X|$ can be checked locally on $|X|$, so we may assume without loss of generality that $X = V/G$. Hence, it is enough to note that the quotient map $V \rightarrow |V/G|$ (induced by the canonical étale atlas $V \rightarrow V/G$) is open. Thus, $|X|$ is Hausdorff.

Finally, let us show that if $|X|$ is Hausdorff and there is an open cover $\{V_i/G_i \subseteq X\}_i$, then X is an orbispace. The maps $V_i \rightarrow V_i/G_i$ are representable étale, so $U := \bigsqcup_i V_i \rightarrow X$ is an étale atlas. To show that the diagonal $X \rightarrow X \times X$ is proper, it is equivalent to show that $U \times_X U \rightarrow U \times U$ is proper. This reduces to showing that $V \times_X V' \rightarrow V \times V'$ is proper for any pair $V/G \hookrightarrow X \leftarrow V'/G'$. As $|X|$ is Hausdorff, the map $V \times_{|X|} V' \rightarrow V \times V'$ a closed embedding (as it is a pullback of the diagonal of $|X|$), so it follows that $V \times_X V' \rightarrow V \times V'$ is proper if and only if $V \times_X V' \rightarrow V \times_{|X|} V'$ is proper. Now for the purposes of studying the latter map $V \times_X V' \rightarrow V \times_{|X|} V'$, we may as well shrink V, V' , and X so that $V/G = X = V'/G'$. Now the diagonal of $V/G = V'/G' = X$ is proper, hence so is $V \times_X V' \rightarrow V \times V'$. \square

COROLLARY 3.4. *Every orbispace X has an étale atlas $U \rightarrow X$ with U Hausdorff; equivalently, every étale atlas $U \rightarrow X$ has U locally Hausdorff.*

Proof. By Proposition 3.3 there is an open cover $\{V_i/G_i \subseteq X\}_i$ with V_i Hausdorff, so $U := \bigsqcup_i V_i \rightarrow X$ is an étale atlas with U Hausdorff. Now for any two étale atlases $U, U' \rightarrow X$, consideration of the surjective étale maps $U \leftarrow U \times_X U' \rightarrow U'$ shows that U is locally Hausdorff if and only if U' is locally Hausdorff. Given any étale atlas $U \rightarrow X$ with U locally Hausdorff and any open cover $\{U_i \subseteq U\}_i$ with U_i Hausdorff, the disjoint union $U' := \bigsqcup_i U_i \rightarrow X$ is an étale atlas with U' Hausdorff. \square

COROLLARY 3.5. *For any étale atlas $U \rightarrow X$ on an orbispace, there exists an open cover $\{V_i/G_i \subseteq X\}_i$ as in Proposition 3.3 such that each map $V_i \rightarrow X$ factors through an open embedding $V_i \rightarrow U$.*

Proof. Let $U \rightarrow X$ be given and fix any open cover $\{V_i/G_i \subseteq X\}_i$ as in Proposition 3.3. As $U \times_X V_i \rightarrow V_i$ is étale, it admits local sections, and hence by replacing each V_i with an open cover of itself, we may assume that each projection $U \times_X V_i \rightarrow V_i$ admits a section (which is thus an open embedding). Now the resulting maps $V_i \rightarrow U$ are étale by the factorization $V_i \rightarrow U \times_X V_i \rightarrow U$, so by again replacing each V_i with an open cover, we may assume they are open embeddings.

Alternatively, we could note that the open cover $\{V_i/G_i \subseteq X\}_i$ produced by the proof of Proposition 3.3 is in fact of the desired form. \square

COROLLARY 3.6. *A map of orbispaces $X \rightarrow Y$ is representable if and only if it is injective on isotropy groups. In particular, an orbispace X is a space if and only if it has trivial isotropy.*

Proof. Any representable morphism of stacks is injective on isotropy groups (just test against points). Thus, we are left with showing that a map of orbispaces $X \rightarrow Y$ which is injective on isotropy groups is representable.

As representability descends under maps admitting local sections, it suffices to show that the fiber product $X' = X \times_Y Y'$ is representable for some étale atlas $Y' \rightarrow Y$. Note that X' has trivial isotropy because Y' has trivial isotropy and $X' \rightarrow Y'$ is injective on isotropy groups (being a pullback of $X \rightarrow Y$).

We claim that X' is an orbispace, provided we take Y' to be Hausdorff (which we can by Corollary 3.4). The pullback of an étale atlas $U \rightarrow X$ is an étale atlas $U' \rightarrow X'$ (because $X' \rightarrow X$ is representable, being a pullback of $Y' \rightarrow Y$). The diagonal of X' is the composition $X' \rightarrow X' \times_X X' \rightarrow X' \times X'$. The second map $X' \times_X X' \rightarrow X' \times X'$ is proper as it is a pullback of $X \rightarrow X \times X$. To analyze the first map $X' \rightarrow X' \times_X X'$, note that it pulls back to $U' \rightarrow U' \times_U U'$ under the map $U' \times_U U' \rightarrow X' \times_X X'$ which admits local sections (being a pullback of $U \rightarrow X$). As $Y \rightarrow Y \times Y$ is separated, its pullback $Y' \times_Y Y' \rightarrow Y' \times Y'$ is also separated, which implies each projection $Y' \times_Y Y' \rightrightarrows Y'$ is separated because Y' is Hausdorff, which implies $Y' \rightarrow Y$ is separated. Hence, its pullback $U' \rightarrow U$ is separated, so $U' \rightarrow U' \times_U U'$ is a closed embedding, hence proper. Thus, $X' \rightarrow X' \times_X X'$ is proper, and we conclude that X' is an orbispace.

We are thus reduced to showing that an orbispace Z with trivial isotropy is a space. By Proposition 3.3, we know that Z is given locally by V/G for V Hausdorff and G finite discrete acting freely (because Z has trivial isotropy). Free actions $G \curvearrowright V$ with V Hausdorff and G finite are locally trivial, so we conclude that the map $Z \rightarrow |Z|$ is an equivalence. Alternatively, we could note that the chart V/G near a given $x \in X$ constructed in the proof of Proposition 3.3 in fact satisfies $G = G_x$ by definition. □

4. Coverings and nerves

We show how Theorem 2.13 implies Theorem 1.1. It is enough to show that a given orbispace admits a representable map to a simplicial complex of groups; this is Proposition 4.9. This simplicial complex of groups is basically just the nerve of a suitable open cover, however its construction is somewhat more delicate than one might initially expect.

A *sieve* on a topological space X is a subset $S \subseteq 2^X$ consisting of open sets such that $U' \subseteq U \in S$ implies $U' \in S$. A *covering sieve* on X is a sieve S such that $\bigcup_{U \in S} U = X$. An open cover $\{U_i \subseteq X\}_i$ is said to *generate* the covering sieve on X consisting of those open sets which are contained in some U_i .

DEFINITION 4.1. A *connection sieve* on a map of spaces $f : X \rightarrow Y$ is a covering sieve S on X such that (1) for $U \in S$, the composition $U \rightarrow X \rightarrow Y$ is an open embedding, and (2) for $U, V \in S$ with $f(U) = f(V)$, either $U = V$ or $U \cap V = \emptyset$.

Note that for sieves satisfying condition (1), condition (2) is equivalent to condition (2') for $U, V \in S$ either $U \cap V = \emptyset$ or $f(U \cap V) = f(U) \cap f(V)$. If $S' \subseteq S$ is an inclusion of covering sieves and S is a connection sieve on $X \rightarrow Y$, then so is S' . In particular, if S and S' are connection sieves on $X \rightarrow Y$, then so is $S \cap S'$. To check that an open cover $\{U_i \subseteq X\}_i$ generates a connection sieve, it is enough to check axioms (1) and (2') for the open sets U_i .

DEFINITION 4.2. A map $X \rightarrow Y$ is called *strongly étale* if and only if it admits a connection sieve.

Open embeddings are strongly étale, and strongly étale maps are separated and étale (however, the converse is false by Example 4.3). Being strongly étale is preserved under pullback

(take the connection sieve generated by the pullback of the original connection sieve), and the class of strongly étale maps is closed under composition (a connection sieve $S_{X/Z}$ for a composition $X \rightarrow Y \rightarrow Z$ is given by those elements of a fixed connection sieve $S_{X/Y}$ for $X \rightarrow Y$ whose image lies in a fixed connection sieve $S_{Y/Z}$ for $Y \rightarrow Z$). A disjoint union $\bigsqcup_i X_i \rightarrow \bigsqcup_i Y_i$ of strongly étale maps $X_i \rightarrow Y_i$ is strongly étale (take disjoint union of connection sieves), and the projection $A \times Y \rightarrow Y$ is strongly étale for any discrete space A . Being strongly étale is *not* local on the target, as shown by the following example.

Example 4.3. Let us construct a finite covering space which is not strongly étale. Let $Z = \{1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\} \cup \{0\} \subseteq \mathbb{R}$ with the subspace topology. Every double cover of Z or $Z \setminus 0$ is trivial. A given double cover of $Z \setminus 0$ has, however, many distinct extensions to Z , indexed by functions $Z \setminus 0 \rightarrow \mathbb{Z}/2$ modulo those functions which extend continuously to zero.

Now suppose given a double cover of Z together with a connection sieve on it. The restriction of this data to $Z \setminus 0$ remembers the extension of the double cover to Z (use the elements of the connection sieve projecting to open sets of the form $Z \cap (0, \varepsilon)$).

Let $\tilde{Z}_\alpha^+ \rightarrow Z^+$ denote the double cover obtained by gluing two copies of $Z \sqcup Z \rightarrow Z$ over $Z \setminus 0$ via some map $\alpha : Z \setminus 0 \rightarrow \mathbb{Z}/2$. If $\tilde{Z}_\alpha^+ \rightarrow Z^+$ is strongly étale, then we can take any connection sieve on it, restrict to the common $Z \setminus 0 \subseteq Z^+$, and deduce that the two extensions of the double cover of $Z \setminus 0$ to Z coincide, in other words that α extends continuously to zero. Hence, the double cover $\tilde{Z}_\alpha^+ \rightarrow Z^+$ is strongly étale if and only if α extends continuously to zero.

Recall that a topological space is called *paracompact* if and only if every open cover admits a locally finite refinement [Die44]. If X is paracompact and Hausdorff, then there exists a *partition of unity* subordinate to any given locally finite open cover $\{U_i \subseteq X\}_i$, namely functions $f_i : X \rightarrow \mathbb{R}_{\geq 0}$ with $\text{supp } f_i \subseteq U_i$ such that $\sum_i f_i \equiv 1$ (recall that the support $\text{supp } f$ of a function $f : X \rightarrow \mathbb{R}_{\geq 0}$ is, by definition, the complement of the largest open set over which f vanishes identically).

LEMMA 4.4. *If Y is paracompact Hausdorff, then a map $X \rightarrow Y$ is strongly étale if and only if there exists an open cover $\{U_i \subseteq Y\}_i$ such that each map $X \times_Y U_i \rightarrow U_i$ is strongly étale.*

Proof. Fix an open cover $\{U_i \subseteq Y\}_i$ and connection sieves S_i on $X \times_Y U_i \rightarrow U_i$. As Y is paracompact, we may assume that our open cover $\{U_i \subseteq Y\}_i$ is locally finite. Using a partition of unity subordinate to this open cover, we may find another open cover $\{V_I \subseteq Y\}_I$ indexed by nonempty finite subsets I of the original index set, such that $V_I \subseteq \bigcap_{i \in I} U_i$ and $V_I \cap V_J = \emptyset$ unless $I \subseteq J$ or $J \subseteq I$ (explicitly, we may take V_I to be the locus where $\min_{i \in I} f_i > \max_{i \notin I} f_i$). We may now define a connection sieve on $X \rightarrow Y$ as the union over I of $2^{X \times_Y V_I} \cap \bigcap_{i \in I} S_i$. □

An orbispace will be called paracompact (respectively, coarsely finite-dimensional, d -dimensional) if and only if its coarse space is.

PROPOSITION 4.5. *Every paracompact orbispace X has an étale atlas $U \rightarrow X$ for which the projections $U \times_X U \rightrightarrows U$ are strongly étale. In fact, there exists such U of the form $U = \bigsqcup_i V_i$ for an open cover $\{V_i/G_i \subseteq X\}_i$ as in Proposition 3.3.*

Proof. Fix an open cover $\{V_i/G_i \hookrightarrow X\}_i$ as in Proposition 3.3. As $|X|$ is paracompact, we may shrink the spaces V_i (G_i -equivariantly) so as to ensure that the associated open cover $\{|V_i/G_i| \subseteq |X|\}_i$ of coarse spaces is locally finite. Choose a partition of unity $\{f_i : |X| \rightarrow \mathbb{R}_{\geq 0}\}_i$ subordinate to the open cover $\{|V_i/G_i| \subseteq |X|\}_i$. Let $V_i^0 \subseteq V_i$ denote the open subset where $f_i > 0$. We show that the étale atlas $U := \bigsqcup_i V_i^0$ has the desired property.

It suffices to show that for any pair of open embeddings $V/G \hookrightarrow X \hookleftarrow W/H$ and pair of functions $f_V, f_W : |X| \rightarrow \mathbb{R}_{\geq 0}$ supported inside $|V/G|$ and $|W/H|$, respectively, the projection $W^0 \times_X V^0 \rightarrow V^0$ is strongly étale. We begin by considering the map $W \times_X V \rightarrow V$, which is a finite covering space over its open image $W/H \times_X V \subseteq V$ (by pullback from $W \rightarrow W/H$, which is a finite covering space by descent from its pullback $H \times W \rightarrow W$). Thus, every point of $W/H \times_X V$ has a neighborhood over which $W \times_X V \rightarrow V$ is strongly étale. Even better, because V is Hausdorff and G is finite, each G -orbit inside $W/H \times_X V$ has a neighborhood over which $W \times_X V \rightarrow V$ is strongly étale. In other words, each point of $|W/H| \cap |V/G| \subseteq |X|$ has a neighborhood over which $W \times_X V \rightarrow V$ is strongly étale.

Now because $|X|$ is paracompact, there exists a locally finite open cover of $|X|$ by $|X| \setminus (\text{supp } f_V \cup \text{supp } f_W)$ together with open subsets $\{A_i \subseteq |W/H| \cap |V/G|\}_i$ over which $W \times_X V \rightarrow V$ is strongly étale. Fix a partition of unity $g : |X| \rightarrow \mathbb{R}_{\geq 0}$ supported inside $|X| \setminus (\text{supp } f_V \cup \text{supp } f_W)$ and $\{g_i : |X| \rightarrow \mathbb{R}_{\geq 0}\}$ supported inside A_i , that is $g + \sum_i g_i \equiv 1$. Now the patching procedure for connection sieves from the proof of Lemma 4.4 shows that $W \times_X V \rightarrow V$ is strongly étale over the complement of $\text{supp } g$. In particular, it follows by restriction that $W^0 \times_X V^0 \rightarrow V^0$ is strongly étale. \square

A *simplicial complex* is a pair $X = (V, S)$ consisting of a set V (‘vertices’) and a set $S \subseteq 2^V \setminus \{\emptyset\}$ (‘simplices’) of finite subsets of V such that S contains all singletons and $\emptyset \neq A \subseteq B \in S$ implies $A \in S$. The *star* $\text{st}(X, \sigma) \subseteq X$ of a simplex σ in a simplicial complex X is the subcomplex consisting of all simplices $\tau \subseteq X$ with $\sigma \cup \tau \in S(X)$.

A map of simplicial complexes $X \rightarrow Y$ is a map of vertex sets $V(X) \rightarrow V(Y)$ which maps simplices to simplices (the image of an element of $S(X)$ is an element of $S(Y)$). A map of simplicial complexes is called *injective* if and only if the map on vertex sets (hence, also the map on simplices) is injective. A map of simplicial complexes $f : X \rightarrow Y$ is called *étale* (respectively, *locally injective*) if and only if the induced maps on stars $\text{st}(X, \sigma) \rightarrow \text{st}(Y, f(\sigma))$ are isomorphisms (respectively, injective). We call a map of simplicial complexes $X \rightarrow Y$ *sufficiently étale* if and only if every simplex $\sigma \subseteq Y$ (equivalently, every vertex) is the image of a simplex $\tau \subseteq X$ at which $X \rightarrow Y$ is étale (this is a useful weakening of the condition of being surjective and étale, which in the context of simplicial complexes is too strong).

The *geometric realization* $\|X\|$ of a simplicial complex X is the set of tuples $t \in \mathbb{R}_{\geq 0}^{V(X)}$ with $\sum_v t_v = 1$ such that $\{v : t_v > 0\} \in S(X)$, topologized by declaring that the realization of the complete simplex on $k + 1$ vertices has the usual topology and that a realization $\|X\|$ is given the strongest topology for which (the realization of) every map from a complete simplex to X is continuous. The geometric realization of an étale map of simplicial complexes is an étale map of spaces.

A *locally injective simplicial complex groupoid* $M \rightrightarrows O$ consists of simplicial complexes O and M together with structure maps satisfying the axioms of a groupoid, where both maps $M \rightrightarrows O$ are locally injective. Local injectivity of the two maps $M \rightrightarrows O$ implies that the natural map $\|M \times_O M\| \xrightarrow{\sim} \|M\| \times_{\|O\|} \|M\|$ is a homeomorphism, and thus the geometric realization $\|M\| \rightrightarrows \|O\|$ is a topological groupoid. If $\partial O \subseteq O$ denotes the subcomplex consisting of those simplices $\sigma \subseteq O$ for which it is not the case that the first projection $M \rightarrow O$ is étale at every simplex $\tau \subseteq M$ mapped to σ under the second projection, then the natural map $\|O\| \setminus \|\partial O\| \rightarrow \|\|M\| \rightrightarrows \|O\|\|$ is étale. A locally injective simplicial complex groupoid $M \rightrightarrows O$ is called *sufficiently étale* if and only if this map is surjective (equivalently, every vertex of O is M -isomorphic to one not in ∂O).

The *abstract simplex category* \mathbf{Simp} has objects finite totally ordered sets and has morphisms weakly order-preserving maps; every object of \mathbf{Simp} is isomorphic to $[n] := \{0 < \dots < n\}$ for

a unique integer $n \geq 0$. A simplicial object in a category \mathbf{C} is a functor $\mathbf{Simp}^{\text{op}} \rightarrow \mathbf{C}$, and the category of simplicial objects in \mathbf{C} is denoted \mathbf{sC} . If \mathbf{C} is complete (respectively, cocomplete), then so is \mathbf{sC} , and limits (respectively, colimits) are calculated pointwise.

We consider only simplicial sets (objects of the category \mathbf{sSet}) and simplicial groupoids (objects of the category \mathbf{sGrpd}). A simplicial set (or groupoid) will be denoted X_\bullet , where X_n is its set (or groupoid) of n -simplices. We denote by $\Delta_\bullet^n \in \mathbf{sSet}$ the standard n -simplex, given by $[m] \mapsto \text{Hom}([m], [n])$. The Yoneda lemma implies that $X_n = \text{Hom}(\Delta_\bullet^n, X_\bullet)$.

A map of simplicial sets $X_\bullet \rightarrow Y_\bullet$ is called injective if and only if it is so levelwise (i.e. every $X_n \rightarrow Y_n$ is injective). A map $X_\bullet \rightarrow Y_\bullet$ is called étale (respectively, locally injective) if and only if for every map $[n] \rightarrow [m]$ in \mathbf{Simp} , the induced map $X_m \xrightarrow{\sim} Y_m \times_{Y_n} X_n$ is a bijection (respectively, injective) (it is equivalent to impose this condition only for $n = 0$). A map $X_\bullet \rightarrow Y_\bullet$ is called étale (respectively, locally injective) at a given n -simplex σ of X_\bullet if and only if $X_m \xrightarrow{\sim} Y_m \times_{Y_n} \{\sigma\}$ is a bijection (respectively, injective) (this condition at a given σ implies the same at any preimage of σ under any structure map of X_\bullet). A map $X_\bullet \rightarrow Y_\bullet$ is called sufficiently étale if and only if the n -simplices of X_\bullet at which the map is étale surject onto the n -simplices of Y_\bullet (it is equivalent to impose this condition only for $n = 0$). These notions generalize to maps of simplicial groupoids by replacing ‘injectivity’ and ‘surjectivity’ for maps of sets with ‘full faithfulness’ and ‘essential surjectivity’ for functors of groupoids. These properties are all preserved under pullback and closed under composition.

The *geometric realization* $\|X_\bullet\|$ of a simplicial set X_\bullet is the colimit of Δ^n over all maps $\Delta_\bullet^n \rightarrow X_\bullet$. Geometric realization is cocontinuous, and the natural map $\|\lim_\alpha (X_\bullet)_\alpha\| \rightarrow \lim_\alpha \| (X_\bullet)_\alpha \|$ is bijective, however it need not be a homeomorphism even for finite limits (for example, the case of binary products is discussed in [May67, Theorem 14.3, Remark 14.4] and [Whi49, Mil56, BT17]). The map $\|X_\bullet \times_{Y_\bullet} Z_\bullet\| \rightarrow \|X_\bullet\| \times_{\|Y_\bullet\|} \|Z_\bullet\|$ is a homeomorphism if at least one of the maps $X_\bullet \rightarrow Y_\bullet$ and $Z_\bullet \rightarrow Y_\bullet$ is locally injective. Thus, a locally injective simplicial set groupoid $M_\bullet \rightrightarrows O_\bullet$ determines a topological groupoid $\|M_\bullet\| \rightrightarrows \|O_\bullet\|$.

Let us now introduce the geometric realization $\|X_\bullet\|$ of any simplicial groupoid X_\bullet which is étale, meaning that it admits a locally injective sufficiently étale map $U_\bullet \rightarrow X_\bullet$ from a simplicial set U_\bullet . For any simplicial groupoid X_\bullet and any locally injective map $U_\bullet \rightarrow X_\bullet$, the pair of simplicial sets $U_\bullet \times_{X_\bullet} U_\bullet \rightrightarrows U_\bullet$ forms a locally injective simplicial set groupoid, whose geometric realization $\|U_\bullet \times_{X_\bullet} U_\bullet\| \rightrightarrows \|U_\bullet\|$ thus defines a topological stack. The geometric realization of X_\bullet is defined as this topological stack $\| \|U_\bullet \times_{X_\bullet} U_\bullet\| \rightrightarrows \|U_\bullet\| \|$ associated to any locally injective sufficiently étale map $U_\bullet \rightarrow X_\bullet$.

LEMMA 4.6. *The geometric realization of an étale simplicial groupoid X_\bullet is well defined.*

Proof. Let $U_\bullet, U'_\bullet \rightarrow X_\bullet$ be locally injective and sufficiently étale. Let $U''_\bullet := U_\bullet \times_{X_\bullet} U'_\bullet \rightarrow X_\bullet$, and consider the map of simplicial set groupoids $(U''_\bullet \times_{X_\bullet} U''_\bullet \rightrightarrows U''_\bullet) \rightarrow (U_\bullet \times_{X_\bullet} U_\bullet \rightrightarrows U_\bullet)$. As $U''_\bullet \rightarrow U_\bullet$ is locally injective and sufficiently étale, it follows that this map induces an isomorphism of topological stacks, and the same applies to U'_\bullet in place of U_\bullet . □

LEMMA 4.7. *The geometric realization $\|X_\bullet\|$ of an étale simplicial groupoid X_\bullet with finite isotropy is an orbispace.*

Proof. All geometric realizations are Hausdorff, so $\|U_\bullet \times_{X_\bullet} U_\bullet\| \rightarrow \|U_\bullet\| \times \|U_\bullet\|$ is separated, hence $\|X_\bullet\|$ has separated diagonal. As X_\bullet has finite isotropy, the map $U_\bullet \times_{X_\bullet} U_\bullet \rightarrow U_\bullet \times U_\bullet$ has finite fibers, which combined with local injectivity of $U_\bullet \times_{X_\bullet} U_\bullet \rightrightarrows U_\bullet$ implies that $\|U_\bullet \times_{X_\bullet} U_\bullet\| \rightarrow \|U_\bullet\| \times \|U_\bullet\|$ is universally closed, hence $\|X_\bullet\|$ has universally closed diagonal. We have thus shown that $\|X_\bullet\|$ has proper diagonal.

To construct an étale atlas for $\|X_\bullet\|$, let $\partial U_\bullet \subseteq U_\bullet$ denote the simplicial subset consisting of those simplices of U_\bullet at which $U_\bullet \rightarrow X_\bullet$ is not étale. Then $\|U_\bullet\| \setminus \|\partial U_\bullet\| \rightarrow \|X_\bullet\|$ is étale. To see that it is surjective, it suffices to show that $\|U_\bullet \times_{X_\bullet} U_\bullet\| \setminus \|U_\bullet \times_{X_\bullet} \partial U_\bullet\| \rightarrow \|U_\bullet\|$ is surjective (note that surjectivity descends under surjective maps such as $\|U_\bullet\| \rightarrow \|X_\bullet\|$), and this follows because $U_\bullet \rightarrow X_\bullet$ is sufficiently étale. \square

A simplicial complex X gives rise to a simplicial set $b_\bullet X$ (its *barycentric subdivision*) whose n -simplices are chains of simplices $\sigma_0 \subseteq \dots \subseteq \sigma_n \subseteq X$ (in other words, $b_\bullet X$ is the nerve of $S(X)$). Barycentric subdivision preserves injectivity, local injectivity, étale, and sufficiently étale. There is a natural identification of geometric realizations $\|X\| = \|b_\bullet X\|$. Moreover, for a locally injective sufficiently étale simplicial complex groupoid $M \rightrightarrows O$, there is a natural identification $\|M\| \rightrightarrows \|O\| = \|[b_\bullet M \rightrightarrows b_\bullet O]\|$ (the simplicial groupoid $[b_\bullet M \rightrightarrows b_\bullet O]$ is étale because $M \rightrightarrows O$ is sufficiently étale).

A simplicial complex of groups (Z, G) also admits a barycentric subdivision $b_\bullet(Z, G)$ which is a simplicial groupoid. In fact, we define $b_\bullet(Z, G)$ for any simplicial complex Z equipped with a functor $G : S(Z)^{\text{op}} \rightarrow \text{Grpd}$ from the face poset to groupoids (a simplicial complex of groups (Z, G) determines such a functor which sends σ to $\mathbb{B}G_\sigma$). The groupoid of n -simplices in the barycentric subdivision $b_\bullet(Z, G)$ is now defined as the groupoid of functors from the category $0 \rightarrow \dots \rightarrow n$ to the category whose objects are pairs $\sigma \in S(Z)$ and $o \in G_\sigma$ and whose morphisms are inclusions $\sigma_1 \subseteq \sigma_2$ covered by maps $o_1 \rightarrow o_2|_{\sigma_1}$. The barycentric subdivision of a simplicial complex of groups is étale, as can be seen as follows. For any $\sigma \subseteq Z$ and $o \in G_\sigma$, we consider the functor $S(\text{st}(Z, \sigma))^{\text{op}} \rightarrow \text{Grpd}$ given by $\tau \mapsto G_{\sigma \cup \tau} \times_{G_\sigma} \{o\}$. Applying the nerve construction from just above to this functor, we obtain a simplicial set mapping to $b_\bullet(Z, G)$, which is the required locally injective map which is étale over $(\sigma, o) \in b_0(Z, G)$. An essentially equivalent discussion (albeit without barycentrically subdividing) appears in [BH99, 12.24–12.25]. As $b_\bullet(Z, G)$ is étale, it has a geometric realization $\|b_\bullet(Z, G)\|$ which we also write as $\|(Z, G)\|$. When G are finite groups (or, more generally, groupoids with finite isotropy), then the geometric realization $\|(Z, G)\|$ is an orbispace by Lemma 4.7, and this is what we have been calling the orbispace presented by the simplicial complex of finite groups (Z, G) .

LEMMA 4.8. *Let $M \rightrightarrows O$ be a locally injective simplicial complex groupoid with the following properties.*

- *The vertices of every simplex of O are pairwise non-isomorphic via M .*
- *If simplices σ and σ' of O have vertex sets which are isomorphic via M , then σ and σ' are themselves isomorphic via M .*

Then there is a simplicial complex of groups giving rise to the same simplicial groupoid as $M \rightrightarrows O$.

Proof. The hypotheses imply that there is a simplicial complex Z whose vertices are the isomorphism classes in the vertex groupoid $V(M) \rightrightarrows O(M)$, and whose simplices are the M -isomorphism classes of simplices of O . Now $M \rightrightarrows O$ defines a functor $G : S(Z)^{\text{op}} \rightarrow \text{Grpd}$, and there is a natural isomorphism between $b_\bullet(Z, G)$ and the simplicial groupoid $[b_\bullet M \rightrightarrows b_\bullet O]$. By definition, all the groupoids G_σ have a single isomorphism class, and all the functors $G_\tau \rightarrow G_\sigma$ are faithful (this follows from local injectivity of $M \rightrightarrows O$). Choosing (independently) a base object of each G_σ shows (Z, G) comes from a simplicial complex of groups. \square

The *nerve* $N(X, \{U_i\}_i)$ of a collection of open sets $\{U_i \subseteq X\}_i$ is the simplicial complex whose vertices V are the indices i with $U_i \neq \emptyset$, and in which a collection I of indices spans a

simplex (i.e. $I \in S$) if and only if $\bigcap_{i \in I} U_i \neq \emptyset$. A partition of unity $\{f_i : X \rightarrow \mathbb{R}_{\geq 0}\}_i$ subordinate to a locally finite open cover $\{U_i \subseteq X\}_i$ defines a map from X to the geometric realization $\|N(X, \{U_i\}_i)\|$ of the nerve of the open cover.

PROPOSITION 4.9. *For any paracompact orbispace X , there exists a simplicial complex of groups (Z, G) and a map $X \rightarrow \|(Z, G)\|$ which is injective on isotropy groups. Moreover, we may take Z to be locally finite, and we may take the groups G_z for vertices $z \in Z$ to be isotropy groups of points of X . If X is coarsely finite-dimensional (respectively, d -dimensional), then Z may be taken to be finite-dimensional (respectively, d -dimensional).*

Proof. We begin with an open cover $\{V_i/G_i \subseteq X\}_i$ with the properties guaranteed by Proposition 4.5, and we set $U := \bigsqcup_i V_i$. As $|X|$ is paracompact, by G_i -equivariantly shrinking the spaces V_i , we may assume that the associated open cover $\{|V_i/G_i| \subseteq |X|\}_i$ of coarse spaces is locally finite. We fix a covering sieve S on $U \times_X U$ which is invariant under the ‘exchange’ (i.e. ‘inverse’) involution of $U \times_X U$ and is a connection sieve for both projections $U \times_X U \rightrightarrows U$. We also fix a partition of unity $\{f_i : |X| \rightarrow \mathbb{R}_{\geq 0}\}_i$ subordinate to the open cover $\{|V_i/G_i| \subseteq |X|\}_i$.

Denote by $o(x) \subseteq U$ and $o_i(x) \subseteq V_i$ the fibers over $x \in |X|$, so $o(x) = \bigsqcup_i o_i(x)$; similarly define $m(x) \subseteq U \times_X U$ and $m_{ij}(x) \subseteq V_i \times_X V_j$ with $m(x) = \bigsqcup_{i,j} m_{ij}(x)$. These sets are finite because the G_i are finite and $\{|V_i/G_i| \subseteq |X|\}_i$ is locally finite. Furthermore, they have the discrete topology, because U is Hausdorff and $U \times_X U$ is Hausdorff (because U is Hausdorff and $X \rightarrow X \times X$ is separated).

The Hausdorff property implies that the inclusions $o_i(x) \subseteq V_i$ and $m_{ij}(x) \subseteq V_i \times_X V_j$ admit retractions defined in some open neighborhood. Now the inverse images of small open neighborhoods $x \in |Z_x| \subseteq |X|$ (i.e. open substacks $Z_x \subseteq X$) form a basis of neighborhoods of $o_i(x)$ and $m_{ij}(x)$, so for sufficiently small Z_x , these inverse images are naturally disjoint unions $U_x = \bigsqcup_{o \in o(x)} U_o$ and $U_x \times_X U_x = \bigsqcup_{m \in m(x)} U_m$. Note that this applies only inside V_i and $V_i \times_X V_j$ for which $o_i(x)$ and $m_{ij}(x)$ are nonempty: the full inverse image of Z_x inside U may intersect other V_i nontrivially. By shrinking Z_x further, we may ensure that the retraction $(U_x \times_X U_x \rightrightarrows U_x) \rightarrow (m(x) \rightrightarrows o(x))$ is a map of groupoids. For later purposes, let us also take Z_x small enough so that:

- if $x \in |V_i/G_i|$, then $|Z_x| \subseteq |V_i/G_i|$;
- if $x \notin \text{supp } f_i$, then $|Z_x| \cap \text{supp } f_i = \emptyset$;
- if $f_i(x) > 0$, then $f_i > 0$ over all of $|Z_x|$.

Each of these conditions can be ensured on its own, and because the open cover $\{|V_i/G_i| \subseteq |X|\}_i$ is locally finite, we can ensure all at once.

We also shrink Z_x so as to ensure that each $U_m \in S$ (our chosen connection sieve), which has the following implication: given $o, o' \in U$ with $U_o \cap U_{o'} \neq \emptyset$, the relation $U_m \cap U_{m'} \neq \emptyset$ is a partial bijection between lifts $m, m' \in U \times_X U$ of o and o' . Furthermore, the domain of this bijection is as large as possible: for $U_o \cap U_{o'} \neq \emptyset$ (so $o \in o_i(x)$ and $o' \in o_i(x')$ for some i) with $x, x' \in |V_j/G_j|$, we get a full bijection between the inverse images of o and o' inside $m_{ij}(x)$ and $m_{ij}(x')$ (this follows because the projection $V_i \times_X V_j \rightarrow V_i$ is a finite covering space of degree $|G_j|$ over $U_o \cup U_{o'}$).

We now consider the nerves $N(U, \{U_o\}_{o \in U})$ and $N(U \times_X U, \{U_m\}_{m \in U \times_X U})$. Note that for simplices in these nerves, namely subsets $O \subseteq U$ or $M \subseteq U \times_X U$ with $\bigcap_{o \in O} U_o \neq \emptyset$ or $\bigcap_{m \in M} U_m \neq \emptyset$, the maps $O \rightarrow |X|$ or $M \rightarrow |X|$ are injective. The natural maps on index sets $U \times_X U \rightrightarrows U$ determine maps of nerves

$$N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \rightrightarrows N(U, \{U_o\}_{o \in U}).$$

These maps are locally injective; indeed, local injectivity means that for every $o, o' \in U$ with $U_o \cap U_{o'} \neq \emptyset$ and every $m \in U \times_X U$ projecting to o , there is at most one lift $m' \in U \times_X U$ of o' with $U_m \cap U_{m'} \neq \emptyset$, and this is a direct consequence of our assumption that every $U_m \in S$. Let us now argue that there is a natural composition map

$$N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \times_{N(U, \{U_o\}_{o \in U})} N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \longrightarrow N(U \times_X U, \{U_m\}_{m \in U \times_X U}).$$

More precisely, we claim that for nonempty finite subsets $M \subseteq U \times_X U$ with $\bigcap_{m \in M} U_m \neq \emptyset$ and $M' \subseteq U \times_X U$ with $\bigcap_{m \in M'} U_m \neq \emptyset$ projecting to $O \subseteq U$ (under the first and second projections, respectively), the subset $M'' \subseteq U \times_X U$ defined by applying the composition map $(U \times_X U) \times_U (U \times_X U) \rightarrow U \times_X U$ to M and M' also satisfies $\bigcap_{m \in M''} U_m \neq \emptyset$. This claim follows from the property that every $U_m \in S$ (indeed, this property implies that $\bigcap_{m \in M} U_m \xrightarrow{\sim} \bigcap_{o \in O} U_o \xleftarrow{\sim} \bigcap_{m \in M'} U_m$). We have thus defined a locally injective simplicial complex groupoid

$$M := N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \rightrightarrows N(U, \{U_o\}_{o \in U}) =: O.$$

We now show that this locally injective simplicial complex groupoid is sufficiently étale. Every isomorphism class of vertex (equivalently, every $x \in |X|$ with $|Z_x| \neq \emptyset$) has a representative $o \in V_i \subseteq U$ with $f_i(o) = f_i(x) > 0$. To show that these representatives are étale, let $m \in U \times_X U$ be a morphism with source o and target o' , and let $U_{o'} \cap U_{o''} \neq \emptyset$. We must show that there is a bijection between morphisms $o \rightarrow o'$ and $o \rightarrow o''$. The morphisms in question all have source inside V_i , so we really can consider just $V_i \times U$ for the present purpose. Now the bulleted conditions on $|Z_x|$ from above imply that because $U_{o'} \cap U_{o''} \neq \emptyset$ and $f_i(o') > 0$, we have $U_{o'} \cup U_{o''} \subseteq |V_i/G_i| \subseteq |X|$. Thus, over $U_{o'} \cap U_{o''}$ the map $V_i \times_X U \rightarrow U$ is a finite covering space of degree $|G_i|$. Thus, all points have the same number of lifts, so it follows that the connection sieve property gives us a bijection between lifts.

We have already seen above that our sufficiently étale locally injective simplicial complex groupoid satisfies the first hypothesis of Lemma 4.8, and the second hypothesis follows from the partial bijection property derived above from the connection sieve. Thus, by Lemma 4.8, there is a simplicial complex of groups (Z, G) giving rise to the same simplicial groupoid $b_\bullet(Z, G) = [b_\bullet M \rightrightarrows b_\bullet O]$ and thus (because $M \rightrightarrows O$ is sufficiently étale) to the same geometric realization $\| (Z, G) \| = [\| M \| \rightrightarrows \| O \|]$. The groups G_z associated to vertices $z \in Z$ are, by definition, isotropy groups of the vertex groupoid $V(M) \rightrightarrows V(O)$, which are, by definition, isotropy groups of points of X .

To conclude, it remains to define a map $X \rightarrow \| (Z, G) \|$ which is injective on isotropy groups. To define this map, we shrink the Z_x so that the open cover $\{|Z_x| \subseteq |X|\}_{x \in |X|}$ is locally finite, and choose a partition of unity $\{g_x : |X| \rightarrow \mathbb{R}_{\geq 0}\}_{x \in |X|}$ subordinate to the open cover $\{|Z_x| \subseteq |X|\}_{x \in |X|}$. These maps g_x lift to maps $g_o : U \rightarrow \mathbb{R}_{\geq 0}$ supported inside U_o and $g_m : U \times_X U \rightarrow \mathbb{R}_{\geq 0}$ supported inside U_m . The collection of these lifts defines a map of topological groupoids

$$(U^0 \times_X U^0 \rightrightarrows U^0) \rightarrow (\| N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \| \rightrightarrows \| N(U, \{U_o\}_{o \in U}) \|),$$

where U^0 is the disjoint union of the open loci $V_i^0 \subseteq V_i$ where $f_i > 0$ (the maps g_o and g_m do not define a map over all of $U \times_X U \rightrightarrows U$ due to the fact that $U_x = \bigsqcup_{o \in o(x)} U_o$ and $U_x \times_X U_x = \bigsqcup_{m \in m(x)} U_m$ may not be the full inverse images of Z_x inside U and $U \times_X U$). It remains to check that this map is injective on isotropy groups; in other words, for $o, o' \in U^0$ we must show that the map $\{o\} \times_X \{o'\} \rightarrow \| N(U \times_X U, \{U_m\}_{m \in U \times_X U}) \|$ is injective. Distinct elements of $\{o\} \times_X \{o'\}$ are, in particular, distinct lifts of o , which therefore cannot lie in any common U_m because $U_m \rightarrow U$ is injective, so we see that the map is indeed injective on isotropy groups. \square

Proof of Theorem 1.1. Apply Proposition 4.9 to find a simplicial complex of finite groups (Z, G) and a map $X \rightarrow \|(Z, G)\|$ which is injective on isotropy groups. Now Theorem 2.13 applies to $\|(Z, G)\|$ to give a vector bundle $V \rightarrow \|(Z, G)\|$, and the pullback of this bundle to X satisfies the desired property because $X \rightarrow \|(Z, G)\|$ is injective on isotropy groups. \square

5. From vector bundles to principal bundles

We derive Corollaries 1.2 and 1.3 from Theorem 1.1.

A hermitian inner product on a complex vector bundle $V \rightarrow X$ is a map $h : V \times_X V \rightarrow \mathbb{C}$ satisfying $h(v, w) = \overline{h(w, v)}$, $h(v, \alpha w) = \alpha h(v, w)$ for $\alpha \in \mathbb{C}$, and $h(v, v) > 0$ for $v \neq 0$ (meaning, these conditions are imposed on the fiber $h_x : V_x \times V_x \rightarrow \mathbb{C}$ over each map $* \xrightarrow{x} X$).

LEMMA 5.1. *A complex vector bundle over a paracompact orbispace X admits a hermitian inner product.*

Proof. Begin with an étale atlas $\bigsqcup_i U_i \rightarrow X$ such that the pullback of V to every U_i is trivial. By Corollary 3.5, we may refine this cover further so that each map $U_i \rightarrow X$ is the composition of $U_i \rightarrow U_i/G_i$ with an open embedding $U_i/G_i \hookrightarrow X$. The pullback of V to each U_i is trivial, hence admits a hermitian inner product; by averaging, we may make it G_i -invariant thus giving a hermitian inner product h_i on the restriction of V to each $U_i/G_i \subseteq X$. As $|X|$ is paracompact, there is a partition of unity φ_i subordinate to this open cover of $|X|$, and hence $\sum_i \varphi_i h_i$ is the desired hermitian inner product on V . \square

COROLLARY 5.2. *Every short exact sequence of vector bundles over a paracompact orbispace splits.*

Given a rank n complex vector bundle $V \rightarrow X$ with hermitian inner product, the associated frame bundle $P \rightarrow X$ is defined by declaring that a map $U \rightarrow P$ (U a topological space) is a map $U \rightarrow X$ together with an isomorphism $\mathbb{C}^n \times U \rightarrow V \times_X U$ under which the pullback of h is the standard hermitian inner product on \mathbb{C}^n . There is an action of $U(n)$ on P (by precomposition with automorphisms of \mathbb{C}^n respecting its hermitian inner product), giving P the structure of a principal bundle over X , meaning that for every map $U \rightarrow X$ from a topological space, there is an open cover $\{U_i \subseteq U\}_i$ such that $P \times_X U_i \rightarrow U_i$ is isomorphic to $U(n) \times U_i \rightarrow U_i$ with $U(n)$ acting by left multiplication on the first factor.

Proof of Corollary 1.2. Let $V \rightarrow X$ be the rank n complex vector bundle produced by Theorem 1.1. By Lemma 5.1, there exists a hermitian inner product on V . The total space P of the associated principal bundle $P \rightarrow X$ has trivial isotropy because the isotropy groups of X act faithfully on the fibers of V .

We claim that P is a Hausdorff topological space. By Corollary 3.6, it is enough to show that P is an orbispace. As $P \rightarrow X$ is representable, the pullback of an étale atlas for X is an étale atlas for P . The diagonal of P may be expressed as the composition $P \rightarrow P \times_X P \rightarrow P \times P$. The second map $P \times_X P \rightarrow P \times P$ is proper, being a pullback of the diagonal of X . To check that the first map $P \rightarrow P \times_X P$ is proper, it suffices by descent to show that its pullback under an étale atlas $U \rightarrow X$ is proper. This pullback is proper because $P \times_X U \rightarrow U$ is a principal $U(n)$ bundle (of topological spaces). We have thus shown that P is a topological space.

The map $P \rightarrow X$ is representable and admits local sections (by definition), so the topological groupoid $P \times_X P \rightrightarrows P$ presents the stack X . It can be seen by inspection that $P \times_X P \rightrightarrows P$ is the action groupoid of the $U(n)$ action on P . \square

Let \mathbf{Sm} denote the category of topological spaces equipped with a maximal atlas of charts from open sets of $\bigsqcup_{n \geq 0} \mathbb{R}^n$ with smooth transition maps (a smooth manifold an object of \mathbf{Sm} whose underlying topological space is Hausdorff). A smooth structure on a stack $F : \mathbf{Top}^{\text{op}} \rightarrow \mathbf{Grpd}$ is a substack F^{sm} of the pullback of F under the forgetful functor $\mathbf{Sm} \rightarrow \mathbf{Top}$ (meaning that $F^{\text{sm}}(U)$ is a full subcategory of $F(U)$ for $U \in \mathbf{Sm}$); maps $U \rightarrow F$ lying in (the essential image of) F^{sm} are then called smooth. Given a map of stacks $F \rightarrow G$ and a smooth structure on G , we may consider the pullback smooth structure on F , defined as those maps $U \rightarrow F$ whose composition with $F \rightarrow G$ is smooth. A map of stacks equipped with smooth structures $(F, F^{\text{sm}}) \rightarrow (G, G^{\text{sm}})$ is called smooth if and only if the composition of any smooth map $U \rightarrow F$ with $F \rightarrow G$ is a smooth map $U \rightarrow G$ (equivalently, F^{sm} is contained inside the pullback of G^{sm}). Stacks with smooth structures form a 2-category just like stacks. This category is complete, and limits $\lim_{\alpha} (F_{\alpha}, F_{\alpha}^{\text{sm}})$ are calculated by taking the limit of the underlying stacks $\lim_{\alpha} F_{\alpha}$ and declaring a map $U \rightarrow \lim_{\alpha} F_{\alpha}$ to be smooth if and only if every induced map $U \rightarrow F_{\alpha}$ is smooth. The category \mathbf{Sm} embeds fully faithfully into the category of stacks with smooth structures, by sending $X \in \mathbf{Sm}$ to the Yoneda functor of its underlying topological space, equipped with the smooth structure consisting of those maps $U \rightarrow X$ which are morphisms in \mathbf{Sm} .

DEFINITION 5.3. A *smooth orbifold* X is an orbispace equipped with a smooth structure such that for every (equivalently, some) étale atlas $U \rightarrow X$, we have $U \in \mathbf{Sm}$ when U is equipped with the pullback of the smooth structure on X .

By Proposition 3.3, a stack X with a smooth structure is a smooth orbifold if and only if $|X|$ is Hausdorff and there is an open cover of X by V_i/G_i for smooth manifolds V_i equipped with smooth actions of finite groups G_i .

Proof of Corollary 1.3. Let $V \rightarrow X$ be the complex vector bundle produced by Theorem 1.1. It suffices to define a smooth structure on V , because then we may follow the arguments of Lemma 5.1 and Corollary 1.2 in the smooth category.

We begin by arguing that a complex vector bundle V over a smooth orbifold X has a smooth structure in a neighborhood of any given point of $|X|$. We may thus assume that $X = U/G$ for a smooth manifold U acted on smoothly by a finite group G , and that the pullback of V to U is trivial. By shrinking this chart further (and adjusting G), we may assume that our given point of $|X|$ corresponds to a point $u \in U$ fixed by G . Choose any trivialization $V_u \times U \rightarrow V \times_X U$ where V_u denotes the fiber over u . By averaging, we may make this map G equivariant, and it remains an isomorphism in a neighborhood of u . We thus obtain a trivialization of $V \times_X U$ near u in which the action of G is constant (independent of the U coordinate). The standard smooth structure in this trivialization is thus, in particular, invariant under the action of G , giving the desired smooth structure on V near our given point of $|X|$.

Having shown the existence of smooth structures locally, we now show how to patch them together. To show that there exists a smooth structure on V over all of X , it suffices to show that for open subsets $A, B \subseteq |X|$ and smooth structures on $V|_A$ and $V|_B$, there exists a smooth structure on $V|_{A \cup B}$ restricting to the given smooth structure on $V|_A$ (indeed, this allows us to patch together smooth structures over arbitrary unions of open sets, by choosing a well ordering and adding one open set at a time). To prove this pairwise patching statement, it suffices to show that an isomorphism of smooth vector bundles may be approximated by a smooth isomorphism (then apply this to the identity map on $V|_{A \cap B}$ equipped with the restrictions of the two given smooth structures). As $|X|$ is locally the quotient of Euclidean space by a finite group action, it is locally metrizable; because it is paracompact and Hausdorff, it is thus metrizable.

Thus, every open subset of $|X|$ is metrizable, hence paracompact. We may now conclude with a smooth partition of unity argument. \square

6. K -theory of orbispaces

We derive Corollary 1.4 from Theorem 1.1. We then derive from Corollary 1.4 some basic properties of the K -theory of finite-rank vector bundles on orbispaces (some of this derivation is borrowed, with various technical differences, from [LO01, § 3] and [Seg68]). In what follows, ‘vector bundle’ can be taken to mean either real vector bundle or complex vector bundle (always of finite rank).

Proof of Corollary 1.4. Denote our given map by $f : X \rightarrow Y$, and let E be the given vector bundle on X . Let F be any vector bundle on Y satisfying the conclusion of Theorem 1.1.

We first consider the local situation on X . Fix $x \in X$, and let $U/G \hookrightarrow X$ be an open embedding sending $u \in U$ fixed by G to $x \in X$ and inducing an isomorphism $G \xrightarrow{\sim} G_x$ (such an open embedding exists by the proof of Proposition 3.3). By shrinking U , we may ensure that the pullbacks of E and F to U are trivial. Choose any map on U from the pullback of E to the pullback of $F^{\oplus N}$ (some integer $N < \infty$) which is injective and G -equivariant at u , and average it to make it equivariant everywhere. This produces over U/G a map from E to $f^*F^{\oplus N}$ which is injective at x . Now because $|X|$ is paracompact Hausdorff, hence normal, there exists a continuous function $\varphi : |X| \rightarrow [0, 1]$ supported inside $|U/G|$ with $\varphi(x) = 1$. Multiplying by this function yields a map $E \rightarrow f^*F^{\oplus N}$ defined on all of X which is injective at x . Note that the integer N may be taken independent of x because the rank of E is bounded.

We now combine the above maps to give an everywhere injective map $E \rightarrow f^*F^{\oplus M}$ as follows. Fix an open cover $|X| = \bigcup_i U_i$ and maps $f_i : E \rightarrow f^*F^{\oplus N}$ defined over all of X which are injective over U_i . As $|X|$ is coarsely finite-dimensional, we may refine this open cover so that it is locally finite and has nerve of dimension $\leq d$ for some integer $d < \infty$. As in the proof of Lemma 4.4, there is yet another open cover of $|X|$ by open sets V_I indexed by the nonempty subsets I of the index set of the U_i , such that $V_I \subseteq \bigcap_{i \in I} U_i$ and $V_I \cap V_J = \emptyset$ unless $I \subseteq J$ or $J \subseteq I$. Note that these conditions imply that $V_I = \emptyset$ unless $|I| \leq d + 1$ and that $V_I \cap V_J = \emptyset$ if $|I| = |J|$ and $I \neq J$. Now let $\sum_I \varphi_I \equiv 1$ be a partition of unity subordinate to the open cover $|X| = \bigcup_I V_I$. We may now define our desired everywhere injective map $E \rightarrow f^*F^{\oplus (d+1)N}$ (so $M = (d + 1)N$) by the formula $\sum_I \alpha_{|I|} \varphi_I f_{i(I)}$, where $i(I)$ is any choice of index in the set I , and $\alpha_r : F^{\oplus N} \hookrightarrow (F^{\oplus N})^{\oplus (d+1)} = F^{\oplus (d+1)N}$ is the inclusion of the r th direct summand.

We have thus constructed an injection $E \rightarrow f^*F^{\oplus M}$ over X . \square

DEFINITION 6.1. For any stack X , denote by $\text{Vect}(X)$ the set of isomorphism classes of vector bundles on X of bounded rank.

Direct sum of vector bundles equips $\text{Vect}(X)$ with the structure of an abelian monoid. A map of stacks $X \rightarrow Y$ induces a pullback map $\text{Vect}(Y) \rightarrow \text{Vect}(X)$.

The reason we restrict attention to vector bundles of bounded rank is so that we may appeal to Corollary 1.4. A counterexample of Lück and Oliver [LO01, Example 3.11] shows that the K -theory of vector bundles of unbounded rank fails to be a cohomology theory on (possibly infinite) simplicial complexes of groups, whereas we show that the K -theory of vector bundles of bounded rank is a cohomology theory for such.

For an abelian monoid M , its group completion $M[-M]$ is the quotient of the free abelian group on the underlying set of M by the subgroup generated by $[a] + [b] - [a + b]$ for $a, b \in M$. The map $M \rightarrow M[-M]$ is universal (initial) among maps from M to an abelian group.

DEFINITION 6.2. We define $K^0(X)$ as the group completion of $\text{Vect}(X)$ for any stack X .

A map of stacks $X \rightarrow Y$ induces a pullback map $K^0(Y) \rightarrow K^0(X)$. The functor K^0 is finitely additive, in the sense that $K^0(\emptyset) = 0$ and the natural map $K^0(X \sqcup Y) \rightarrow K^0(X) \oplus K^0(Y)$ is an isomorphism. In the present context of vector bundles of bounded rank, additivity does not hold for general infinite disjoint unions.

Eventually, we will restrict attention to the K -theory of orbispaces satisfying the hypothesis of Theorem 1.1. However, we impose such hypotheses gradually, as they become relevant.

To show that K -theory is homotopy invariant, the following is the key assertion.

LEMMA 6.3. *For any paracompact orbispace X , every vector bundle on $X \times [0, 1]$ is pulled back from X .*

Proof. Let a vector bundle V over $X \times [0, 1]$ be given.

We first discuss the local structure around a given point $x \in X$. Fix an open embedding $Y/G \hookrightarrow X$ and a lift $y \in Y$ of x fixed by G . We consider the pullback bundle $V|_{Y \times [0, 1]}$ on $Y \times [0, 1]$, which is a G -equivariant vector bundle. For each $t \in [0, 1]$, there exist open neighborhoods $t \in T \subseteq [0, 1]$ and $y \in U \subseteq Y$ such that $V|_{U \times T}$ is trivial. By the compactness of $[0, 1]$, we may assume U is independent of t . Replacing Y with U , we may assume that for each $t \in [0, 1]$ there exists an open neighborhood $t \in T \subseteq [0, 1]$ such that $V|_{Y \times T}$ is trivial. A trivialization induces a map $V|_{Y \times T} \rightarrow p_T^* V|_{\{y\} \times T}$ which is the identity over $\{y\} \times T$. Averaging makes this map G -equivariant, so it descends to a map $V|_{Y/G \times T} \rightarrow p_T^* V|_{\{y\}/G \times T}$ which is still the identity over $\{y\} \times T$. It is thus an isomorphism over some $U \times T'$; another compactness argument and shrinking of Y ensures that we have a collection of isomorphisms $V|_{Y/G \times T} \rightarrow p_T^* V|_{\{y\}/G \times T}$ which are the identity over $\{y\} \times T$. Patching these together via a partition of unity on $[0, 1]$ and further shrinking Y produces an isomorphism

$$V|_{Y/G \times [0, 1]} \xrightarrow{\sim} p_{[0, 1]}^* V|_{\{y\}/G \times [0, 1]},$$

which is the identity over $\{y\} \times [0, 1]$. Any vector bundle over $\mathbb{B}G \times [0, 1]$ is pulled back from $\mathbb{B}G$, and hence we conclude that $V|_{Y/G \times [0, 1]}$ is pulled back from Y/G , for some neighborhood Y/G of $x \in X$.

We now globalize. Begin with an open cover $|X| = \bigcup_i U_i$ and over each U_i an isomorphism $\xi_i : V|_{U_i \times [0, 1]} \rightarrow (p_X^* V|_{X \times \{0\}})|_{U_i \times [0, 1]}$ which is the identity over $U_i \times \{0\}$. Note that for any $x \in U_i$ and any $t, t' \in [0, 1]$, the map ξ_i determines an isomorphism

$$V_{(x, t)} \xrightarrow{\xi_i^{-1} \circ \xi_i} V_{(x, t')},$$

which is the specialization of an isomorphism between the pullbacks of V under the two projections $X \times [0, 1]^2 \rightarrow X \times [0, 1]$. Using this observation, we may now patch together the ξ_i into an isomorphism $V \rightarrow p_X^* V|_{X \times \{0\}}$ as follows. Fix a partition of unity $\varphi_i : |X| \rightarrow \mathbb{R}_{\geq 0}$ subordinate to the open cover $|X| = \bigcup_i U_i$, and fix an arbitrary total ordering of the index set of the open cover. For $(x, t) \in X \times [0, 1]$, let $i = 1, \dots, k$ denote the indices of the open sets U_i containing x , ordered as in the fixed total order, and consider the composition of isomorphisms

$$V_{(x, t)} \xrightarrow{\xi_1^{-1} \circ \xi_1} V_{(x, (1 - \varphi_1(x)) \cdot t)} \xrightarrow{\xi_2^{-1} \circ \xi_2} V_{(x, (1 - \varphi_1(x) - \varphi_2(x)) \cdot t)} \xrightarrow{\xi_3^{-1} \circ \xi_3} \dots \xrightarrow{\xi_k^{-1} \circ \xi_k} V_{(x, 0)}.$$

This fiberwise description now translates into the desired isomorphism of vector bundles $V \rightarrow p_X^* V|_{X \times \{0\}}$. □

Two maps $X \rightarrow Y$ will be called homotopic if and only if there exists a map $X \times [0, 1] \rightarrow Y$ whose restrictions to $X \times \{0\}$ and $X \times \{1\}$ are the two given maps. A map $X \rightarrow Y$ is called

a homotopy equivalence if and only if there exists a map $Y \rightarrow X$ such that the compositions $X \rightarrow Y \rightarrow X$ and $Y \rightarrow X \rightarrow Y$ are homotopic to the respective identity maps.

COROLLARY 6.4 (Homotopy invariance). *If X is a paracompact orbispace, then homotopic maps $X \rightarrow Y$ induce the same map $K^0(Y) \rightarrow K^0(X)$.*

For any map of abelian monoids $M' \rightarrow M$, the quotient M/M' is, as a set, the quotient of M by the equivalence relation $x \sim_{M'} y$ if and only if there exist $a, b \in M'$ with $x + a = y + b$, equipped with the descent of the monoid operation from M . The map $M \rightarrow M/M'$ is initial among maps from M to an abelian monoid sending M' to zero. If (the image inside M of) M' is an abelian group, then the kernel of the quotient map $M \rightarrow M/M'$ is precisely M' . The group completion $M[-M]$ is the quotient of $M \times M$ by the diagonal submonoid M .

DEFINITION 6.5. For any map of stacks $f : Y \rightarrow X$, the relative K -theory $K^0(X, Y)$ is defined as follows. We consider the set $\text{Tri}(X, Y)$ of isomorphism classes of triples (E_0, E_1, i) where E_0, E_1 are vector bundles on X of bounded rank and $i : f^*E_0 \rightarrow f^*E_1$ is an isomorphism between their pullbacks to Y . Now $\text{Tri}(X, Y)$ is an abelian monoid under direct sum, and it contains $\text{Vect}(X)$ as the submonoid of triples of the form $(E, E, f^* \text{id}_E)$. The relative K -theory is the quotient

$$K^0(X, Y) := \text{Tri}(X, Y) / \text{Vect}(X).$$

Note that *a priori* $K^0(X, Y)$ is merely an abelian monoid, not an abelian group.

A map $(Y \rightarrow X) \rightarrow (Y' \rightarrow X')$ (i.e. a commutative square) induces a map $K^0(X', Y') \rightarrow K^0(X, Y)$. There is a natural map $K^0(X, Y) \rightarrow K^0(X)$ given by sending (E_0, E_1, i) to the formal difference $[E_0] - [E_1]$, and the composition $K^0(X, Y) \rightarrow K^0(X) \rightarrow K^0(Y)$ is zero. The map $K^0(X, \emptyset) \rightarrow K^0(X)$ is an isomorphism (because $\text{Tri}(X, \emptyset)$ is simply $\text{Vect}(X) \times \text{Vect}(X)$).

A *pair* (X, A) shall mean that A is a closed substack of X . A map of pairs $(X, A) \rightarrow (Y, B)$ means a map $X \rightarrow Y$ sending A inside B . Note that paracompactness, coarse finite-dimensionality, being an orbispace, and the hypothesis of Theorem 1.1 all pass to closed substacks.

Remark 6.6. Even for very nice orbispace pairs (X, A) , we *cannot* in general form the quotient X/A in a reasonable way. For example, for an orbispace X , if we could reasonably define $(X \times [0, 1]) / (X \times \{0\})$ as an orbispace, it would follow that X is (globally) the quotient of a topological space by a finite group action (a very special property).

For any pair (X, A) , let $\text{cyl}(X, A)$ denote the pair $((X \times \{0\}) \cup (A \times [0, 1]), A \times \{1\})$. Here $(X \times \{0\}) \cup (A \times [0, 1])$ denotes the union of closed substacks of $X \times [0, 1]$ (recall that closed substacks of a stack Z are in bijective correspondence with closed subsets of $|Z|$, so by union of closed substacks we mean union of subsets of $|Z|$). There is a natural map of pairs $\text{cyl}(X, A) \rightarrow (X, A)$. A map of pairs $f : (X, A) \rightarrow (X', A')$ induces a map $\text{cyl}(f) : \text{cyl}(X, A) \rightarrow \text{cyl}(X', A')$.

LEMMA 6.7. *For any pair (X, A) where A is a paracompact orbispace, $K^0(\text{cyl}(X, A))$ is an abelian group.*

Proof. Let $(E_0, E_1, i : E_0|_{A \times \{1\}} \rightarrow E_1|_{A \times \{1\}})$ be a triple representing an arbitrary element of $K^0(\text{cyl}(X, A))$. We claim that the triple $(E_1, E_0, -i)$ is an inverse to it. It suffices to show that

$$(E_0 \oplus E_1, E_0 \oplus E_1, \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}) \quad \text{and} \quad (E_0 \oplus E_1, E_0 \oplus E_1, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix})$$

are isomorphic. In other words, it suffices to show that $\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ is the restriction to $A \times \{1\}$ of an automorphism of $E_0 \oplus E_1$. By Lemma 6.3, there exist isomorphisms $E_i|_{A \times [0, 1]} = p_A^*(E_i|_{A \times \{1\}})$.

In such coordinates, the desired automorphism of $E_0 \oplus E_1$ may be given by

$$\begin{pmatrix} \cos \frac{\pi}{2}t & -i \sin \frac{\pi}{2}t \\ i \sin \frac{\pi}{2}t & \cos \frac{\pi}{2}t \end{pmatrix}$$

on $A \times [0, 1]$ and the identity on $X \times \{0\}$ (note that to specify a map of vector bundles over a stack, it suffices to specify its restriction to each member of a finite cover by closed substacks, subject to the requirement that these restrictions agree on their pairwise overlaps). \square

LEMMA 6.8. *For any pair (X, A) where A is a paracompact orbispace, the map $K^0(X, A) \rightarrow K^0(\text{cyl}(X, A))$ is surjective.*

Proof. We claim that every vector bundle on $(X \times \{0\}) \cup (A \times [0, 1])$ is pulled back from X . Let E be a vector bundle on $\text{cyl}(X, A)$, let F denote its restriction to $X = X \times \{0\}$, and let us show that $E = p_X^*F$. The identity map is an identification of E and p_X^*F over $X \times \{0\}$, and by Lemma 6.3 there is an identification of E and p_X^*F over $A \times [0, 1]$ which agrees with the identity over $A \times \{0\}$. These isomorphisms $E \rightarrow p_X^*F$ thus patch together to define the desired isomorphism.

As every vector bundle on $(X \times \{0\}) \cup (A \times [0, 1])$ is pulled back from X , it follows that $\text{Tri}(X, A) \rightarrow \text{Tri}(\text{cyl}(X, A))$ is surjective, so we are done. \square

LEMMA 6.9. *For every vector bundle E over an orbispace X , there exists an open cover $X = \bigcup_i Z_i/G_i$ such that the restriction of E to each Z_i/G_i is pulled back from $*/G_i$ (equivalently, the pullback of E to Z_i is G_i -equivariantly trivial).*

Proof. Every point $x \in X$ has an open neighborhood of the form Z/G by Proposition 3.3. Choose a lift $z \in Z$ of x , and by shrinking Z and replacing G with the stabilizer of z , assume that G fixes z . Choose a map from $E|_Z$ to the trivial bundle $E_z \times Z$ which is the identity at z , average this map to make it G -equivariant, and shrink Z so that this map is an isomorphism over all of Z . \square

LEMMA 6.10. *Let $G \curvearrowright Z$ be a finite group action on a Hausdorff topological space. If Z/G is paracompact, then so is Z .*

Proof. Let an open covering $Z = \bigcup_\alpha V_\alpha$ be given. For every $z \in Z$, there exists an open neighborhood U of z which is G_z -invariant and whose G/G_z -translates are disjoint and each contained in some (possibly different) V_α . The images of such U in Z/G form an open covering of Z/G . Refining this to a locally finite covering and taking inverse images in Z gives the desired locally finite refinement of the original covering. \square

LEMMA 6.11. *Let X be a paracompact orbispace, let E be a vector bundle over X , and let $Y \subseteq X$ be a closed substack. Every section of $E|_Y$ extends to a section of E .*

Proof. Fix an open cover $X = \bigcup_i Z_i/G_i$ such that the pullback of E to each Z_i is trivialized G_i -equivariantly (Lemma 6.9). Let $\varphi_i : |X| \rightarrow [0, 1]$ be a partition of unity subordinate to this covering. Now $\text{supp } \varphi_i$ is a closed subset of a paracompact Hausdorff space, hence paracompact Hausdorff; its inverse image inside Z_i is, thus, paracompact Hausdorff by Lemma 6.10 (recall Z_i is Hausdorff). Thus, by the Tietze extension theorem, our given section on $Z_i \times_X (Y \cap \text{supp } \varphi_i)$ extends to $Z_i \times_X \text{supp } \varphi_i$. We can make it G_i -equivariant by averaging, so our given section on $Y \cap \text{supp } \varphi_i$ extends to $\text{supp } \varphi_i$. Now $(\text{supp } \varphi_i)^\circ$ is an open cover of X , so pick another partition of unity ψ_i subordinate to this cover, and use it to patch together the extended sections on each $\text{supp } \varphi_i$. \square

LEMMA 6.12. *For any paracompact orbispace pair (X, A) , the map $K^0(X, A) \rightarrow K^0(\text{cyl}(X, A))$ is an isomorphism.*

Proof. We know from the proof of Lemma 6.8 that every vector bundle on $(X \times \{0\}) \cup (A \times [0, 1])$ is pulled back from X . Given this fact, it suffices to show that the pullback map $\text{Tri}(X, A) \rightarrow \text{Tri}(\text{cyl}(X, A))$ is injective. Thus, suppose that we are given two triples $(E_0, E_1, i : E_0|_A \rightarrow E_1|_A)$ and $(E'_0, E'_1, i' : E'_0|_A \rightarrow E'_1|_A)$ whose pullbacks to $\text{cyl}(X, A)$ coincide, meaning that there are isomorphisms $\alpha_i : p_X^* E_i \rightarrow p_X^* E'_i$ on $(X \times \{0\}) \cup (A \times [0, 1])$ intertwining i and i' over $A \times \{1\}$. Now these maps α_i extend to all of $X \times [0, 1]$ by Lemma 6.11, and these extended maps are isomorphisms over open neighborhood of $(X \times \{0\}) \cup (A \times [0, 1])$. As $X \times [0, 1] \rightarrow X$ is universally closed (being a pullback of $[0, 1] \rightarrow *$), this neighborhood contains $U \times [0, 1]$ for some open neighborhood U of $A \subseteq X$. Now a paracompact Hausdorff space is normal, so by Urysohn's lemma there exists a continuous function $\varphi : |X| \rightarrow [0, 1]$ supported inside U which is identically 1 on A . Pulling back α_i under the graph of φ defines isomorphisms $E_i \rightarrow E'_i$ whose restrictions to A intertwine i and i' , showing that the original triples on (X, A) are isomorphic. \square

PROPOSITION 6.13 (Relative homotopy invariance). *For any map of paracompact orbispace pairs $(X, A) \rightarrow (X', A')$ whose constituent maps $X \rightarrow X'$ and $A \rightarrow A'$ are individually homotopy equivalences, the map $K^0(X', A') \rightarrow K^0(X, A)$ is an isomorphism.*

Proof. We begin by showing that the two maps $K^0(X \times [0, 1], A \times [0, 1]) \rightarrow K^0(X, A)$ (pullback under $\times\{0\}$ and $\times\{1\}$) coincide (from which it follows that homotopic maps of pairs induce the same map on relative K^0). As vector bundles on $X \times [0, 1]$ are pulled back from X by Proposition 6.3, this amounts to showing that triples $(E_0, E_1, i), (E_0, E_1, j) \in \text{Tri}(X, A)$ represent the same element of $K^0(X, A)$ if i and j are homotopic. The construction from the proof of Lemma 6.7 shows that the homotopy between $i^{-1} \circ j$ and the identity gives rise to an isomorphism between the pullbacks of these triples to $\text{cyl}(X, A)$, which is enough by Lemma 6.12.

As homotopic maps of pairs induce the same map on relative K^0 , it follows that homotopy equivalences of pairs induce isomorphisms on relative K^0 . In light of Lemma 6.12, it thus suffices to show that a map of pairs $(X, A) \rightarrow (X', A')$ whose constituent maps $X \rightarrow X'$ and $A \rightarrow A'$ are individually homotopy equivalences induces a homotopy equivalence of pairs $\text{cyl}(X, A) \rightarrow \text{cyl}(X', A')$. This is well known: given $(f, g) : (X, A) \rightarrow (X', A')$, let $p : X' \rightarrow X$ and $q : A' \rightarrow A$ be homotopy inverses to f and g . To define a map $\text{cyl}(X', A') \rightarrow \text{cyl}(X, A)$ which is p on $X' \times \{0\}$ and q on $A' \times \{1\}$, we need a homotopy between the two maps $p|_{A'}, q : A' \rightarrow X$. A distinguished homotopy class of such homotopies is furnished by composing further with the homotopy equivalence $f : X \rightarrow X'$ and fixing homotopies between $f \circ p$ and $\text{id}_{X'}$ and between $f|_A \circ q = g \circ q$ and $\text{id}_{A'}$. One then checks that this map $\text{cyl}(X', A') \rightarrow \text{cyl}(X, A)$ is a homotopy inverse to the map $\text{cyl}(X, A) \rightarrow \text{cyl}(X', A')$. \square

LEMMA 6.14. *Let X be a paracompact orbispace. Every vector bundle E over a closed substack $Y \subseteq X$ is the restriction of a vector bundle over some open substack $U \subseteq X$ containing Y .*

Proof. Fix a locally finite open cover $X = \bigcup_i Z_i/G_i$ such that the pullback of E to each $(Z_i)_Y := Z_i \times_X Y \subseteq Z_i$ is trivialized G_i -equivariantly (Lemma 6.9). Such trivializations evidently extend $E|_{Y \cap (Z_i/G_i)}$ to Z_i/G_i . We patch together these extensions E_i on Z_i/G_i as follows.

Choose closed substacks $K_i \subseteq X$ with $K_i \subseteq Z_i/G_i$ and whose interiors cover X . We thus have a collection of transition functions $\alpha_{ij} : K_i \cap K_j \cap Y \rightarrow \text{Hom}(E_i, E_j)$ satisfying $\alpha_{ii} = \text{id}$ and $\alpha_{ij} \alpha_{jk} \alpha_{ki} = \text{id}$ over $K_i \cap K_j \cap K_k \cap Y$. We now execute the following operation for every pair (i, j) inductively according to an arbitrary well-ordering of such pairs. For $i = j$, do nothing. For $i \neq j$, choose an extension of α_{ij} from $K_i \cap K_j \cap Y$ to $K_i \cap K_j$ using Lemma 6.11. This extension

remains an isomorphism in a neighborhood of $K_i \cap K_j \cap Y$. As (the coarse space of) $K_i \cap K_j$ is paracompact Hausdorff, hence normal, we may choose a closed substack $A \subseteq K_i \cap K_j$ whose interior contains $K_i \cap K_j \cap Y$ and over which the extension of α_{ij} is an isomorphism. Now this new α_{ij} over A gives unique extensions of the remaining $\alpha_{k\ell}$ to $K_k \cap K_\ell \cap (Y \cup A)$ such that the cocycle condition is satisfied. We now replace Y with $Y \cup A$ and go on to the next pair of indices. Note that because our cover $X = \bigcup_i Z_i/G_i$ is locally finite, the set Y remains closed even after possibly infinitely many steps.

After processing every pair (i, j) , our extended transition functions define a vector bundle on a closed substack $\bar{Y} \subseteq X$ whose restriction to $Y \subseteq \bar{Y}$ is E , and by construction Y is contained in the interior of \bar{Y} . □

We now come to the exactness and excision properties of K -theory, where we finally make use of Corollary 1.4.

For the proof of exactness, we make use of the following notion. Let us call a map of abelian monoids $f : M \rightarrow N$ *cofinal* if and only if for every $n \in N$ there exist $m \in M$ and $n' \in N$ with $f(m) = n + n'$. The quotient M/M' is an abelian group if and only if $M' \rightarrow M$ is cofinal. The conclusion of Corollary 1.4 for a map $Y \rightarrow X$ is that the pullback map $\text{Vect}(X) \rightarrow \text{Vect}(Y)$ is cofinal (recalling from Corollary 5.2 that every inclusion of vector bundles over a paracompact orbispace is split).

PROPOSITION 6.15 (Exactness). *Let X be an orbispace satisfying the hypothesis of Theorem 1.1, let $A \subseteq X \supseteq Y$ be closed substacks, let $B := A \cap Y$ be their intersection, and let $A \cup_B Y \subseteq X$ be their union. The following sequence is exact:*

$$K^0(X, A \cup_B Y) \rightarrow K^0(X, A) \rightarrow K^0(Y, B).$$

Proof. The sequence

$$\text{Tri}(X, A \cup_B Y) \rightarrow \text{Tri}(X, A) \rightarrow (\text{Tri}(Y, B), \text{Vect}(Y))$$

is exact in the sense that the image of the first map coincides with the inverse image of $\text{Vect}(Y)$ under the second map. Indeed, given a triple $(E_0, E_1, i) \in \text{Tri}(X, A)$ for which $i|_B$ extends to Y , we may glue this extension to i over B to lift (E_0, E_1, i) from $\text{Tri}(X, A)$ to $\text{Tri}(X, A \cup_B Y)$. Quotienting this sequence by $\text{Vect}(X)$, we conclude that

$$K^0(X, A \cup_B Y) \rightarrow K^0(X, A) \rightarrow (\text{Tri}(Y, B)/\text{Vect}(X), \text{Vect}(Y)/\text{Vect}(X))$$

is exact. Now $\text{Vect}(X) \rightarrow \text{Vect}(Y)$ is cofinal by Corollary 1.4, so $\text{Vect}(Y)/\text{Vect}(X)$ is an abelian group, so the kernel of the map from $\text{Tri}(Y, B)/\text{Vect}(X)$ to its quotient by $\text{Vect}(Y)/\text{Vect}(X)$ is precisely $\text{Vect}(Y)/\text{Vect}(X)$. This quotient is simply $\text{Tri}(Y, B)/\text{Vect}(Y) = K^0(Y, B)$, so we are done. □

For the proof of excision, we make use of the following alternative description of relative K -theory in terms of direct limits. Let $\widehat{\text{Vect}}(X)$ denote the category whose objects are vector bundles on X of bounded rank and whose morphisms are homotopy classes of injective maps. The category $\widehat{\text{Vect}}(X)$ is *filtered*, meaning that (1) it is nonempty, (2) for every pair of objects x, y , there exist morphisms $x \rightarrow z \leftarrow y$, and (3) for every pair of morphisms $x \rightrightarrows y$ there exists a morphism $y \rightarrow z$ such that the two compositions $x \rightarrow z$ coincide. Indeed, (1) we have the zero vector bundle, (2) given vector bundles E and F , they both admit a morphism to $E \oplus F$, and (3) given two injections $E \hookrightarrow F$, they become homotopic after composing with the inclusion $F \hookrightarrow F \oplus E$.

For any map of stacks $f : Y \rightarrow X$ and any vector bundle F on Y , let $\text{Vect}(X, F)$ denote the set of isomorphism classes of pairs (V, i) consisting of a vector bundle V on X of bounded rank and an isomorphism $i : f^*V \xrightarrow{\sim} F$. If X is a paracompact orbispace, $\text{Vect}(X, f^*E)$ forms a directed system over $E \in \widehat{\text{Vect}}(X)$. Indeed, given an inclusion $E \hookrightarrow E'$, there is a natural map $\text{Vect}(X, f^*E) \xrightarrow{\oplus(E'/E)} \text{Vect}(X, f^*E')$ because the inclusion $E \hookrightarrow E'$ has a(n automatically unique up to homotopy) splitting by Corollary 5.2. For such X , we claim that $K^0(X, Y)$ may be expressed as the direct limit

$$K^0(X, Y) = \varinjlim_{E \in \widehat{\text{Vect}}(X)} \text{Vect}(X, f^*E).$$

Elements of the direct limit above are, by definition, equivalence classes of triples $(E, V, i) \in \text{Tri}(X, Y)$. Two triples (E, V, i) and (E', V', i') are equivalent in the above direct limit if and only if they become isomorphic after pushing both to a common $E \hookrightarrow E'' \hookleftarrow E'$, which is the same as saying (E, V, i) and (E', V', i') are isomorphic after adding elements of $\text{Vect}(X)$.

Recall that a functor F between filtered categories is called *cofinal* if and only if (1) for every x in the target there exists a morphism $x \rightarrow F(y)$, and (2) for every two morphisms $x \rightrightarrows F(y)$, there exists a morphism $y \rightarrow z$ such that the compositions $x \rightarrow F(z)$ coincide. The significance of cofinality is that pulling back under a cofinal functor induces an isomorphism on direct limits:

$$\varinjlim_{x \in C} A(F(x)) \xrightarrow{\sim} \varinjlim_{x \in D} A(x)$$

is an isomorphism for any cofinal functor $F : C \rightarrow D$ between filtered categories C and D and any directed system of sets A over D .

The conclusion of Corollary 1.4 for $f : Y \rightarrow X$ is equivalent to the assertion that the pullback functor $f^* : \widehat{\text{Vect}}(X) \rightarrow \widehat{\text{Vect}}(Y)$ is cofinal. Indeed, f^* always satisfies condition (2) because any two inclusions $E \hookrightarrow f^*F$ become homotopic upon postcomposing with $f^*(F \hookrightarrow F \oplus F)$, and condition (1) is precisely the conclusion of Corollary 1.4.

PROPOSITION 6.16 (Excision). *Let $X' \rightarrow X$ be a representable map of orbispaces satisfying the hypothesis of Theorem 1.1. Let $Y \rightarrow X$ be arbitrary, and let $Y' \rightarrow X'$ denote its pullback along $X' \rightarrow X$. If there is an open cover $X = U \cup V$ such that $Y \rightarrow X$ is an isomorphism over U and $X' \rightarrow X$ is an isomorphism over V , then the natural map $K^0(X, Y) \rightarrow K^0(X', Y')$ is an isomorphism.*

Proof. We may write the map in question in terms of direct limits as

$$\varinjlim_{E \in \widehat{\text{Vect}}(X)} \text{Vect}(X, f^*E) \rightarrow \varinjlim_{E' \in \widehat{\text{Vect}}(X')} \text{Vect}(X', (f')^*E').$$

By Corollary 1.4, we may replace the second direct limit with the corresponding direct limit over $\widehat{\text{Vect}}(X)$. It thus suffices to show that the pullback map $\text{Vect}(X, f^*E) \rightarrow \text{Vect}(X', (f')^*E')$ is an isomorphism for every vector bundle E on X of bounded rank with $E' := E \times_X X'$. Since $Y \rightarrow X$ is an isomorphism over U and $X = U \cup V$, this coincides with the map $\text{Vect}(V, f^*E|_V) \rightarrow \text{Vect}(V \times_X X', (f')^*E|_V)$, which is a bijection since $X' \rightarrow X$ is an isomorphism over V . \square

Given the significance of the hypothesis of Theorem 1.1, it is essential to show that this property is preserved under various natural operations.

LEMMA 6.17. *For any topological stack X and any locally compact Hausdorff space R , the natural map $|X \times R| \rightarrow |X| \times R$ is an isomorphism.*

Proof. For any atlas $U \rightarrow X$, the induced map $U \rightarrow |X|$ is a topological quotient map, meaning it is surjective and a subset of the target is open if and only if its inverse image in the source is open. Topological quotient maps are preserved by taking product with a locally compact Hausdorff space [Whi48, Lemma 4], so $U \times R \rightarrow |X| \times R$ is also a topological quotient map. As $U \times R \rightarrow X \times R$ is also an atlas, the map $U \times R \rightarrow |X \times R|$ is also a topological quotient map. Now $|X \times R| \rightarrow |X| \times R$ is a bijection, so we are done. \square

LEMMA 6.18. *If a topological space X is coarsely finite-dimensional, then so is $X \times R$ for any closed subset $R \subseteq \mathbb{R}^n$.*

Proof. We begin with the case $R = [0, 1]$. Let a cover $X \times [0, 1] = \bigcup_{\alpha} U_{\alpha}$ be given. For every $x \in X$ and $t \in [0, 1]$, there exists a pair of open neighborhoods $x \in V \subseteq X$ and $t \in W \subseteq [0, 1]$ such that $V \times W$ is contained in some U_{α} . Fixing $x \in X$ and using compactness of $[0, 1]$, we see that there are finitely many such $V_i \times W_i$ covering $\{x\} \times [0, 1]$. Hence, there exists an open neighborhood $x \in V \subseteq X$ and an $\varepsilon > 0$ such that $V \times ((t - \varepsilon, t + \varepsilon) \cap [0, 1])$ is contained in some U_{α} for every $t \in [0, 1]$. We consider the collection of all such pairs (V, ε) . As X is coarsely finite-dimensional, there is a collection of such pairs $(V_{\alpha}, \varepsilon_{\alpha})$ such that the nerve of the covering $X = \bigcup_{\alpha} V_{\alpha}$ is finite-dimensional. Now cover $X \times [0, 1]$ by $V_{\alpha} \times ([0, 1] \cap \varepsilon_{\alpha} \cdot (k, k + 2))$ for integers k .

Next, we consider the case $R = \mathbb{R}$. Let a cover of $X \times \mathbb{R}$ be given. The intersection of this cover with $X \times [n, n + 2]$ has a refinement with finite-dimensional nerve by the case $R = [0, 1]$. Consider this refinement intersected with $X \times (n, n + 2)$, and take union over all integers n to obtain a refinement of the original cover of $X \times \mathbb{R}$ with finite-dimensional nerve.

Finally, by induction, we obtain the case $R = \mathbb{R}^n$, and the general case follows because coarse finite-dimensionality passes to closed subsets. \square

COROLLARY 6.19. *If X satisfies the hypothesis of Theorem 1.1, then so does $X \times R$ for any closed subset $R \subseteq \mathbb{R}^n$.*

Let us now recall the Puppe sequence, which produces from Propositions 6.15 and 6.16 a long exact sequence. Let (X, A) be any pair of orbispaces satisfying the hypothesis of Theorem 1.1. Let $I := [0, 1]$. We now have maps of pairs

$$\begin{aligned} (A, \emptyset) &\rightarrow (X, \emptyset) \rightarrow (X, A) \\ (A \times I, A \times \partial I) &\rightarrow (X \times I, X \times \partial I) \rightarrow (X \times I, (X \times \partial I) \cup (A \times I)) \\ (A \times I^2, A \times \partial I^2) &\rightarrow (X \times I^2, X \times \partial I^2) \rightarrow (X \times I^2, (X \times \partial I^2) \cup (A \times I^2)) \\ &\vdots \end{aligned}$$

(note that these pairs satisfy the hypothesis of Theorem 1.1 by Corollary 6.19). We also have maps $(X \times I^k, (X \times \partial I^k) \cup (A \times I^k)) \rightarrow (A \times I^{k+1}, A \times \partial I^{k+1})$ up to inverting maps inducing isomorphisms on K^0 . Namely, these ‘connecting maps’ are given by

$$\begin{aligned} (X \times I^k \times I, (X \times \partial I^k \times I) \cup (A \times I^k \times \{1\})) \\ \downarrow \\ (X \times I^k \times I, (X \times \partial I^k \times I) \cup (A \times I^k \times \{1\}) \cup (X \times I^k \times \{0\})) \end{aligned}$$

where we note that the domain admits a natural map to $(X \times I^k, (X \times \partial I^k) \cup (A \times I^k))$ (which induces an isomorphism on K^0 by Proposition 6.13) and the target a natural map from $(A \times I^{k+1}, A \times \partial I^{k+1})$. To see that this second map also induces an isomorphism on K^0 , factor

it as

$$\begin{aligned}
 &(A \times I^k \times I, (A \times \partial I^k \times I) \cup (A \times I^k \times \{1\}) \cup (A \times I^k \times \{0\})) \\
 &\quad \downarrow \\
 &((X \times I^k \times \{0\}) \cup (A \times I^k \times I), (A \times \partial I^k \times I) \cup (A \times I^k \times \{1\}) \cup (X \times I^k \times \{0\})) \\
 &\quad \downarrow \\
 &((X \times I^k \times I), (X \times \partial I^k \times I) \cup (A \times I^k \times \{1\}) \cup (X \times I^k \times \{0\}))
 \end{aligned}$$

and observe that the first map is an isomorphism by excision Proposition 6.16 (and some manipulation involving adding $A \times I^k \times [0, \frac{1}{2}]$ to the second space of both pairs, which does not change their homotopy type but makes it possible to apply Proposition 6.16) and the second map is an isomorphism by Proposition 6.13 (a homotopy inverse to $(X \times I^k \times \{0\}) \cup (A \times I^k \times I) \rightarrow X \times I^k \times I$ is given by projection to $X \times I^k \times \{0\}$, and the second terms are both homotopy equivalent to $(X \times D^k) \cup (A \times S^k)$).

DEFINITION 6.20. For any pair (X, A) , we define²

$$\begin{aligned}
 K^{-n}(X) &:= K^0(X \times I^n, X \times \partial I^n), \\
 K^{-n}(X, A) &:= K^0(X \times I^n, (X \times \partial I^n) \cup (A \times I^n)).
 \end{aligned}$$

Given this definition, we can write the Puppe sequence above in a more familiar form. Namely, for X an orbispace satisfying the hypothesis of Theorem 1.1, we have a sequence of the form

$$\dots \rightarrow K^{-2}(A) \rightarrow K^{-1}(X, A) \rightarrow K^{-1}(X) \rightarrow K^{-1}(A) \rightarrow K^0(X, A) \rightarrow K^0(X) \rightarrow K^0(A)$$

functorial in the pair (X, A) .

PROPOSITION 6.21 (Long exact sequence). *The sequence above is exact.*

Proof. Each of the following three triples satisfies the hypotheses of Proposition 6.15:

$$\begin{aligned}
 &(A \times I^k, A \times \partial I^k) \rightarrow (X \times I^k, X \times \partial I^k) \rightarrow (X \times I^k, (X \times \partial I^k) \cup (A \times I^k)), \\
 &(X \times I^k, X \times \partial I^k) \xrightarrow{\times\{0\}} (X \times I^k \times I, (X \times \partial I^k \times I) \cup (A \times I^k \times \{1\})) \\
 &\quad \rightarrow (X \times I^k \times I, (X \times \partial I^k \times I) \cup (A \times I^k \times \{1\}) \cup (X \times I^k \times \{0\})), \\
 &(X \times I^k \times [\frac{1}{2}, 1], (X \times \partial I^k \times [\frac{1}{2}, 1]) \cup (A \times I^k \times \{1\})) \\
 &\quad \rightarrow (X \times I^k \times I, (X \times \partial I^k \times I) \cup (X \times I^k \times \{0\}) \cup (A \times I^k \times \{1\})) \\
 &\quad \rightarrow (X \times I^k \times I, (X \times \partial I^k \times I) \cup (X \times I^k \times \{0\}) \cup (X \times I^k \times [\frac{1}{2}, 1]))
 \end{aligned}$$

and the last pair is homotopy equivalent to $(X \times I^k \times [0, \frac{1}{2}], (X \times \partial I^k \times [0, \frac{1}{2}]) \cup (X \times I^k \times \{0\}) \cup (X \times I^k \times \{\frac{1}{2}\}))$. □

We now define a multiplicative structure on K -theory. There is a natural map $K^0(X) \otimes K^0(Y) \rightarrow K^0(X \times Y)$ sending $[E] \otimes [F]$ to $[E \boxtimes F]$. To define the tensor product map on relative K -theory, we consider the following alternative model based on chain complexes.

DEFINITION 6.22. Let (X, A) be a pair. Consider bounded complexes of vector bundles on X which are exact over A . Let us call two such complexes homotopic if and only if they are the restrictions to $\times\{0\}$ and $\times\{1\}$ of a complex on $X \times [0, 1]$ acyclic over $A \times [0, 1]$ (homotopy is

² To keep track of signs, one should really write this as $K^{-n}(X) \otimes H^n(I^n, \partial I^n) = K^0(X \times I^n, X \times \partial I^n)$ etc., however for the purposes of our presentation here, we will not be so precise.

an equivalence relation). Denote the set of homotopy classes of such complexes by $\text{Kom}(X, A)$, which is an abelian monoid under direct sum, and define

$$K_{\text{kom}}^0(X, A) := \text{Kom}(X, A) / \text{Kom}(X, X).$$

This quotient is an abelian group: indeed, the sum of a complex and its shift is homotopic to the mapping cone of the identity map.

PROPOSITION 6.23. *There is a natural isomorphism $K^0(X, A) = K_{\text{kom}}^0(X, A)$ for paracompact orbispace pairs (X, A) .*

Proof. There is a natural map $K^0(X, A) \rightarrow K_{\text{kom}}^0(X, A)$ defined as follows. Given a triple $(E_0, E_1, i) \in \text{Tri}(X, A)$, extend the isomorphism i to a neighborhood of A using Lemma 6.11, and multiply by a cutoff function supported inside this neighborhood to obtain a globally defined map $d : E_0 \rightarrow E_1$ which is an isomorphism over A . This complex is well defined up to homotopy, so we have defined a map $\text{Tri}(X, A) \rightarrow \text{Kom}(X, A)$. This map evidently sends $\text{Vect}(X)$ to (the image of) $\text{Kom}(X, X)$, so it defines a map $K^0(X, A) \rightarrow K_{\text{kom}}^0(X, A)$ as desired.

We may also define a natural map $K_{\text{kom}}^0(X, A) \rightarrow K^0(X, A)$ as follows. Given a complex (E_\bullet, d) , choose a metric on each E_i using Lemma 5.1, and note that the map $d + d^* : E_{\text{even}} := \bigoplus_{i \text{ even}} E_i \rightarrow \bigoplus_{i \text{ odd}} E_i =: E_{\text{odd}}$ is an isomorphism wherever d is exact. As all metrics are homotopic and $K^0(X, A)$ is homotopy invariant by Proposition 6.13, sending (E_\bullet, d) to the triple $(E_{\text{even}}, E_{\text{odd}}, (d + d^*)|_A)$ gives a well-defined map $K_{\text{kom}}^0(X, A) \rightarrow K^0(X, A)$. The composition $K^0(X, A) \rightarrow K_{\text{kom}}^0(X, A) \rightarrow K^0(X, A)$ is evidently the identity.

To finish, it suffices to show that $K^0(X, A) \rightarrow K_{\text{kom}}^0(X, A)$ is surjective. Equivalently, we are to show that every element of $K_{\text{kom}}^0(X, A)$ is represented by a complex concentrated in degrees $[0, 1]$. To do this, we use the following ‘folding’ operation. Given a complex concentrated in degrees $[s, r]$, the differential $d_r : E_r \rightarrow E_{r-1}$ is injective over A , hence over an open neighborhood U of A inside X . If d_r is, in fact, everywhere injective, then we may choose a metric on E_{r-1} and split it as $\text{im}(E_r) \oplus \text{im}(E_r)^\perp$, showing that the subcomplex $E_r \rightarrow \text{im}(E_r)$ is a direct summand, so our complex represents the same class in K_{kom}^0 as one concentrated in degrees $[s, r - 1]$. Now consider the general case where d_r is not assumed everywhere injective. By replacing our given complex by its direct sum with the mapping cone of the identity map $E_r \rightarrow E_r$ concentrated in degrees $[r - 2, r - 1]$, we may assume that there exists an injection $i : E_r \rightarrow E_{r-1}$ which agrees with d_r over a neighborhood U of A . Now multiply d by a cutoff function φ supported inside U and positive on A (this is homotopic to the original differential), and take the convex interpolation between φd_r and i as a further homotopy. This ensures that the differential $E_r \rightarrow E_{r-1}$ is everywhere injective, so our complex represents the same class in K_{kom}^0 as one concentrated in degrees $[\min(s, r - 2), r - 1]$. The dual operation shows that any complex concentrated in degrees $[s, r]$ represents the same class in K_{kom}^0 as one concentrated in degrees $[s + 1, \max(s + 2, r)]$. Combining these two operations, we can put anything in degrees $[0, 1]$. \square

Tensor product at the level of K_{kom}^0 evidently defines commutative and associative maps

$$K^0(X, A) \otimes K^0(Y, B) \rightarrow K^0(X \times Y, (A \times Y) \cup (X \times B))$$

for paracompact orbispace pairs (X, A) and (Y, B) for which $X \times Y$ is paracompact. These induce, by inspection, associative and graded commutative maps

$$K^{-n}(X, A) \otimes K^{-m}(Y, B) \rightarrow K^{-n-m}(X \times Y, (A \times Y) \cup (X \times B))$$

under the same hypotheses (note that a paracompact space times a compact space is paracompact).

We conclude with a discussion of Bott periodicity. So far, our discussion has applied equally to complex vector bundles as to real vector bundles, however we now restrict to complex vector bundles. The *Bott element* $\beta \in K^{-2}(\mathbb{C}) = K^0(D^2, \partial D^2)$ is represented by the complex of vector bundles $\underline{\mathbb{C}} \xrightarrow{z} \underline{\mathbb{C}}$ on \mathbb{C} (which contains D^2 as the unit disk).

PROPOSITION 6.24. *For any finite simplicial complex of groups X with subcomplex A , multiplication by the Bott element $K^{-k}(X, A) \rightarrow K^{-k-2}(X, A)$ is an isomorphism for all $k \geq 0$.*

Proof. By the long exact sequence, excision, the five lemma, and finite additivity, we are reduced to the case of $(D^i, \partial D^i) \times \mathbb{B}G$. By the definition of K^{-n} , we are further reduced to the case of K^0 , namely to showing that multiplication by the Bott element

$$K^0((D^k, \partial D^k) \times \mathbb{B}G) \rightarrow K^0((D^{k+2}, \partial D^{k+2}) \times \mathbb{B}G)$$

is an isomorphism for all $k \geq 0$, which is well known (see [AB64], [Ati68, Theorem 4.3], and [Seg68, §3]). □

In fact, this argument shows more generally that for any complex line bundle L over a finite simplicial complex of groups X , pullback followed by multiplication with the relative Bott class in $K^0(L, \infty)$ (represented by the complex $\underline{\mathbb{C}} \xrightarrow{z} L$ on L) defines an isomorphism $K^{-k}(X) \rightarrow K^{-k}(L, \infty)$ (and, yet more generally, that pullback and multiplication with the Koszul complex defines an isomorphism $K^{-k}(X) \rightarrow K^{-k}(E, \infty)$ for any complex vector bundle E).

DEFINITION 6.25. The periodic K -theory of a pair (X, A) is the direct limit

$$K_{\text{per}}^*(X, A) := \varinjlim_j K^{*-2j}(X, A)$$

over multiplication by β . Whereas K^* is defined only in nonpositive degrees, K_{per}^* is defined in all degrees.

As direct limits are exact, K_{per}^* is, similarly to K^* , a cohomology theory for orbispace pairs satisfying the hypothesis of Theorem 1.1. When the natural map $K^* \rightarrow K_{\text{per}}^*$ is an isomorphism, K_{per}^* provides a natural extension of K^* from nonpositive degrees to all degrees.

COROLLARY 6.26. *For any finite simplicial complex of groups X with subcomplex A , the natural map $K^*(X, A) \rightarrow K_{\text{per}}^*(X, A)$ is an isomorphism in nonpositive degrees.*

ACKNOWLEDGEMENTS

I would like to thank André Henriques for discussions about orbispaces and topological stacks, David Treumann for conversations about the algebraic case, and the referees for their remarks. I also thank the Department of Mathematics at Cambridge University for their hospitality during the time when this paper was conceived.

REFERENCES

AR03 A. Adem and Y. Ruan, *Twisted orbifold K-theory*, Comm. Math. Phys. **237** (2003), 533–556; MR 1993337.

Ati68 M. F. Atiyah, *Bott periodicity and the index of elliptic operators*, Q. J. Math. Oxford Ser. (2) **19** (1968), 113–140; MR 228000.

Ati69 M. F. Atiyah, *Algebraic topology and operators in Hilbert space*, Lectures in Modern Analysis and Applications. I (Springer, Berlin, 1969), 101–121; MR 0248803.

AB64 M. Atiyah and R. Bott, *On the periodicity theorem for complex vector bundles*, Acta Math. **112** (1964), 229–247; MR 178470.

- AS69 M. F. Atiyah and I. M. Singer, *Index theory for skew-adjoint Fredholm operators*, Publ. Math. Inst. Hautes Études Sci. **37** (1969), 5–26; [MR 285033](#).
- Beh14 K. Behrend, *Introduction to algebraic stacks*, in *Moduli spaces*, London Mathematical Society Lecture Note Series, vol. 411 (Cambridge University Press, Cambridge, 2014), 1–131; [MR 3221293](#).
- BN06 K. Behrend and B. Noohi, *Uniformization of Deligne–Mumford curves*, J. Reine Angew. Math. **599** (2006), 111–153; [MR 2279100](#).
- BH99 M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, in *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 319 (Springer, Berlin, 1999); [MR 1744486](#).
- BT17 A. D. Brooke-Taylor, *Products of CW complexes*, Preprint (2017), [arXiv:1710.05296](#).
- CTS21 J.-L. Colliot-Thélène and A. N. Skorobogatov, *The Brauer–Grothendieck group*, in *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, vol. 71 (Springer, Cham, 2021); [MR 4304038](#).
- Cor92 J. M. Corson, *Complexes of groups*, Proc. Lond. Math. Soc. (3) **65** (1992), 199–224; [MR 1162493](#).
- dJ03 A. J. de Jong, *A result of Gabber*, Preprint (2003), <https://www.math.columbia.edu/dejong/>.
- Die44 J. Dieudonné, *Une généralisation des espaces compacts*, J. Math. Pures Appl. (9) **23** (1944), 65–76; [MR 0013297](#).
- EHKV01 D. Edidin, B. Hassett, A. Kresch and A. Vistoli, *Brauer groups and quotient stacks*, Amer. J. Math. **123** (2001), 761–777; [MR 1844577](#).
- Hae90 A. Haefliger, *Orbi-espaces*, in *Sur les groupes hyperboliques d’après Mikhael Gromov (Bern, 1988)*, Progress in Mathematics, vol. 83 (Birkhäuser, Boston, MA, 1990), 203–213; [MR 1086659](#).
- Hae91 A. Haefliger, *Complexes of groups and orbihedra*, in *Group theory from a geometrical viewpoint* (World Scientific Publishing, River Edge, NJ, 1991), 504–540; [MR 1170375](#).
- Hei05 J. Heinloth, *Notes on differentiable stacks*, in *Mathematisches Institut, Georg-August-Universität Göttingen: Seminars winter term 2004/2005* (Universitätsdrucke Göttingen, Göttingen, 2005), 1–32; [MR 2206877](#).
- HG07 A. Henriques and D. Gepner, *Homotopy theory of orbispaces*, Preprint (2007), [arXiv:math/0701916](#).
- HM04 A. Henriques and D. S. Metzler, *Presentations of noneffective orbifolds*, Trans. Amer. Math. Soc. **356** (2004), 2481–2499; [MR 2048526](#).
- Hen05 A. G. Henriques, *Orbispaces*, PhD thesis, Massachusetts Institute of Technology (2005).
- Kre09 A. Kresch, *On the geometry of Deligne–Mumford stacks*, in *Algebraic geometry—Seattle 2005. Part 1*, Proceedings of Symposia in Pure Mathematics, vol. 80 (American Mathematical Society, Providence, RI, 2009), 259–271; [MR 2483938](#).
- KV04 A. Kresch and A. Vistoli, *On coverings of Deligne–Mumford stacks and surjectivity of the Brauer map*, Bull. Lond. Math. Soc. **36** (2004), 188–192; [MR 2026412](#).
- LO01 W. Lück and B. Oliver, *The completion theorem in K-theory for proper actions of a discrete group*, Topology **40** (2001), 585–616; [MR 1838997](#).
- LMB00 G. Laumon and L. Moret-Bailly, *Champs algébriques*, in *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, vol. 39 (Springer, Berlin, 2000); [MR 1771927](#).
- Mat11 A. Mathew, *Bott periodicity and integrality theorems*, <https://amathew.wordpress.com/2011/12/28/bott-periodicity-and-integrality-theorems/>.
- Mat71 T. Matumoto, *Equivariant K-theory and Fredholm operators*, J. Fac. Sci. Univ. Tokyo Sect. I A Math. **18** (1971), 109–125; [MR 0290354](#).

- May67 J. P. May, *Simplicial objects in algebraic topology*, Van Nostrand Mathematical Studies, No. 11 (Van Nostrand, Princeton, NJ, 1967); [MR 0222892](#).
- Met03 D. Metzler, *Topological and smooth stacks*, Preprint (2003), [arXiv:math/0306176](#).
- Mil56 J. Milnor, *Construction of universal bundles. I*, Ann. of Math. (2) **63** (1956), 272–284; [MR 77122](#).
- Noo05 B. Noohi, *Foundations of topological stacks I*, Preprint (2005), [arXiv:math/0503247](#).
- Sch18 S. Schwede, *Global homotopy theory*, New Mathematical Monographs, vol. 34 (Cambridge University Press, Cambridge, 2018); [MR 3838307](#).
- Sch20 S. Schwede, *Orbispace, orthogonal spaces, and the universal compact Lie group*, Math. Z. **294** (2020), 71–107; [MR 4050064](#).
- Seg68 G. Segal, *Equivariant K-theory*, Publ. Math. Inst. Hautes Études Sci. **34** (1968), 129–151; [MR 234452](#).
- Seg70 G. Segal, *Fredholm complexes*, Q. J. Math. Oxford Ser. (2) **21** (1970), 385–402; [MR 271930](#).
- Sie98 B. Siebert, *Gromov–Witten invariants for general symplectic manifolds*, Preprint (1998), [arXiv:dg-ga/9608005v2](#).
- Tot04 B. Totaro, *The resolution property for schemes and stacks*, J. Reine Angew. Math. **577** (2004), 1–22; [MR 2108211](#).
- Whi48 J. H. C. Whitehead, *Note on a theorem due to Borsuk*, Bull. Amer. Math. Soc. (N.S.) **54** (1948), 1125–1132; [MR 29503](#).
- Whi49 J. H. C. Whitehead, *Combinatorial homotopy. I*, Bull. Amer. Math. Soc. (N.S.) **55** (1949), 213–245; [MR 30759](#).

John Pardon

Department of Mathematics, Princeton University, Princeton, NJ 08544, USA