

NRO Legacy Project: M33 all Disk Survey of Giant Molecular Clouds with NRO 45-m and ASTE 10-m telescopes

N. Kuno^{1,2}, T. Tosaki³, S. Onodera¹, R. Miura⁴, K. Muraoka⁵,
S. Komugi⁶, T. Sawada⁶, K. Nakanishi⁶, K. Kohno^{7,8}, H. Kaneko^{1,2},
A. Hirota¹, N. Arimoto^{2,6}, H. Nakanishi⁹, and R. Kawabe¹

¹Nobeyama Radio Observatory, National Astronomical Observatory of Japan,
Minamimaki, Minamisaku, Nagano, 384-1305, Japan

²The Graduate University for Advanced Studies (SOKENDAI),
Osawa, Mitaka, Tokyo, 181-8588, Japan

³Joetsu University of Education, Yamayashiki-machi, Joetsu, Niigata, 943-8512, Japan

⁴University of Tokyo, Department of Astronomy, School of Science,
Hongo, Bunkyo, Tokyo, 113-0033, Japan

⁵Department of Physical Science, Osaka Prefecture University,
Gakuen 1-1, Sakai, Osaka, 599-8531, Japan

⁶National Astronomical Observatory of Japan, Osawa, Mitaka, Tokyo, 181-8588, Japan

⁷Institute of Astronomy, School of Science, The University of Tokyo,
Osawa, Mitaka, Tokyo, 181-0015, Japan

⁸Research Center for Early Universe, School of Science, The University of Tokyo,
Hongo, Bunkyo, Tokyo, 113-0033, Japan

⁹Graduate School of Science and Engineering, Kagoshima University,
1-21-35 Korimoto, Kagoshima, Kagoshima 890-0065, Japan

Abstract. We have conducted all disk imaging of M33 in $^{12}\text{CO}(1-0)$ using the 45-m telescope at Nobeyama Radio Observatory. We present preliminary results of this project. The spatial resolution of ~ 80 pc is comparable to the size of GMCs. The identified GMCs show wide variety in star forming activity. The variety can be regarded as the difference of their evolutionary stage. We found that Kennicutt-Schmidt law breaks in GMC scale (~ 80 pc), although it is still valid in 1 kpc scale. The correlation between molecular gas fraction, $f_{mol} = \Sigma(\text{H}_2)/\Sigma(\text{HI}+\text{H}_2)$ and gas surface density shows two distinct sequences and shows that f_{mol} tends to be higher near the center. We also made partial mapping $^{12}\text{CO}(3-2)$ with ASTE telescope. These data show that the variation of physical properties of molecular gas are correlated with the GMC evolution and mass. That is, GMCs with more active star formation and more mass tend to have higher fraction of dense gas.

Keywords. galaxies: individual(M33), ISM, GMC

1. Introduction

Since M33 is one of the nearest spiral galaxies in the local group and moderately face-on, the galaxy is suitable to resolve each GMCs and investigate their properties. Furthermore, there are many other frequency data of tracers of atomic gas, star forming regions, and stars, for comparisons. Therefore, M33 is the best target for studying GMCs and star formation within a whole galaxy. Our observations are performed with Nobeyama 45m and ASTE telescopes. The 25 multi-beam receiver, BEARS, mounted

on the 45m telescope and CATS345 installed in ASTE were used. We can get $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(3-2)$ data with highly uniform quality by using these receivers and On-The-Fly technique. Our scientific objectives are to investigate the basic properties of GMCs and their spatial variation, and to understand the evolutionary process of GMCs and massive star formation. We will make GMC catalog with the information about their evolutionary stage using our data and other frequency data (e.g., optical data with SUBARU). For these purposes, we have to observe the variation of the internal structure of GMCs along their evolutionary stage eventually. Observations of the internal structure of GMCs in M33 will be achieved by ALMA.

2. Preliminary results

Fig. 1 shows the total integrated intensity map of $^{12}\text{CO}(1-0)$ and intensity weighted mean velocity map. Many GMCs are identified in the map and global galactic rotation can be seen. Fig. 2 shows the comparison between CO and HI, and CO and star formation rate. We can see the variation of evolutionary stage of GMCs in fig. 2. Namely, there are some GMCs with no star forming activity and HII regions without associated GMCs. GMCs are located on the HI peaks in the outer region, but not in the inner region. We present some preliminary results of our project below.

2.1. Breakdown of Kennicutt-Schmidt law

Kennicutt-Schmidt law is a well-known correlation between surface gas density and star formation rate in galaxies. But, it is not clear in what scale the Kennicutt-Schmidt law breaks down. To answer the question, Onodera *et al.* (2010) compared the surface density of molecular gas and that of the star formation rate in various spatial resolution (fig. 3). These plots show that the correlation is evident in 1 kpc and 500 pc resolution. The correlation becomes looser with higher resolution and finally breaks down in 80 pc resolution. These results indicate that the resolution of 80pc, which is comparable to GMC scales, is the scale in which the Kennicutt-Schmidt law becomes invalid. The comparison between distributions of CO and star forming regions implies that the main cause of the breakdown is that the variation of evolutionary stage of GMCs becomes apparent in this scale as seen in fig. 2.




Figure 1. $^{12}\text{CO}(1-0)$ integrated intensity map of M33 (left) and velocity field (right). (Tosaki *et al.* 2011)




Figure 2. The $^{12}\text{CO}(1-0)$ map superposed on HI (Rosolowsky *et al.* 2007) (left) and star formation rate derived from H α luminosity (Hoopes and Walterbos 2000) with extinction-correction using 24 μm data (Rieke *et al.* 2004) (right). (Tosaki *et al.* 2011)

2.2. HI to H $_2$ transition

Tosaki *et al.* (2011) make comparison between the CO and HI maps and show that molecular gas fraction f_{mol} is higher in the inner region than in the outer region (fig. 2). Fig. 4 shows correlation between gas surface density and f_{mol} . It is apparent that there




Figure 3. Star formation rate per unit area vs. surface density of H $_2$ gas for four different resolutions. The broken lines represent 2σ of surface density of H $_2$. The line in d) shows the best fit to the ~ 1 kpc resolution data. (Onodera *et al.* 2010)




Figure 4. Correlation between gas surface density and f_{mol} . Data of $R \leq 1.5$ kpc, $1.5 \text{ kpc} \leq R \leq 3 \text{ kpc}$, and $R \geq 3 \text{ kpc}$. Dotted and dashed curves indicate correlation for $U = 10U_0$ in the case of $Z = Z_0$ and $Z = 0.5Z_0$ (left), and for $Z = 1.0Z$ in the case of $U = 10U_0$ and $U = 48Z_0$ (right), respectively. (Tosaki *et al.* 2011)

are two distinct sequences depending on the distance from the galactic center. f_{mol} is higher in the inner region $R < 1.5$ kpc than in the outer region at the same gas surface density. Most probable cause of the difference of f_{mol} is high metallicity in the inner region. Actually, sharp increase of metallicity in the central 1 kpc region in M33 has been reported by Vilchez *et al.* (1988). On the other hand, if we attribute the difference to the variation of radiation field, radiation field has to increase with distance from the center. But, radial distribution of tracers of stars and star forming regions does not show such a trend. Another possibility is variation of gas volume density. In fig. 4, gas surface density is used as an indicator of gas volume density assuming that the scale height is constant. But, if the scale height is larger in the outer region than in the inner region as seen in our Galaxy, volume density is higher in the inner region at the same surface density.

2.3. Variation of CO(1-0)/CO(3-2) ratio in GMCs

Onodera (2009) and Onodera *et al.* (2011) investigated the variation of CO(1-0)/CO(3-2) ratio in GMCs and its dependence on the GMC properties. They show that the ratio depends on star forming activity in GMCs and GMC mass. Namely, GMCs with more active star formation and more massive tend to have higher fraction of dense gas.

References

- Hoopes, C. G. & Walterbos, R. A. M. 2000, *ApJ*, 541, 597
 Onodera, S. 2009, PhD thesis, University of Tokyo
 Onodera, S., *et al.* 2010, *ApJ(Letters)* 722, L127
 Onodera, S., *et al.* 2011, in prep.
 Rieke, G. H., *et al.* 2004, *ApJS*, 154, 25
 Tosaki, T., *et al.* 2011, *PASJ*, submitted
 Vilchez, J. M., *et al.* 1988, *MNRAS*, 235, 633