

CONTRACTED IDEALS FROM INTEGRAL EXTENSIONS OF REGULAR RINGS

M. HOCHSTER¹

0. Introduction. The purpose of this paper is to consider the following question: if R is a regular Noetherian ring and $S \supset R$ is a module-finite R -algebra, is R a direct summand of S as R -modules? An affirmative answer is given if R contains a field, and it is shown that if the completions of the local rings of S possess maximal Cohen-Macaulay modules in the sense of §1 of [6] then the conclusion is valid in this case too. Hence, if Conjecture E of [6] is true then the question raised here has an affirmative answer without further restriction on the regular Noetherian ring R , and it will be shown here that only a greatly weakened version of Conjecture E is needed.

It follows from our results on direct summands that if R contains a field, its local rings are regular, and S is an extension algebra integral over R , then every ideal of R is contracted, i.e. if $I \subset R$ then $IS \cap R = I$. In fact, we prove the direct summand result by proving first that certain key ideals I of a regular local ring have this property. In the final section of the paper we consider briefly some propositions about the class of domains such that every ideal is contracted from every integral extension.

Throughout this paper, all rings are commutative, with identity, all modules are unitary, and ring homomorphisms are assumed to preserve the identity.

1. Regular rings. Our first reductions in the problem are consequences of the following:

LEMMA 1. *Let $R \subset S$ be rings and assume that S is finitely presented as an R -module. Then R is a direct summand of S if and only if for*

Received November 6, 1972.

¹This research was partially supported by NSF grant GP-29224X.

each maximal ideal P of R , R_P is a direct summand of S_P .

Moreover, if T is a faithfully flat R -algebra then R is a direct summand of S if and only if $T = R \otimes_R T$ is a direct summand of $S \otimes_R T$.

Proof. Consider the exact sequence

$$0 \rightarrow R \rightarrow S \rightarrow S/R \rightarrow 0.$$

To ask whether R is a direct summand of S is the same as to ask whether

$$\text{Hom}_R(S/R, S) \rightarrow \text{Hom}_R(S/R, S/R)$$

is surjective. This map is surjective if and only if for each maximal ideal P of R ,

$$\text{Hom}_R(S/R, S) \otimes_R R_P \rightarrow \text{Hom}_R(S/R, S/R) \otimes_R R_P$$

is surjective, and since S (and hence S/R) is finitely presented, we have the commutative diagram:

$$\begin{array}{ccc} \text{Hom}_R(S/R, S) \otimes_R R_P & \cong & \text{Hom}_{R_P}(S_P/R_P, S_P) \\ \downarrow & & \downarrow \\ \text{Hom}_R(S/R, S/R) \otimes_R R_P & \cong & \text{Hom}_{R_P}(S_P/R_P, S_P/R_P). \end{array}$$

The first part of the theorem follows at once.

For the second part we note that since T is faithfully flat

$$\text{Hom}_R(S/R, S) \rightarrow \text{Hom}_R(S/R, S/R)$$

is surjective if and only if

$$\text{Hom}_R(S/R, S) \otimes_R T \rightarrow \text{Hom}_R(S/R, S/R) \otimes_R T$$

is surjective, and since $S, S/R$ are finitely presented and T is flat we have the commutative diagram:

$$\begin{array}{ccc} \text{Hom}_R(S/R, S) \otimes_R T & \cong & \text{Hom}_T(S \otimes T/R \otimes T, S \otimes T) \\ \downarrow & & \downarrow \\ \text{Hom}_R(S/R, S/R) \otimes_R T & \cong & \text{Hom}_T(S \otimes T/R \otimes T, S \otimes T/R \otimes T) \end{array}$$

and so the second part follows.

Remark 1. The criteria of Lemma 1 apply to determine whether

an arbitrary short exact sequence of finitely presented R -modules is split, not just

$$0 \rightarrow R \rightarrow S \rightarrow S/R \rightarrow 0.$$

This is evident from the proof.

PROPOSITION 1. *If $R \subset S$ are rings such that R is a direct summand of S then for each ideal I of R , $IS \cap R = I$.*

Proof. See [7], Proposition 10.

The condition that R be a direct summand of an extension ring S is equivalent, in the terminology of [7], to the existence of a Reynolds operator from S to R . The consequences of the existence of a Reynolds operator are discussed in detail in [7] Propositions 9–12. Such operators occur frequently in invariant theory over fields of characteristic zero.

We note that if R is a domain which is a direct summand of every module-finite extension, then R is integrally closed. For if $b, a \neq 0$ are in R and b/a is integral over R , then since $R[b/a]$ is module-finite over R we have $b \in (aR[b/a]) \cap R = aR$ and $b/a \in R$.

The following two results show that if R contains the rationals integral closure is also sufficient. The next result is well known but we include a short proof for the sake of completeness.

LEMMA 2. *If R is an integrally closed domain which contains the rationals Q and S is an integral extension domain of R of finite degree, then R is a direct summand of S .*

Proof. There is no harm in enlarging S . Hence, we might as well assume that S is the integral closure of R in a finite normal field extension L of the fraction field K of R . Let $d = [L : K]$. Then $(1/d) \operatorname{Tr}_{L/K}$ retracts S onto R . In fact, if $r \in R$ (or K) $\operatorname{Tr}_{L/K}(r) = dr$. On the other hand, the trace of an integral element is integral and every element of K integral over R is in R .

Lemma 2 shows that the hypothesis of regularity is much more than sufficient in case R contains the rationals. To dispel the idea that anything much weaker than regularity might be enough in characteristic p we give an example.


EXAMPLE 1. Let K be an algebraically closed field of characteristic 2. Let u, v, x, t , and z be indeterminates. Let $D = K[u^2, v^2, u^3 + v^3] \subset$

$K[u, v]$. The K -homomorphism from $K[x, y, z]$ onto D which takes x, y, z to $u^2, v^2, u^3 + v^3$, respectively, has as its kernel the principal ideal $(x^3 + y^3 + z^2)$, so that D is Gorenstein (and, in fact, a complete intersection), and it is easy to see that D is integrally closed in its fraction field. In D , $u^3 + v^3 \notin (u^2, v^2)D$, but in $K[u, v]$, which is integral over D , $u^3 + v^3 \in (u^2, v^2)K[u, v]$. Thus, (u^2, v^2) is not contracted from $K[u, v]$, and it follows that D is *not* a direct summand of its integral extension $K[u, v]$, by Proposition 1. The essential properties of this example are preserved by localizing at the maximal ideals $((u, v) \cap D$ and (u, v)) and even by, furthermore, completing the resulting local rings. Hence, even an analytically normal Gorenstein algebro-geometric local ring fails to be a summand of all of its module-finite extensions in characteristic p . The author has not found any natural condition weaker than regularity which suffices (although regularity is not necessary).

The next lemma shows that we only need to look at domain extensions.

LEMMA 3. *Let R be a domain and S an integral extension of R . Then there is a prime P of S such that $P \cap R = (0)$, and if R is a direct summand of S/P (under the induced inclusion), then R is a direct summand of S .*

Proof. Since $R - \{0\}$ is a multiplicative system in S we can take P to be any ideal of S maximal with respect to disjointness from $R - \{0\}$. We have a commutative diagram:



and it follows that if $f: S/P \rightarrow R$ is an R -module retraction then $fg: S \rightarrow R$ is an R -module retraction.

Suppose we are given, now, a regular Noetherian ring R and a module-finite R -algebra $S \supset R$, and we want to decide whether R is a direct summand of S . By using the first part of Lemma 1 we can reduce to the case where R is local, and by using the second part to the case where R is complete, taking $T = \hat{R}$, the completion of R . Using Lemma 3 we can reduce to the case where S is a domain. However, for the moment, we shall not make all of these reductions but we give

instead a criterion for R to be a direct summand of S when R is a regular local ring and S is module-finite over R .

THEOREM 1. *Let (R, P) be a regular local ring and let x_1, \dots, x_n be a regular system of parameters for R (i.e. $\dim R = n$ and $P = (x_1, \dots, x_n)R$). Let $R \subset S$, where S is a module-finite R -algebra. Then R is a direct summand of S if and only if for every positive integer k ,*

$$(x_1 \cdots x_n)^k \notin (x_1^{k+1}, \dots, x_n^{k+1})S$$

Proof. Let $I_k = (x_1^k, \dots, x_n^k)R$. Since R is regular and x_1^k, \dots, x_n^k is a system of parameters, $R_k = R/I_k$ is Gorenstein and zero-dimensional: let $P_k = P/I_k$. Then $\text{Ann}_{R_k} P_k$ will be isomorphic to a single copy of R/P , and every nonzero ideal of R_k will contain $\text{Ann}_{R_k} P_k$. It is quite easy to see, in fact, that $\text{Ann}_{R_k} P_k$ is generated by the residue class modulo I_k of $u_k = (x_1 \cdots x_n)^{k-1}$. It follows that every ideal of R strictly larger than I_k contains u_k .

Now, if R is a direct summand of S then every ideal of R will be contracted. Since $u_{k+1} \notin I_{k+1}R$, we will then have $u_{k+1} \notin I_{k+1}S$, which is precisely the condition asserted in Theorem 1.

To prove the converse, assume that $u_{k+1} \notin I_{k+1}S$, all k . Let \hat{R} be the completion of R . By the second statement of Lemma 1, it suffices to show that \hat{R} is a direct summand of $S \otimes_R \hat{R}$, which is a module-finite extension algebra of \hat{R} . Since \hat{R} is faithfully flat over R , $S \otimes_R \hat{R}$ is faithfully flat over S . Then $u_{k+1} \notin (I_{k+1}(S \otimes_R \hat{R})) \cap \hat{R}$, because if it did, we would have $u_{k+1} \in ((I_{k+1}(S \otimes_R \hat{R})) \cap S) \cap \hat{R}$ and since $S \otimes_R \hat{R}$ is faithfully flat over S , $(I_{k+1}(S \otimes_R \hat{R})) \cap S = I_{k+1}S$ and $u_{k+1} \in (I_{k+1}S \cap R) \cap R = (I_{k+1}S) \cap R$, contradiction. Hence, we may assume without loss of generality that R is complete.

Since the ideals I_k are cofinal with the powers of P and R is complete,

$$(*) \quad \text{Hom}_R(S, R) = \lim \text{Hom}_R(S_k, R_k)$$

where $S_k = S/I_kS$. We also note the isomorphisms

$$\text{Hom}_R(S_k, R_k) \cong \text{Hom}_{R_k}(S_k, R_k).$$

Now, $I_kS \cap R = I_k$, because if $J = I_kS \cap R$ and $J \supsetneq I_k$, then $u_k \in J$, and this contradicts our hypothesis. Hence, the inclusion $R \rightarrow S$ induces an inclusion $R_k \rightarrow S_k$ for each k . Since R_k is a zero-dimensional Gorenstein

ring, it is injective as an R_k -module, and it follows that for each k , the inclusion $R_k \rightarrow S_k$ splits, i.e. R_k is a direct summand of S_k . For each k , let $h_k: \text{Hom}_R(S_k, R_k) \rightarrow R_k$ by $h_k(\phi) = \phi(1_k)$ where $1_k \in R_k \subset S_k$ is the image of $1 \in R \subset S$, and let $h: \text{Hom}_R(S, R) \rightarrow R$ by $h(\phi) = \phi(1)$. Then $H_k = h_k^{-1}(1)$ (respectively, $H = h^{-1}(1)$) is the set of splittings of $R_k \rightarrow S_k$ (respectively, $R \rightarrow S$) and the inverse limit relation $(*)$ induces

$$(**) \quad H = \lim_{\leftarrow k} H_k .$$

Here H_k (respectively, H) is a coset (or translate) of a submodule of $\text{Hom}_R(S, R_k)$ (respectively, $\text{Hom}_R(S, R)$) and the maps are restricted module homomorphisms. All we need to show to complete the proof is that $H \neq \phi$. Since each R_k is a summand of S_k , each $H_k \neq \phi$. But an inverse limit of nonempty cosets in Artinian modules is nonempty. To see this, note that for each k the decreasing sequence of nonempty subcosets $\text{Im}(H_{i+k} \rightarrow H_k)$ of H_k stabilizes, since their lengths must stabilize. Denote this subset of H_k by E_k . Then the E_k form a subsystem of nonempty subcosets and *surjective* maps so that

$$\phi \neq \lim_{\leftarrow k} E \subset H .$$

This completes the proof.

Remark 2. Theorem 1 generalizes in various ways. We indicate a number of these generalizations. First, it is not necessary to assume that S is an R -algebra. Assume that R is regular, as before, let x_1, \dots, x_n be as before, and let E be a finitely generated R -module. Let $e \in E$ be such that $\text{Ann}_R e = (0)$, i.e. $R \rightarrow E$ by $r \mapsto re$ is a monomorphism. Then Re is a direct summand of E if and only if for every integer $k \geq 0$,

$$(x_1 \cdots x_n)^k e \notin (x_1^{k+1}, \dots, x_n^{k+1})E .$$

The proof is essentially the same.

Moreover, it is not necessary to assume that R is regular. Let (R, P) be any local ring, and let $\{I_k\}$ be a sequence of irreducible P -primary ideals cofinal with the powers of P . (If R is Gorenstein and x_1, \dots, x_n is any system of parameters, we can take I_k to be (x_1^k, \dots, x_n^k) .) Let $u_k \in R$ generate the minimum nonzero ideal of R/I_k modulo I_k . (Note that a P -primary ideal I is irreducible if and only if R/I is a 0-dimensional Gorenstein ring.) Let E be a finitely generated R -module and $e \in E$ be

such that $\text{Ann}_R e = (0)$. Then Re is a direct summand of E if and only if for every integer $k \geq 0$.

$$u_k e \notin I_k E.$$

(If R is Gorenstein, x_1, \dots, x_n is a system of parameters, and $I_k = (x_1^k, \dots, x_n^k)$, we may always take $u_{k+1} = (x_1 \cdots x_n)^k u_1$). The proof of this result follows precisely the lines of the proof of Theorem 1.

Remark 3. We also note the following result. Let R be a regular local ring and x_1, \dots, x_n a regular system of parameters. Let E be a finitely generated R -module. Then E has a direct summand isomorphic to R if and only if $\text{Ann}_R E = 0$ and for every integer $k \geq 0$,

$$(x_1 \cdots x_n)^k E \not\subset (x_1^{k+1}, \dots, x_n^{k+1}) E.$$

To see why this is true let

$$E_k = (x_1^{k+1}, \dots, x_n^{k+1}) E : (x_1 \cdots x_n)^k R \subset E.$$

Then $\{E_k\}$ is an increasing sequence of submodules of E and we can choose k such that $E_k = E_{k+1} = \dots$. Then $E_k \neq E$. Also, if $T =$

$$\{e \in E : \text{for some } r \in R - \{0\}, re = 0\}$$

then T is a proper submodule of E . Hence, $E \neq E_k \cup T$, and we can choose e not in E_k or T . Then $Re \cong R$ is a direct summand of E , by the first part of Remark 2.

We now are ready to handle the “geometric case.”

THEOREM 2. Let R be a ring which contains a field and suppose that for every prime P of R , R_P is a regular local ring. Let $S \supset R$ be an R -algebra which is finitely presented as an R -module. Then R is a direct summand of S .

Hence, if R is a regular Noetherian ring which contains a field and S is a module-finite extension algebra, then R is a direct summand of S .

Proof. We note that the second statement follows from the first simply because a finitely generated module over a Noetherian ring is finitely presented.

As for the proof of the first statement, we first handle the trivial case where R contains a field of characteristic zero. By Lemma 1, we

can assume that R is local, by Lemma 3 we can assume that S is a domain, and the result follows at once from Lemma 2.

Now assume that R contains a field of characteristic $p > 0$, so that for every t the map $h_t: S \rightarrow S$ by $h_t(r) = r^{(p^t)}$ is a ring homomorphism. As above, we can assume that R is local, and from the second part of Lemma 1 we can assume that R is complete. From Lemma 3 we can also assume that S is a domain. Thus, S will also be a complete local domain (module-finite over R). Let x_1, \dots, x_n be a regular system of parameters for R . To complete the proof we need only show that there is no relation of the form

$$(\#) \quad (x_1 \cdots x_n)^k = \sum_{i=1}^n s_i x_i^{k+1},$$

where the $s_i \in S$, by Theorem 1. We shall give two different proofs that $(\#)$ is impossible. The first is more elementary but the second yields somewhat more information.

For the first proof, we note that since S is a domain, it is torsion-free over R , and consequently can be embedded in a free R -module. Hence, there is an R -homomorphism $g: S \rightarrow R$ such that $a = g(1) \neq 0$. It follows that we can choose t so large that $a \notin P^{p^t}$. We can raise both sides of $(\#)$ to the p^t power to obtain:

$$(*) \quad (x_1 \cdots x_n)^{kp^t} \cdot 1 = \sum_{i=1}^n s_i^{p^t} x_i^{k p^t + p^t}$$

Apply g to both sides:

$$(x_1 \cdots x_n)^{k'} a = \sum_{i=1}^n r_i x_i^{k' + p^t}$$

where $k' = kp^t$ and $r_i = g(s_i^{p^t}) \in R$. Hence, in the regular ring R ,

$$a \in (x_1^{k' + p^t}, \dots, x_n^{k' + p^t}) : (x_1 \cdots x_n)^{k'}$$

and since x_1, \dots, x_n is an R -sequence it follows that this colon of ideals can be computed precisely as though x_1, \dots, x_n were indeterminates over Z (see [3] or [9]), so that

$$a \in (x_1^{p^t}, \dots, x_n^{p^t}) \subset P^{p^t},$$

a contradiction. This completes the proof of Theorem 2.

Before giving the second proof of the impossibility of $(\#)$, we discuss

briefly the problem of proving Theorem 2 without the hypothesis that R contains a field. We reduce at once to the case where R is a complete local ring and S is a domain module-finite over R , and all we need to show is that $(\#)$ cannot hold. Since x_1, \dots, x_n is a system of parameters of S (S is local because R is complete) all we need is the following:

CONJECTURE 1. *If S is any local ring and x_1, \dots, x_n is a system of parameters for X , then for every integer $k \geq 0$,*

$$(x_1 \cdots x_n)^k \notin (x_1^{k+1}, \dots, x_n^{k+1}).$$

Evidently, if we give a proof of this conjecture when S contains a field of characteristic $p > 0$, then we have a new way of completing the last part of the proof of Theorem 2.

Note that in Conjecture 1, if S is a counterexample, so is \hat{S} , its completion, and if P is a prime of \hat{S} of coheight $\dim \hat{S} = \dim S$, then \hat{S}/P is a complete local domain which is a counterexample.

We now give a proof of Conjecture 1 when S contains a field of characteristic $p > 0$ which differs from the earlier argument in the proof of Theorem 2. It is based on:

PROPOSITION 2. *If y_1, \dots, y_n is a system of parameters for the local ring (S, P) and N is any sufficiently large positive integer, then the system of parameters $x_1 = y_1^N, \dots, x_n = y_n^N$ satisfies the conclusion of Conjecture 1.*

Proof. We can assume that S is complete. Then $n = \dim S$, and $H_p^n(S) \neq 0$, where H_p^i denotes the i th local cohomology module with respect to the maximal ideal P of S . (See Part 4 of Proposition 6.4 of [4].) By Theorem 2.3 of [4], $H_p^i(S)$ can be interpreted as a direct limit of Koszul homology modules, and in the special case at hand we obtain

$$H_p^n(S) = \lim_{\leftarrow k} S/(y_1^k, \dots, y_n^k)$$

where the map

$$S/(y_1^k, \dots, y_n^k) \rightarrow S/(y_1^{i+k}, \dots, y_n^{i+k})$$

is induced by multiplication by $(y_1 \cdots y_n)^i$. Since $H_p^n(S) \neq 0$, we can choose N_0 such that if $N \geq N_0$

$$\text{Im}(S/(y_1^N, \dots, y_n^N) \rightarrow H_p^n(S)) \neq 0.$$

Hence, if $N \geq N_0$ the image of 1_N in $S/(y_1^{N+i}, \dots, y_n^{N+i})$ is nonzero for every $i \geq 0$, where 1_N is the residue of 1 modulo (y_1^N, \dots, y_n^N) . But this says that for $N \geq N_0$ and every $i \geq 0$,

$$(y_1 \cdots y_n)^i \notin (y_1^{N+i}, \dots, y_n^{N+i}).$$

In particular if $i = Nk$ we have that

$$(y_1^N \cdots y_n^N)^k \notin ((y_1^N)^{k+1}, \dots, (y_n^N)^{k+1})$$

for all k and $N \geq N_0$, which is just what we wanted to prove.

Thus, the nonvanishing of local cohomology in dimension n is relevant to our problem.

We now give our second method for completing the proof of Theorem 2. If S contains a field of characteristic p and y_1, \dots, y_n is a system of parameters such that

$$(y_1 \cdots y_n)^k = \sum_{i=1}^n s_i y_i^{k+1}$$

choose N of the form p^t as in Proposition 2. Raising both sides to the N th power gives:

$$(y_1^N \cdots y_n^N)^k = \sum_{i=1}^n s_i^N (y_i^N)^{k+1},$$

contradicting Proposition 2.

We have shown that Conjecture 1 holds if S contains a field of characteristic $p > 0$. We now prove it if S contains any field, by working backwards from Theorem 2.

PROPOSITION 3. *If S is a local ring which contains a field and x_1, \dots, x_n is a system of parameters for S , then for every integer $k \geq 0$,*

$$(x_1 \cdots x_n)^k \notin (x_1^{k+1}, \dots, x_n^{k+1}).$$

Proof. We can assume without loss of generality that S is a complete equicharacteristic local domain, so that S is module-finite over its subring $R = K[[x_1, \dots, x_n]]$, where K is a coefficient field for S . Then $(x_1^{k+1}, \dots, x_n^{k+1})R$ is contracted from S , by Theorem 2 and Proposition 1, and since $(x_1 \cdots x_n)^k \notin (x_1^{k+1}, \dots, x_n^{k+1})R$, we also have that $(x_1 \cdots x_n)^k \notin (x_1^{k+1}, \dots, x_n^{k+1})S$, as required.

We next give a discussion of the relevance of Cohen-Macaulay modules. The following proposition is the key:

PROPOSITION 4. Let S be a local ring and y_1, \dots, y_n a system of parameters for S . Suppose that there is a (possibly non-Noetherian) module E over S such that

- 1) $(y_1, \dots, y_n)E \neq E$
- 2) The first Koszul homology module

$$H_1(E; y_1, \dots, y_n) = 0$$

((2) is satisfied if y_1, \dots, y_n is an E -sequence.) Then for every integer $k \geq 1$,

$$(y_1 \cdots y_n)^k \notin (y_1^{k+1}, \dots, y_n^{k+1})$$

Proof. Let Z be the integers, let x_1, \dots, x_n be indeterminates over Z , let $B = Z[x_1, \dots, x_n]$ and make S into a B -algebra by mapping x_i to y_i , $1 \leq i \leq n$. We can think of $Z = B/(x_1, \dots, x_n)$ as a B -module, and then

$$H_1(E; y_1, \dots, y_n) \cong \text{Tor}_1^B(Z, E).$$

Let $I = (x_1^k, \dots, x_n^k) \subset B$ and let $J = (x_1^k, \dots, x_n^k, (x_1 \cdots x_n)^{k+1}) \subset B$. It is easy to see that $J/I \cong Z$.

Let I_0 be any ideal of B generated by monomials in x_1, \dots, x_n which contains a power of each x_i , $1 \leq i \leq n$. We shall show that B/I_0 has a filtration in which all the factors are copies of Z . In fact, it is easy to see that if $I_0 \neq B$, there is a monomial $u \notin I_0$ such that $x_1 u, \dots, x_n u \in I_0$, that $(I_0 + uB)/I_0 \cong Z$, and the result follows from Noetherian induction on I_0 , since $I_0 + B$ is a larger ideal of the same form.

Since 2) yields that $\text{Tor}_1^B(Z, E) = 0$, we see that for I_0 as above, $\text{Tor}_1^B(B/I_0, E) = 0$. Hence, since $0 \rightarrow I_0 \rightarrow B \rightarrow B/I_0 \rightarrow 0$ is exact, so is

$$0 \rightarrow I_0 \otimes_B E \rightarrow B \otimes_B E \rightarrow (B/I_0) \otimes_B E \rightarrow 0,$$

and hence $I_0 \otimes_B E \cong I_0 E$. Thus, $I \otimes_B E \cong IE$ and $J \otimes_B E \cong JE$. Now we have the exact sequence:

$$0 \rightarrow I \rightarrow J \rightarrow J/I \rightarrow 0,$$

and $J/I \cong Z$. Applying $\otimes_B E$ and recalling $\text{Tor}_1^B(Z, E) = 0$, we have

$$0 \rightarrow IE \rightarrow JE \rightarrow Z \otimes E \rightarrow 0$$

is exact, and $Z \otimes_B E \cong E/(y_1, \dots, y_n)E \neq 0$, by 1), so that $IE \subsetneq JE$. But then $IS \subsetneq JS$, and so $(y_1 \cdots y_n)^k \notin (y_1^{k+1}, \dots, y_n^{k+1})$.

THEOREM 3. *Let R be a ring such that R_P is a regular local ring for each maximal ideal P of R , let S be an extension ring such that S is finitely presented as an R -module, and suppose that for each maximal ideal M of S and each system of parameters y_1, \dots, y_n of M , there is an S -module E such that*

- 1) $(y_1, \dots, y_n)E \neq E$
- 2) $H_1(E; y_1, \dots, y_n) = 0$.

Then R is a direct summand of S .

Proof. Let P be a maximal ideal of R . It suffices to show that R_P is a direct summand of $R_P \otimes S$ for each such P . Thus, we can reduce to the case where R is local. Let y_1, \dots, y_n be a regular system of parameters for R . Since S is module-finite over R , there is a maximal ideal M of S lying over P such that $\dim S_M = \dim R_P$, and it follows that y_1, \dots, y_n is a system of parameters for S_M . Hence, by Proposition 4, y_1, \dots, y_n have no relation like (#) in S_M , and hence no such relation in S . The result now follows from Theorem 1.

Remark 4. Suppose that for each maximal ideal M of S , the completion of S_M possesses a Cohen-Macaulay module whose dimension is equal to that of S_M . Then this module certainly satisfies the requirements of Theorem 3. Moreover, it is not known whether every complete local ring possesses a maximal Cohen-Macaulay module (this is Conjecture E of [6]). Thus, Conjecture E implies Conjecture 1 here, and also implies that the restriction that R contain a field can be removed from the hypothesis of Theorem 2.

Remark 5. Conjecture 1 is equivalent to a superficially stronger statement, Conjecture 1° below. If $(a) = (a_1, \dots, a_n)$ is an n -tuple of nonnegative integers and $x = (x_1, \dots, x_n)$ is an n -tuple of elements of a ring S , let $x^{(a)} = x_1^{a_1} \dots x_n^{a_n}$. Write $(a) \geq (b)$ if $a_i \geq b_i, 1 \leq i \leq n$. Then:

CONJECTURE 1°. *If S is a local ring and x_1, \dots, x_n is a system of parameters, then $x^{(a)} \in (x^{(b)} : 1 \leq t \leq m)S$ if and only if for some $t, (a) \geq (b_t)$.*

To see that Conjecture 1° follows from Conjecture 1, suppose 1° fails and

$$(*) \quad x^{(a)} = \sum_{t=1}^n s_t x^{(b_t)}, s_t \in S.$$

Choose $(a') \geq (a)$ with all entries of (a') equal, say $(a') = (k, \dots, k)$. If we multiply both sides of $(*)$ by $x^{(a')-(a)}$ we obtain a contradictory example in which the left hand side is $(x_1 \cdots x_n)^k$. (For each t , $(a) \not\geq (b_t)$ implies that $(a') \not\geq (b_t) + (a') - (a)$). Hence, we can assume without loss of generality that $(a) = (k, \dots, k)$. Now $(k, \dots, k) \not\geq (b_t)$ implies that some entry of b_t is at least $k + 1$, and hence $(x^{(b_t)} : 1 \leq t \leq m)S \subset (x_1^{k+1}, \dots, x_n^{k+1})S$, and $(x_1 \cdots x_n)^k \in (x_1^{k+1}, \dots, x_n^{k+1})S$, contradicting Conjecture 1.

Conjecture 1° is known if the monomials involved all have the same degree [10], Theorem 21, p. 292. This result is used to prove the analytic independence of a system of parameters, and is really a somewhat stronger and more general statement. Hence, Conjecture 1 (or 1°) is quite a bit stronger than the analytic independence of a system of parameters.

Remark 6. Theorem 21 of [10] has a module-theoretic extension which is quite easy to prove. It is worth noting this extension, and also that the corresponding extension of Conjecture 1° fails.

The module-theoretic extension of Theorem 21 of [10] is as follows: Let $x^{(a)}, x^{(b_1)}, \dots, x^{(b_m)}$ be distinct monomials of the same degree d in a system of parameters x_1, \dots, x_n of a local ring S , and let E be an S -module of dimension n , so that x_1, \dots, x_n is a system of parameters for E . Then

$$x^{(a)}E \not\subset (x^{(b_1)}, \dots, x^{(b_m)})E.$$

To prove this, let N_e be the number of monomials of degree e in x_1, \dots, x_n . If $e \geq d$, then $(x_1, \dots, x_n)^e E$ is $\subset (S_e)E$, where S_e is the set of monomials of degree e which are not multiples of $x^{(a)}$, so that S_e has $N_e - N_{e-d}$ elements. Hence, $E_e = (x_1, \dots, x_n)^e E / (x_1, \dots, x_n)^{e+1} E$ has a filtration whose factors are $N_e - N_{e-d}$ homomorphic images of $E / (x_1, \dots, x_n)E$, and $\ell(E_e) \leq b(N_e - N_{e-d})$, where $b = \ell(E / (x_1, \dots, x_n)E)$. It follows at once that the degree of the Hilbert polynomial $f(e)$ of E (which agrees with $\ell(E / (x_1, \dots, x_n)^e E)$ for large e) is equal to the degree of $N_e, n - 1$, when it should have degree n .

To see that this does not generalize to the context of Conjecture 1°, note that if $\dim S \geq 2$, x_1, x_2 is part of a system of parameters, and E is the ideal $(x_1^2, x_2^2)S$, then

$$x_1^2 x_2^2 E \subset (x_1^3, x_2^3) E.$$

Remark 7. Conjecture E is known for local rings of dimension ≤ 2 . Hence, Conjectures 1 and 1° hold in dimension ≤ 2 , and the hypothesis that R contains a field can be dropped in Theorem 2 if $\dim R \leq 2$.

2. Ideally integrally closed domains. Call a domain D *ideally integrally closed* (or *IIC*) if for every integral extension ring S of D and every ideal I of D , $IS \cap D = I$. Thus, every ideal of D is contracted from every integral extension. In this section we briefly explore this notion. We begin with some rather straight forward observations:

A. *An IIC domain D is integrally closed.* (For if b/a is in the integral closure S of D , b is in $(aD)R \cap D = aD$, and b/a is in D .)

B. *To determine whether D is IIC it suffices to consider only domains S integral over D . Hence, it suffices to consider the integral closure S of D in an algebraic closure of the fraction field of D .* (For if $IS \cap D = J$, and J properly contains I , we can choose a prime P of S maximal with respect to disjointness from the multiplicative system $D - \{0\}$, and then $I(S/P) \cap D$ contains J .)

C. *It suffices to consider only module-finite domain extensions of D , since each domain integral over D is a directed union of these. Likewise, it suffices to consider only finitely generated ideals of D . Hence, Prüfer domains are IIC.*

D. *If D is IIC, then so is $T^{-1}D$ for any multiplicative system T in D .* (For if S is integral over $T^{-1}D$ and I is an ideal of $T^{-1}D$ such that $IS \cap T^{-1}D = J$ properly contains I , then if S_0 is the integral closure of D in S , it is easy to see that $((I \cap D)S_0) \cap D = J \cap D$ properly contains $I \cap D$.)

E. *If for each maximal ideal M of the domain D the domain D_M is IIC, then D is IIC.* (For suppose S is a domain integral over D and $IS \cap D = J$ properly contains I . Choose a maximal ideal M of D which contains $I:J$ in D . $((ID_M)(D - M)^{-1}S) \cap D_M$ contains JD_M which properly contains ID_M .)

F. *If D is a domain which is a directed union of IIC domains, then D is an IIC domain. In fact, it suffices that each subdomain of D*

which is finitely generated over the prime subdomain D_0 of D be contained in a subdomain of D which is IIC. (For suppose $j \notin (i_1, \dots, i_n)D$ but S is a domain integral over D such that $J = \sum_t s_t i_t$, $s_t \in S$ for each t . Clearly, we can choose a ring $D_1 \subset D$ finitely generated over D_0 which contains j, i_1, \dots, i_n and sufficiently many elements so that r_1, \dots, r_n are integral over D_1 . Then we can choose an IIC subdomain D_2 of D with $D_1 \subset D_2$. Let $R_2 = D_2[r_1, \dots, r_n]$. Then $j \notin (i_1, \dots, i_n)D_2$, but $j \in (i_1, \dots, i_n)R_2$, contradiction.)

G. If D is a domain such that every module-finite extension domain S of D has D as a direct summand (as a D -module), then D is IIC. (This is immediate from Proposition 1.)

H. If D is a domain which contains the rationals, Q , then D is IIC if and only if D is integrally closed. (This is immediate from A., G., above, and Lemma 2.)

I. If $D \subset R$, R is an IIC domain, and for every ideal I of D , $IR \cap D = I$, then D is IIC. In particular, if R is IIC and faithfully flat over D , or if R is IIC and D is a direct summand of R , then D is IIC. It follows that if the completion \hat{D} of a Noetherian local domain D is an IIC domain, then D is an IIC domain. (Let L be an algebraically closed field which contains R . If C is a domain integral over D , there is a copy of C in L , and $R[C]$ is a domain integral over R . Hence, for any ideal I of D , we have $IC \cap D \subset (IR[C]) \cap (R \cap D) = (IR[C] \cap R) \cap D = (IR) \cap D = I$.)

Let us say that a proper ideal I of a commutative ring R is *big* if it is not an intersection of properly larger ideals. Thus, I is big if and only if there exists $x \notin I$ such that x is in every ideal properly larger than I . Obviously, I is big if and only if (0) is big in R/I . By a trivial application of Zorn's lemma, every proper ideal of a commutative ring is an intersection of big ideals. Hence:

J. A domain D is IIC if and only if for every module-finite domain extension S of D and every big ideal I of D , $IS \cap D = I$.

It is easy to show that if R is Noetherian then $I \subseteq R$ is big if and only if I is primary to a maximal ideal and irreducible. Thus, (0) is big in R if and only if R is a zero-dimensional Gorenstein local ring.

K. A Gorenstein local domain (D, P) is IIC if and only if for every module-finite domain extension S of D , D is a direct summand of S . Moreover, if $\{I_k\}$ is a sequence of P -primary irreducible ideals of D cofinal with $\{P^k\}$, then D is IIC if and only if for every module-finite domain extension S of D , each I_k is contracted. (It suffices to show that if each I_k is contracted from S then D is a direct summand of S . See the second paragraph of Remark 2.)

L. Let (D, P) be a two-dimensional Gorenstein local domain and let x_1, x_2 be a system of parameters. If (x_1, x_2) is contracted from every module-finite extension domain S of D , then D is IIC. (Let u generate $(x_1, x_2):P$ modulo (x_1, x_2) . It suffices to show that we cannot have $x_1^k x_2^k u = s_1 x_1^{k+1} + s_2 x_2^{k+1}$. Let S' be the integral closure of S . Then S' is Noetherian [8], Theorem (33.12), p. 120, and semilocal, and x_2 is not a zerodivisor modulo $x_1 S'$, for S' is integrally closed and $\dim S' = 2$, so that S' is Cohen-Macaulay. Since x_1, x_2 is an S' -sequence, we must have that x_1^k divides s_2 in S' , and that x_2^k divides s_1 in S' . Hence, $x_1^k x_2^k u = x_1^{k+1} s'_1 + x_2^{k+1} s'_2$, $s'_1, s'_2 \in S'$, and $u = x_1 s'_1 + x_2 s'_2$, so that $(x_1, x_2)D$ is not contracted from $D[s'_1, s'_2]$, a contradiction.)

This completes our list of observations about IIC domains. A somewhat more interesting result (although the real work was done in the proof of Theorem 2) is:

PROPOSITION 5. Let D be a domain such that for each maximal ideal M of D , D_M is a regular local ring, and suppose that D contains a field. Then D is IIC.

Proof. This follows at once from observations E. and G. and Theorem 2.

The following proposition shows that there are interesting non-regular examples of IIC domains in characteristic p .

PROPOSITION 6. Let k be any field and let x_1, \dots, x_n be indeterminates over k . Let \mathfrak{m} be a set of monomials (expressions of the form $x_1^{i_1} \cdots x_n^{i_n}$) in x_1, \dots, x_n such that the subring $D = k[\mathfrak{m}] \subset k[x_1, \dots, x_n]$ is integrally closed (in the fraction field of D). Then D is IIC.

Proof. We use the results of [5]. By Proposition 1 of [5], the fact that $k[\mathfrak{m}]$ is normal implies that \mathfrak{m} generates a normal semigroup

\mathfrak{S} of monomials and hence that \mathfrak{S} is isomorphic to a full semigroup \mathfrak{T} of monomials (possibly in different indeterminates). But then $k[\mathfrak{T}]$ is a direct summand of a polynomial ring, by Lemma 1 of [5], and it follows from Theorem 2 here and observation I. that $k[m] \cong k[\mathfrak{T}]$ is *IIC*.

The rest of this paper is devoted primarily to a discussion of open questions and examples.

Example 1, revisited. The ring D of Example 1 is an integrally closed Gorenstein ring which is not *IIC*. D has, moreover, a module-finite extension which is *IIC*.

QUESTION 1. Is there some nice class of domains which have minimal or minimum *IIC* extensions (analogues of integral closures)?

In connection with this we note that the intersection of a decreasing sequence of *IIC* domains need not be *IIC*. In Example 1, D is countable, and hence has only countably many height 1 primes: say P_1, P_2, P_3, \dots is an enumeration of them. Let $T_i = D - (\cup_{t=1}^i P_t)$ and let $D_i = T_i^{-1}D$. Then D_i is an integrally closed semilocal domain of Krull dimension 1, and hence D_i is *IIC* by Remark 7. But $D = \cap_i D_i$.

However, I do not know the answer to

QUESTION 2. If D_1, D_2 are *IIC* subdomains of D , is $D_1 \cap D_2$ *IIC*?
Some other obvious lines of inquiry are:

QUESTION 3. If D is *IIC*, is $D[x]$ *IIC*? Is $D[[x]]$ *IIC*? Here, x is an indeterminate.

QUESTION 4. If D is finitely generated over a field, is

$$\{P \in \text{Spec } D : D_P \text{ is } IIC\}$$

Zariski open in $\text{Spec } D$?

QUESTION 5. If D_1 and D_2 are *IIC* domains finitely generated over an algebraically closed field k , is $D_1 \otimes_k D_2$ *IIC*? (If $D_2 = k[x]$, Question 5 becomes a special case of Question 3.)

EXAMPLE 2. An interesting example of *UFD* which is not *IIC* comes out of invariant theory. We first need:

PROPOSITION 7. If D is local and *IIC* and R is a module-finite equidimensional extension of D which is Cohen-Macaulay, then D is Cohen-Macaulay.

Proof. Let x_1, \dots, x_n be a system of parameters for D . It follows easily that x_1, \dots, x_n is an R -sequence in R . But then, since D is *IIC*, x_1, \dots, x_n is an R -sequence in D , for if $dx_{k+1} \in (x_1, \dots, x_k)D$, then $d \in (x_1, \dots, x_k)R \cap D = (x_1, \dots, x_k)D$.

Now let K be a field of characteristic 2 and let $R_1 = K[x_1, x_2, x_3, x_4]$. Let g be the K -automorphism of R_1 such that $g(x_i) = x_i$ while $g(x_i) = x_i + x_{i+1}$, $i = 1, 2, 3$. Let G be the cyclic group of order 4 generated by g , so that we have an action of G on R_1 . The ring of invariants D_1 is a graded K -algebra which is known [1] to be a non-Cohen-Macaulay *UFD* of dimension 4. Let D, R be the localizations of D_1, R_1 at their respective irrelevant maximal ideals. Since R is module-finite over D and Cohen-Macaulay (in fact, regular) while D is not Cohen-Macaulay, it follows from Proposition 7 that D is not *IIC*.

On the other hand we note:

PROPOSITION 8. *If G is a finite group acting on an *IIC* domain D and the order d of G is invertible in D , then the ring of invariants*

$$D^G = \{r \in D : g(r) = r \text{ for all } g \in G\}$$

*is *IIC*.*

Proof. The map $h : D \rightarrow D^G$ by

$$h(r) = (1/d) \sum_{g \in G} g(r)$$

is a D^G -module retraction of D onto D^G .

EXAMPLE 3. Proposition 6 provides a number of interesting examples of non-regular *IIC* domains in characteristic $p > 0$. One example is the usual homogeneous coordinate ring for a multiprojective space

$$\mathbf{P}^{t_1} \times \dots \times \mathbf{P}^{t_n}$$

namely, the Segre product (see [2] and [7], §13) of n polynomial rings in $t_1 + 1, \dots, t_n + 1$ variables, respectively. A particularly simple case is $\mathbf{P}^1 \times \mathbf{P}^1$ where the homogeneous coordinate ring is

$$\begin{aligned} D &= k[x_1y_1, x_1y_2, x_2y_1, x_2y_2] \\ &\cong k[p, q, r, s]/(ps - qr) \end{aligned}$$

Another example is the homogeneous coordinate ring for \mathbf{P}^{n-1} spanned

by all forms whose degree is a multiple of d in $k[x_1, \dots, x_n]$. The simplest case is

$$D = k[x^2, xy, y^2] \cong k[p, q, r]/(pr - q^2)$$

Finally, we note that some of the examples above have an invariant-theoretic flavor, and we raise the following question:

QUESTION 6. If k is an algebraically closed field of characteristic p and G is a connected linear algebraic group acting rationally on an n -dimensional vector space V , so that the symmetric algebra $R = S(V) \cong k[x_1, \dots, x_n]$, then is the ring of invariants R^G of the induced action of G on R *IIC*? More generally, if R is an *IIC* domain and D is integrally closed in R , is D *IIC*?

REFERENCES

- [1] M^m M. Bertin, Anneaux d'invariants d'anneaux de polynomes, en caracterisque p . Comptes Rendus. Ser. A. **264**:2 (1967), 653–656.
- [2] W. L. Chow, On unmixedness theorem, Amer. J. Math. **86** (1964), 799–822.
- [3] J. A. Eagon and M. Hochster, R-sequences and indeterminates, preprint.
- [4] A. Grothendieck (notes by R. Hartshorne), Local cohomology, Springer-Verlag Lecture Notes in Mathematics No. 41, 1967.
- [5] M. Hochster, Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes, Annals of Math., **96** (1972), 318–337.
- [6] M. Hochster, Cohen-Macaulay modules, Proc. Kansas Commutative Algebra Conference, Springer-Verlag, Lecture Notes in Mathematics No. 311, 1973.
- [7] M. Hochster and J. A. Eagon, Cohen-Macaulay rings, invariant theory, and the generic perfection of determinantal loci, Amer. J. Math. **93** (1971), 1020–1058.
- [8] M. Nagata, Local rings, Interscience, New York, 1962.
- [9] D. Taylor, Ideals generated by monomials in an R-sequence, Thesis, University of Chicago, 1966.
- [10] O. Zariski and P. Samuel, Commutative algebra, Vol. II, Van Nostrand, Princeton, 1960.

University of Minnesota