

Bridging Galaxy Dynamics and Baryon Efficiency of 40 EDGE-CALIFA galaxies

Veselina Kalinova¹, Dario Colombo¹ and Erik Rosolowsky¹

¹Department of Physics 4-181 CCIS, University of Alberta, Edmonton AB T6G2E1, Canada
email: veselina@ualberta.ca

Abstract. We apply the Jeans Axisymmetric Multi-Gaussian Expansion method to the stellar kinematic maps of 40 Sa–Sd EDGE-CALIFA galaxies and derive their circular velocity curves (CVCs). The CVCs are classified using the Dynamical Classification method developed in Kalinova *et al.* (2015). We also calculate the observational baryon efficiency, OBE, where $M_*/M_b = M_*/(M_* + M_{HI} + M_{H_2})$ of the galaxies using their stellar mass, total neutral hydrogen mass and total molecular gas from CO luminosities. Slow-rising, Flat and Round-peaked CVC types correspond to specific OBEs, stellar and dark matter (DM) halo mass values, while the Sharp-peaked CVCs span in the whole DM halo mass range of $10^{11} - 10^{14} M_\odot$.

Keywords. Galaxies: kinematics and dynamics, galaxies: spiral, surveys

Introduction. Galaxy formation and evolution are regulated by the interaction between three fundamental components: stars, gas, and dark matter. To understand the link between these components, modern simulations have used the “abundance matching” where empirical relationships between stellar mass, star formation (SF) efficiencies, and the DM halo mass (M_h) are calibrated using populations of galaxies at different redshifts (Moster *et al.* 2013). This model proposes two domains based on SF feedback (low-mass) vs. Active Galactic Nuclei feedback (high-mass) tied to specific values of M_h and baryon efficiency. The parameters in the relationship imply globally important physics as fundamental as the Kennicutt-Schmidt law (Kennicutt *et al.* 1998), and the origin of the cold gas phase with respect to the stellar disc (Davis *et al.* 2011).

Data and Method. To observationally test the theoretical prediction of Moster *et al.* (2013) we focused on 40 Sa–Sd galaxies from the optical integral-field spectroscopic survey CALIFA (Sánchez *et al.* 2012) and the CARMA-CO survey EDGE† (Bolatto *et al.*, in prep). Dynamical models and CVCs of the galaxies are constructed using the Multi-Gaussian Expansion method (MGE; Emsellem *et al.* 1994) on SDSS images and Jeans Anisotropic MGE model (JAM; Cappellari 2008) on CALIFA stellar kinematics (Falcón-Barroso *et al.*, in prep). Further, we classify the CVCs of the 40 galaxies by shape and amplitude using the Dynamical classification method, described in Kalinova *et al.* 2015. The principal components of our sample are obtained through the eigenvectors of the prototype curves in Figure 7 of their paper, i.e. a given curve is associated to a specific dynamical class (Slow-rising, Flat, Sharp-peaked, Round-peaked) based on the minimum Euclidean distance to the k-means centroids that define the prototype curves in the five-dimensional principal components clustering space. The final dynamical classification of our 40 galaxies is shown in Fig. 1a. To understand how these four dynamical classes are related to the OBE and M_h of the galaxies we use also measurements of the total stellar mass (M_*) adopted from González-Delgado *et al.* (2015), while the M_{HI} are taken from Springob *et al.* (2005). DM halo mass estimations are extrapolated from the $M_* - M_h$ relation of Moster *et al.* (2013) using our adopted stellar masses (see Fig. 1b, top). The CO luminosity of the molecular gas is derived from the CARMA CO(1-0) EDGE integrated intensity maps (Bolatto *et al.*, in prep), and then converted in total M_{H_2} using the factor $\alpha_{CO} = 4.4 M_\odot (\text{pc}^2 \text{ K km s}^{-1})^{-1}$.

Results. Our analysis shows that Slow-rising CVCs are mainly presented by low M_h ($10^{11} - 10^{12} M_\odot$) and low observational baryon efficiencies, while Round-peaked CVCs correspond to galaxies with high OBE and high implied M_h ($10^{12} - 10^{14} M_\odot$). Flat CVCs cluster around intermediate DM halos, i.e., M_h ($10^{12} - 10^{13} M_\odot$) with approximately constant and high OBEs. Sharp-peaked CVCs, instead, span the entire OBE range and DM halo mass range ($10^{11} -$

† <http://www.astro.umd.edu/~bolatto/EDGE/>




Figure 1. (a) Dynamical Classification of 40 (Sa–Sd) EDGE-CALIFA galaxies. (b) *Top*: Extrapolated DM Halo masses from the $M_* - M_h$ relation of Moster *et al.* (2013) using the stellar mass of the galaxies from González-Delgado *et al.* (2015). *Bottom*: Observational Baryon Efficiencies (M_*/M_b) of the galaxies throughout their dynamical classes, shown in different color.

$10^{14} M_\odot$, see Fig 1b). The dynamics of the 40 galaxies, and in particular, the shape and the amplitude of their CVCs might be linked to the cooling of the hot gas and formation of the stellar disc in the different DM halos mass ranges. This represents the key to unveil the connection between internal and external processes in the galaxies. Better conclusions will be drawn via DM halo masses measured by high resolution HI rotation curves and modelling of the hot gas content.

References

- Cappellari, M. 2008, *MNRAS*, 390, 71
 Davis, T. A., Alatalo, K., Sarzi, M., et al. 2011, *MNRAS*, 417, 882
 Emsellem, E., Monnet, G., & Bacon, R. 1994, *A&A*, 285, 723
 González Delgado, R. M., García-Benito, R., Pérez, E., et al. 2015, *A&A*, 581, A103
 Kalinova, V., Colombo, D., Rosolowsky, E., et al. 2015, arXiv:1509.03352
 Kennicutt, R. C., Jr. 1998, *ApJ*, 498, 541
 Moster, B. P., Naab, T., & White, S. D. M. 2013, *MNRAS*, 428, 3121
 Sánchez, S. F., Kennicutt, R. C., Gil de Paz, A., et al. 2012, *A&A*, 538, A8
 Springob, C. M., Haynes, M. P., Giovanelli, R., & Kent, B. R. 2005, *ApJS*, 160, 149