

## High Resolution CO Images of Seyfert Galaxies

Margaret Meixner and Melvyn Wright  
*Astronomy Dept., Univ. of California, Berkeley, CA 94720 USA*

Rich Puchalsky and Leo Blitz  
*Astronomy Program, Univ. of Maryland, College Park, MD 20742 USA*

Tim Heckmen  
*Dept. of Physics and Astronomy, Johns Hopkins Univ., Baltimore, MD 21218 USA*

### Summary

The Berkeley-Illinois-Maryland millimeter array has been used to image the CO (J=1-0) emission of three Seyfert galaxies, NGC 3227, NGC 7469 and NGC 5033, selected for their strong single-dish CO detections. Extensive u-v coverage was obtained for all three galaxies resulting in 2"-3" resolution. The CO emission in NGC 3227 and NGC 7469 appears as compact structures centered on the active nuclei, containing substantial fractions ( $\sim 80\%$  and  $\sim 50\%$ ) of the single-dish flux. In NGC 3227, 10% of the CO flux detected by the interferometer is contained within the ionized narrow line region. The unresolved molecular gas concentrations in the nucleus of NGC 3227 imply a CO mass of  $6.5 \times 10^7 M_{\odot}$  concentrated within a diameter  $\lesssim 50$  pc. The CO emission in NGC 5033 is not detected at this resolution, implying a CO structure size of 20"-60" (1.2-3.6 kpc). Continuum emission at 2.7mm is not detected in any of the three galaxies. In the center of NGC 7469, the H<sub>2</sub> mass is comparable to the dynamical mass. Kinematic studies of the detected gas reveal a rotational motion of the gas in NGC 3227 and NGC 7469, allowing identification of the gas in NGC 7469 with a nuclear starburst. These data are consistent with the idea that interactions between galaxies cause gas to concentrate in their nuclei thereby feeding starburst and Seyfert activity.

**Fig.** NGC 3227: CO contours overlain on H $\alpha$  emission. Line marks major axis of galaxy.

