

implicated in depressive symptomatology, parasuicide and suicide; a similar association with depression has been found with vigabatrin. In contrast, carbamazepine (CBZ) has been shown to ameliorate depression and reduce anxiety in PWE. Valproate and lamotrigine may also be associated with less depressed mood. It is also important to note that both parasuicide and suicide are significantly more common in PWE than in the general population. Treatment of depression in PWE may be problematic as the majority of antidepressants lower the seizure threshold. Thus, rationalising AEDs with the reduction of polypharmacy, and prescription, where possible, of monotherapy (with special reference to CBZ) may well be the first step. After that, the judicious prescription of "safer" antidepressants (with respect to both lowering the seizure threshold and safety in overdose) may be indicated.

THERAPEUTIC ASPECTS OF DEPRESSION AND EPILEPSY: NEW VS. OLD ANTIDEPRESSANT DRUGS

R. Torta, R. Borio, R.A. Cicolin, F. Monaco, L. Ravizza. Dep. of Neurosciences, University of Turin v. Cherasco 11, 10126 Torino, Italy

Depressive symptoms can affect clinical prognosis of epilepsy and increase risk of suicide: therefore antidepressant drugs (AD) have to be frequently used in depressed epileptic patients. Nevertheless data obtained in vitro and in animals indicate that most tricyclic antidepressants (TCA) exert anticonvulsant, proconvulsant or convulsant effects in relationship with the dose used: higher doses are related with more common convulsant effect. Particularly some old AD are characterized by an intrinsic proconvulsant activity (i.e. amoxapine, maprotiline, etc.), others show a lower seizure risk (nomifensine, trazodone, viloxazine, etc.). At last, in some cases, pro- or anticonvulsant properties seem to be dose related. A significant proportion of drug-related seizures occurs in patients with an identifiable predisposition such as previous history of epilepsy, an abnormal EEG, the presence of cerebropathy, an alcohol abuse, concomitant medications.

The effects of Selective Serotonin Reuptake Inhibitors (SSRI) have to be well defined, even if experimental studies show interesting results about an antiepileptic activity for some of these compounds. Particularly fluoxetine exerts anticonvulsant actions against maximal electroshock convulsions in genetically seizure-prone rodents and enhances the anticonvulsant effects of phenytoin and carbamazepine. A discrepancy is clinically reported in the literature between studies that observe an anticonvulsant action of fluoxetine in epileptic patients and others reporting a lack of potentiation of these antidepressant in drug-resistant epilepsy.

Otherwise some Authors report isolated proconvulsant effects shown by zimelidine, fluvoxamine and fluoxetine or toxic reactions following the combined administration of fluoxetine and phenytoin.

In our depressed epileptic patients, treated with fluoxetine, we observed a statistically significant improvement of depressive symptoms and anxiety, without changes of EEG, seizure frequency and anticonvulsant plasma levels.

FORCED NORMALISATION AND ITS RELEVANCE FOR NEUROPSYCHIATRY

Michael R. Trimble. Institute of Neurology, University Department of Clinical Neurology, Queen Square, London WC1N 3BG

The relationship between epilepsy and psychiatry has several interesting interfaces. One of these is the clinical observation that suddenly stopping seizures in patients who have habitual seizures may be associated with the onset of an acute behaviour disorder. Landolt, in the 1960's identified the phenomenon of forced normal-

isation. This was essentially an EEG concept, in which suppression of seizures, with "normalisation" of the EEG was associated with the presentation of a schizophrenia like psychosis. On resolution of the psychosis with a seizure, the EEG abnormalities returned.

The observations on forced normalisation have been confirmed by many anecdotes but the phenomenon has been poorly studied. In part this is because of the necessity to examine EEG data while patients are psychiatrically disturbed. However, the theoretical interest of these observations in relationship to the opposite, namely the resolution of the psychosis in psychiatry following the administration of ECT is obviously of relevance for psychiatry.

In this presentation the background to forced normalisation, and the clinical observations related to it will be presented. This will be followed by some personal examples and then the theoretical underpinnings of this condition will be examined. The relationship between these phenomena in epilepsy and ECT will then be explored.

S53. Recent empirical data informing prevention strategies

Chairmen: R Jenkins, D de Leo

PREVENTION OF DEPRESSION AND SUICIDE IN PRIMARY CARE

W. Rutz, Jan Wålinder, Lars von Knorring, Hans Pihlgren, Zoltan Rihmer. Department of Psychiatry, Box 1343, S-621 24 Visby, Sweden

Depression is an endemic disease with high morbidity and a high mortality in depression related suicide. Suicidality, especially amongst males is increasing today in East European countries. General practitioners are considered to have a most important role in the prevention and treatment of depression. Matters of depression are today more openly discussed in the society. Patients refuse today less to see depression as a causative factor behind their symptoms when showing up in primary care. Taboos concerning the stigma of depression and suicidality are reduced. In spite of this still today important problems exist regarding poor diagnostic and treatment routines in primary care and the inability of depressive and suicidal males to seek for help or to be recognized.

In the years 1983–1984, the Swedish Committee for Prevention and Treatment of Depression (PTD) offered an educational program to all general practitioners (GP:s) on the Swedish island of Gotland. During the 80:ies this education has been shown to lead to a significant decrease in inpatient care, morbidity, suicide, mortality and costs caused by depressive illness on the island. Unspecific anxiolytic and sedative medication decreased and specific antidepressive medication increased. Thus, evidence was found that a quality improving educational program in primary care with focus on depression and suicide was effective. A shortcoming was, however, that the number of male suicides was almost unaffected by the education and the GP:s improved ability to diagnose and treat depressions.

The role of the Gotland study as a possible model for preventing depression and depression related suicide is described. Strategies to avoid failures concerning its efficacy on the prevention of male suicides are discussed.