

GROUPS ACTING ON TREES WITH PRESCRIBED LOCAL ACTION

STEPHAN TORNIER 

(Received 12 October 2021; accepted 21 May 2022; first published online 12 September 2022)

Communicated by Michael Giudici

Abstract

We extend the Burger–Mozes theory of closed, nondiscrete, locally quasiprimitive automorphism groups of locally finite, connected graphs to the semiprimitive case, and develop a generalization of Burger–Mozes universal groups acting on the regular tree T_d of degree $d \in \mathbb{N}_{\geq 3}$. Three applications are given. First, we characterize the automorphism types that the quasicentre of a nondiscrete subgroup of $\text{Aut}(T_d)$ may feature in terms of the group’s local action. In doing so, we explicitly construct closed, nondiscrete, compactly generated subgroups of $\text{Aut}(T_d)$ with nontrivial quasicentre, and see that the Burger–Mozes theory does not extend further to the transitive case. We then characterize the (P_k) -closures of locally transitive subgroups of $\text{Aut}(T_d)$ containing an involutive inversion, and thereby partially answer two questions by Banks *et al.* [‘Simple groups of automorphisms of trees determined by their actions on finite subtrees’, *J. Group Theory* **18**(2) (2015), 235–261]. Finally, we offer a new view on the Weiss conjecture.

2020 Mathematics subject classification: primary 20E08; secondary 22D05.

Keywords and phrases: groups acting on trees, totally disconnected locally compact group, semiprimitive, local action, quasicentre, independence property, Weiss conjecture.

1. Introduction

In the structure theory of locally compact (l.c.) groups, totally disconnected (t.d.) ones are in focus because any locally compact group G is an extension of its connected component G_0 by the totally disconnected quotient G/G_0 ,

$$1 \longrightarrow G_0 \longrightarrow G \longrightarrow G/G_0 \longrightarrow 1,$$

and connected l.c. groups have been identified as inverse limits of Lie groups in seminal work by Gleason [13], Montgomery and Zippin [20] and Yamabe [35].

The author was supported by the Swiss National Science Foundation (Doc.mobility fellowship 172120) and the Australian Research Council (DP120100996, FL170100032 and DE210100180).

© The Author(s), 2022. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Every t.d.l.c. group can be viewed as a directed union of compactly generated open subgroups. Among the latter, groups acting on regular graphs and trees stand out due to the Cayley–Abels graph construction: every compactly generated t.d.l.c. group G acts vertex-transitively on a connected regular graph Γ of finite degree d with compact kernel K . In particular, the universal cover of Γ is the d -regular tree T_d and we obtain a cocompact subgroup \tilde{G} of its automorphism group $\text{Aut}(T_d)$,

$$1 \longrightarrow \pi_1(\Gamma) \longrightarrow \tilde{G} \longrightarrow G/K \longrightarrow 1,$$

as an extension of $\pi_1(\Gamma)$ by G/K ; see [19, Section 11.3] and [15] for details.

In studying the automorphism group $\text{Aut}(\Gamma)$ of a locally finite, connected graph $\Gamma = (V, E)$, we follow the notation of Serre [25]. The group $\text{Aut}(\Gamma)$ is t.d.l.c. when equipped with the permutation topology for its action on $V \cup E$; see Section 2.1. Given a subgroup $H \leq \text{Aut}(\Gamma)$ and a vertex $x \in V$, the stabilizer H_x of x in H induces a permutation group on the set $E(x) := \{e \in E \mid o(e) = x\}$ of edges issuing from x . We say that H is locally ‘X’ if for every $x \in V$ the said permutation group satisfies property ‘X’ (for example, being transitive, semiprimitive or quasiprimitive).

In [2], Burger and Mozes develop a remarkable structure theory of closed, nondiscrete, locally quasiprimitive subgroups of $\text{Aut}(\Gamma)$, which resembles the theory of semisimple Lie groups; see Theorem 2.2. In Section 3 (specifically Theorem 3.14) we show that this theory readily carries over to the semiprimitive case.

Let Ω be a set of cardinality $d \in \mathbb{N}_{\geq 3}$ and let $T_d = (V, E)$ be the d -regular tree. Burger and Mozes complement their structure theory with a particularly accessible class of subgroups of $\text{Aut}(T_d)$ with prescribed local action. Given $F \leq \text{Sym}(\Omega)$, their universal group $U(F)$ is closed in $\text{Aut}(T_d)$, vertex-transitive, compactly generated and locally permutation isomorphic to F . It is discrete if and only if F is semiregular. When F is transitive, $U(F)$ is maximal up to conjugation among vertex-transitive subgroups of $\text{Aut}(T_d)$ that are locally permutation isomorphic to F , hence *universal*.

We generalize the universal groups by prescribing the local action on balls of a given radius $k \in \mathbb{N}$, the Burger–Mozes construction corresponding to the case $k=1$. Equip T_d with a labelling, that is, a map $l : E \rightarrow \Omega$ such that for every $x \in V$ the map $l_x : E(x) \rightarrow \Omega$, $e \mapsto l(e)$ is a bijection, and $l(e) = l(\bar{e})$ for all $e \in E$. Also, fix a tree $B_{d,k}$ that is isomorphic to a ball of radius k around a vertex in the labelled tree T_d and let $l_x^k : B(x, k) \rightarrow B_{d,k}$ ($x \in V$) be the unique label-respecting isomorphism. Then

$$\sigma_k : \text{Aut}(T_d) \times V \rightarrow \text{Aut}(B_{d,k}), (g, x) \mapsto l_{gx}^k \circ g \circ (l_x^k)^{-1}$$

captures the k -local action of g at the vertex $x \in V$.

DEFINITION 1.1. Let $F \leq \text{Aut}(B_{d,k})$. Define

$$U_k(F) := \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V : \sigma_k(g, x) \in F\}.$$

While $U_k(F)$ is always closed, vertex-transitive and compactly generated, other properties of $U(F)$ need not carry over. In particular, the group $U_k(F)$ need not be locally action isomorphic to F ; we say that $F \leq \text{Aut}(B_{d,k})$ satisfies *condition (C)* if it is.

This can be viewed as an interchangeability condition on neighbouring local actions; see Section 4.4. There also is a discreteness condition (D) on $F \leq \text{Aut}(B_{d,k})$ in terms of certain stabilizers in F under which $U_k(F)$ is discrete; see Section 4.2.2. Finally, the groups $U_k(F)$ are universal in a sense akin to the above by Theorem 4.34.

For $\tilde{F} \leq \text{Aut}(B_{d,k})$, let $F := \pi\tilde{F} \leq \text{Sym}(\Omega)$ denote its projection to $\text{Aut}(B_{d,1})$, which is naturally permutation isomorphic to $\text{Sym}(\Omega)$ via the labelling of $B_{d,1}$. The following rigidity theorem is inspired by [2, Proposition 3.3.1].

THEOREM 1.2. *Let $F \leq \text{Sym}(\Omega)$ be 2-transitive and F_ω ($\omega \in \Omega$) simple nonabelian. Further, let $\tilde{F} \leq \text{Aut}(B_{d,k})$ with $\pi\tilde{F} = F$ satisfy (C). Then $U_k(\tilde{F})$ equals either*

$$U_2(\Gamma(F)), \quad U_2(\Delta(F)) \quad \text{or} \quad U_1(F).$$

Here, the groups $\Gamma(F), \Delta(F) \leq \text{Aut}(B_{d,2})$ of Section 4.4 satisfy both (C) and (D) and therefore yield discrete universal groups. Illustrating the necessity of the assumptions in Theorem 4.32, we construct further universal groups in the case where either point-stabilizers in F are not simple, F is not primitive, or F is not perfect; see, for example, $\Phi(F, N), \Phi(F, \mathcal{P}), \Pi(F, \rho, X) \leq \text{Aut}(B_{d,2})$ in Section 4.4.

In Section 5 we present three applications of the framework of universal groups. First, we study the quasicentre of subgroups of $\text{Aut}(T_d)$. The quasicentre $\text{QZ}(G)$ of a topological group G consists of those elements whose centralizer in G is open. It plays a major role in the Burger–Mozes structure theorem (Theorem 2.2): a nondiscrete, locally quasiprimitive subgroup of $\text{Aut}(T_d)$ does not feature any nontrivial quasiceutral elliptic elements. We extend this fact to the following local-to-global-type characterization of the automorphism types that the quasicentre of a nondiscrete subgroup of $\text{Aut}(T_d)$ may feature in terms of the group’s local action.

THEOREM 1.3. *Let $H \leq \text{Aut}(T_d)$ be nondiscrete. If H is locally:*

- (i) *Transitive, then $\text{QZ}(H)$ contains no inversion.*
- (ii) *Semiprimitive, then $\text{QZ}(H)$ contains no nontrivial edge-fixating element.*
- (iii) *Quasiprimitive, then $\text{QZ}(H)$ contains no nontrivial elliptic element.*
- (iv) *k -transitive, ($k \in \mathbb{N}$) then $\text{QZ}(H)$ contains no hyperbolic element of length k .*

More importantly, the proof of the above theorem suggests using groups of the form $\bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$ for appropriate local actions $F^{(k)} \leq \text{Aut}(B_{d,k})$ in order to explicitly construct nondiscrete subgroups of $\text{Aut}(T_d)$ whose quasicentres contain certain types of automorphisms. This leads to the following sharpness result.

THEOREM 1.4. *There exist a $d \in \mathbb{N}_{\geq 3}$ and a closed, nondiscrete, compactly generated subgroup of $\text{Aut}(T_d)$ that is locally:*

- (i) *Intransitive and contains a quasiceutral inversion.*
- (ii) *Transitive and contains a nontrivial quasiceutral edge-fixating element.*
- (iii) *Semiprimitive and contains a nontrivial quasiceutral elliptic element.*
- (iv) (a) *Intransitive and contains a quasiceutral hyperbolic element of length 1.*
 (b) *Quasiprimitive and contains a quasiceutral hyperbolic element of length 2.*

Part (ii) of this theorem can be strengthened to the following result which shows that the Burger–Mozes theory does not extend further to locally transitive groups.

THEOREM 1.5. *There exist $d \in \mathbb{N}_{\geq 3}$ and a closed, nondiscrete, compactly generated, locally transitive subgroup of $\text{Aut}(T_d)$ with open, hence nondiscrete, quasicentre.*

We also give an algebraic characterization of the (P_k) -closures of locally transitive subgroups of $\text{Aut}(T_d)$ which contain an involutive inversion. Thereby, we partially answer two questions by Banks *et al.* [1, page 259] who introduced the term (P_k) -closure in [1] and called it k -closure; however the term k -closure has an established meaning for permutation groups due to Wielandt, so we use (P_k) -closure here. Recall (Section 2.2.2) that the (P_k) -closure ($k \in \mathbb{N}$) of a subgroup $H \leq \text{Aut}(T_d)$ is given by

$$H^{(P_k)} = \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V(T_d) \text{ there exists } h \in H : g|_{B(x,k)} = h|_{B(x,k)}\}.$$

THEOREM 1.6. *Let $H \leq \text{Aut}(T_d)$ be locally transitive and contain an involutive inversion. Then $H^{(P_k)} = U_k(F^{(k)})$ for some labelling l of T_d and $F^{(k)} \leq \text{Aut}(B_{d,k})$.*

Combined with the independence properties (P_k) , $k \in \mathbb{N}$ (Section 2.2.2), introduced by Banks *et al.* [1] as generalizations of Tits' independence property, Theorem 5.31 entails the following characterization of universal groups.

COROLLARY 1.7. *Let $H \leq \text{Aut}(T_d)$ be closed, locally transitive and contain an involutive inversion. Then $H = U_k(F^{(k)})$ if and only if H satisfies property (P_k) .*

Banks, Elder and Willis use subgroups of $\text{Aut}(T_d)$ with pairwise distinct (P_k) -closures to construct infinitely many, pairwise nonconjugate, nondiscrete simple subgroups of $\text{Aut}(T_d)$ via Theorem 2.1 and ask whether they are also pairwise nonisomorphic as topological groups. We partially answer this question in the following theorem.

THEOREM 1.8. *Let $H \leq \text{Aut}(T_d)$ be nondiscrete, locally permutation isomorphic to $F \leq \text{Sym}(\Omega)$ and contain an involutive inversion. Suppose that F is transitive and that every nontrivial subnormal subgroup of F_ω ($\omega \in \Omega$) is transitive on $\Omega \setminus \{\omega\}$. If $H^{(P_k)} \neq H^{(P_l)}$ for some $k, l \in \mathbb{N}$ then $(H^{(P_k)})^{+k}$ and $(H^{(P_l)})^{+l}$ are nonisomorphic.*

Infinitely many families of pairwise nonisomorphic simple groups of this type, each sharing a certain transitive local action, are constructed in Example 5.37.

Finally, Section 5.3 offers a new view on the Weiss conjecture [33] which states that there are only finitely many conjugacy classes of discrete, locally primitive and vertex-transitive subgroups of $\text{Aut}(T_d)$ for a given $d \in \mathbb{N}_{\geq 3}$. This conjecture was extended by Potočnik *et al.* in [21] to semiprimitive local actions, and impressive partial results have been obtained by the same authors as well as by Giudici and Morgan [11]. We show that under the additional assumption that each group contains an involutive inversion, it suffices to show that for every semiprimitive $F \leq \text{Sym}(\Omega)$

there are only finitely many $\tilde{F} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) with $\pi\tilde{F} = F$ that satisfy conditions (C) and (D) in a minimal fashion; see Definition 5.42.

2. Preliminaries

This section gathers together preliminaries on permutation groups, graph theory and Burger–Mozes theory. References are given in each subsection.

2.1. Permutation groups. Let Ω be a set. In this section we give definitions and results concerning the group $\text{Sym}(\Omega)$ of bijections of Ω . Refer to [6, 12, 22] and [15, Section 1.2] for further details.

Let $F \leq \text{Sym}(\Omega)$. The *degree* of F is $|\Omega|$. For $\omega \in \Omega$, the *stabilizer* of ω in F is $F_\omega := \{\sigma \in F \mid \sigma\omega = \omega\}$. The subgroup of F generated by its point-stabilizers is denoted by $F^+ := \langle \{F_\omega \mid \omega \in \Omega\} \rangle$. The permutation group F is *semiregular*, or *free*, if $F_\omega = \{\text{id}\}$ for all $\omega \in \Omega$; equivalently, if F^+ is trivial. It is *transitive* if its action on Ω is transitive, and *regular* if it is both semiregular and transitive.

Let $F \leq \text{Sym}(\Omega)$ be transitive. The *rank* of F is the number $\text{rank}(F) := |F \backslash \Omega^2|$ of orbits of the diagonal action $\sigma \cdot (\omega, \omega') := (\sigma\omega, \sigma\omega')$ of F on Ω^2 . Equivalently, $\text{rank}(F) = |F_\omega \backslash \Omega|$ for all $\omega \in \Omega$. Note that the diagonal $\Delta(\Omega) := \{(\omega, \omega) \mid \omega \in \Omega\}$ is always an orbit of the diagonal action $F \curvearrowright \Omega^2$. The permutation group F is *2-transitive* if it acts transitively on $\Omega^2 \setminus \Delta(\Omega)$. In other words, $\text{rank}(F) = 2$.

We now define several classes of permutation groups lying in between the classes of transitive and 2-transitive permutation groups. Let $F \leq \text{Sym}(\Omega)$. A partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω is *preserved* by F , or *F-invariant*, if for all $\sigma \in F$ we have that $\{\sigma\Omega_i \mid i \in I\} = \{\Omega_i \mid i \in I\}$. The partitions $\Omega = \Omega$ and $\Omega = \bigsqcup_{\omega \in \Omega} \{\omega\}$ are *trivial*. A map $a : \Omega \rightarrow F$ is *constant with respect to* (w.r.t.) \mathcal{P} if $a(\omega) = a(\omega')$ whenever $\omega, \omega' \in \Omega_i$ for some $i \in I$. The permutation group F is *primitive* if it is transitive and preserves no nontrivial partition of Ω . Equivalently, F is transitive and its point-stabilizers are maximal subgroups. Given a normal subgroup N of F , the partition of Ω into N -orbits is F -invariant. Consequently, every nontrivial normal subgroup of a primitive group is transitive. The permutation group F is *quasiprimitive* if it is transitive and all its nontrivial normal subgroups are transitive. Finally, F is *semiprimitive* if it is transitive and all its normal subgroups are either transitive or semiregular. The following implications among the above properties follow from the definitions; we list examples illustrating that each implication is strict:

$$2\text{-transitive} \Rightarrow \text{primitive} \Rightarrow \text{quasiprimitive} \Rightarrow \text{semiprimitive} \Rightarrow \text{transitive} .$$

$$A_5 \curvearrowright A_5/D_5 \quad A_5 \curvearrowright A_5/C_5 \quad C_4 \triangleright C_2 \quad D_4 \triangleright C_2 \times C_2$$

Note that A_5 is simple and that $C_5 \leq D_5 \leq A_5$ is a nonmaximal subgroup of A_5 .

2.1.1. Permutation topology. Let X be a set and $H \leq \text{Sym}(X)$. The basic open sets of the *permutation topology* on H are $U_{x,y} := \{h \in H \mid \text{for all } i \in \{1, \dots, n\} : h(x_i) = y_i\}$, where $n \in \mathbb{N}$ and $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in X^n$. This turns H into a Hausdorff, t.d. group and makes the action map $H \times X \rightarrow X$ continuous for the discrete topology

on X . The group H is discrete if and only if the stabilizer in H of a finite subset of X is trivial. It is compact if and only if it is closed and all its orbits are finite. Finally, $\text{Sym}(X)$ is second-countable if and only if X is countable.

2.2. Graph theory. We first recall Serre's [25] notation and definitions in the context of graphs and trees, and then give generalities about automorphisms of trees. We conclude with an important simplicity criterion.

2.2.1. Definitions and notation. A graph Γ is a tuple (V, E) consisting of a vertex set V and an edge set E , together with a fixed-point-free involution of E , denoted by $e \mapsto \bar{e}$, and maps $o, t : E \rightarrow V$, providing the *origin* and *terminus* of an edge, such that $o(\bar{e}) = t(e)$ and $t(\bar{e}) = o(e)$ for all $e \in E$. Given $e \in E$, the pair $\{e, \bar{e}\}$ is a *geometric edge*. For $x \in V$, we let $E(x) := o^{-1}(x) = \{e \in E \mid o(e) = x\}$ be the set of edges issuing from x . The *valency* of $x \in V$ is $|E(x)|$. A vertex of valency 1 is a *leaf*. A *morphism* between graphs $\Gamma_1 = (V_1, E_1)$ and $\Gamma_2 = (V_2, E_2)$ is a pair (α_V, α_E) of maps $\alpha_V : V_1 \rightarrow V_2$ and $\alpha_E : E_1 \rightarrow E_2$ preserving the graph structure, that is, $\alpha_V(o(e)) = o(\alpha_E(e))$ and $\alpha_V(t(e)) = t(\alpha_E(e))$ for all $e \in E$.

For $n \in \mathbb{N}$, let Path_n denote the graph with vertex set $\{0, \dots, n\}$ and edge set $\{(k, k+1), (\bar{k}, \bar{k}+1) \mid k \in \{0, \dots, n-1\}\}$. A *path* of length n in a graph Γ is a morphism γ from Path_n to Γ . It can be identified with $(e_1, \dots, e_n) \in E(\Gamma)^n$, where $e_k = \gamma((k-1, k))$ for $k \in \{1, \dots, n\}$. In this case, γ is a path from $o(e_1)$ to $t(e_n)$.

Similarly, let $\text{Path}_{\mathbb{N}_0}$ and $\text{Path}_{\mathbb{Z}}$ be the graphs with vertex sets \mathbb{N}_0 and \mathbb{Z} , and edge sets $\{(k, k+1), (\bar{k}, \bar{k}+1) \mid k \in \mathbb{N}_0\}$ and $\{(k, k+1), (\bar{k}, \bar{k}+1) \mid k \in \mathbb{Z}\}$, respectively. A *half-infinite path*, or *ray*, in a graph Γ is a morphism γ from $\text{Path}_{\mathbb{N}_0}$ to Γ . It can be identified with $(e_k)_{k \in \mathbb{N}} \in E(\Gamma)^{\mathbb{N}}$ where $e_k = \gamma((k-1, k))$ for $k \in \mathbb{N}$. In this case, γ *originates at*, or *issues from*, $o(e_1)$. An *infinite path*, or *line*, in a graph Γ is a morphism from $\text{Path}_{\mathbb{Z}}$ to Γ . A pair $(e_k, e_{k+1}) = (e_k, \bar{e}_k)$ of edges in a path is a *backtracking*. A graph is *connected* if any two of its vertices can be joined by a path. The maximal connected subgraphs of a graph are its *connected components*.

A *forest* is a graph in which there are no nonbacktracking paths (e_1, \dots, e_n) with $o(e_1) = t(e_n)$, $n \in \mathbb{N}$. Consequently, a morphism of forests is determined by the underlying vertex map. In particular, a path of length $n \in \mathbb{N}$ in a forest is determined by the images of the vertices of Path_n .

A *tree* is a connected forest. As a consequence of the above, the vertex set V of a tree T admits a natural metric. Given $x, y \in V$, define $d(x, y)$ as the minimal length of a path from x to y . A tree in which every vertex has valency $d \in \mathbb{N}$ is *d-regular*. It is unique up to isomorphism and denoted by T_d .

Let $T = (V, E)$ be a tree. For $S \subseteq V \cup E$, the *subtree spanned by S* is the unique minimal subtree of T containing S . For $x \in V$ and $n \in \mathbb{N}_0$, the subtree spanned by $\{y \in V \mid d(y, x) \leq n\}$ is the *ball* of radius n around x , denoted by $B(x, n)$. Similarly, $S(x, n) = \{y \in V \mid d(y, x) = n\}$ is the *sphere* of radius n around x , and the set of edges within distance n of x is $E(x, n) := \{e \in E \mid d(o(e), x), d(t(e), x) \leq n\}$. For a subtree $T' \subseteq T$, let $\pi : V \rightarrow V(T')$ denote the closest point projection, that is, $\pi(x) = y$ whenever

$d(x, y) = \min_{z \in V(T')} \{d(x, z)\}$. In the case of an edge $e = (x, y) \in E$, the *half-trees* T_x and T_y are the subtrees spanned by $\pi^{-1}(x)$ and $\pi^{-1}(y)$, respectively.

Two nonbacktracking rays $\gamma_1, \gamma_2 : \text{Path}_{\mathbb{N}} \rightarrow T$ in T are *equivalent*, $\gamma_1 \sim \gamma_2$, if there exist $N, d \in \mathbb{N}$ such that $\gamma_1(n) = \gamma_2(n + d)$ for all $n \geq N$. The *boundary*, or *set of ends*, of T is the set ∂T of equivalence classes of nonbacktracking rays in T .

2.2.2. Automorphism groups of graphs. Let $\Gamma = (V, E)$ be a graph. We equip the group $\text{Aut}(\Gamma)$ of automorphisms of Γ with the permutation topology for its action on $V \cup E$.

Notation. Let $H \leq \text{Aut}(\Gamma)$. Given a subgraph $\Gamma' \subseteq \Gamma$, the *pointwise stabilizer* of Γ' in H is denoted by $H_{\Gamma'}$. Similarly, the *setwise stabilizer* of Γ' in H is denoted by $H_{[\Gamma']}$. In the case where Γ' is a single vertex x , the permutation group that H_x induces on $E(x)$ is denoted by $H_x^{(1)} \leq \text{Sym}(E(x))$. Given a property ‘X’ of permutation groups, the group H is *locally ‘X’* if for every $x \in V$ the permutation group $H_x^{(1)}$ has ‘X’; with the exception that H is *locally k -transitive* ($k \in \mathbb{N}_{\geq 3}$) if H_x acts transitively on the set of nonbacktracking paths of length k issuing from x . It is *locally ∞ -transitive* if it is locally k -transitive for all $k \in \mathbb{N}$.

Let $d \in \mathbb{N}_{\geq 3}$ and $T_d = (V, E)$ the d -regular tree. Then $\text{Aut}(T_d)$ acts on ∂T_d by $g \cdot [\gamma] := [g \circ \gamma]$. Given $[\gamma] \in \partial T_d$, the *stabilizer* of $[\gamma]$ in H is $H_{[\gamma]} = \{h \in H \mid h \circ \gamma \sim \gamma\}$.

We let ${}^+H = \langle \{H_x \mid x \in V\} \rangle$ denote the subgroup of H generated by vertex-stabilizers and $H^+ = \langle \{H_e \mid e \in E\} \rangle$ the subgroup generated by edge-stabilizers. For a subtree $T \subseteq T_d$ and $k \in \mathbb{N}$, let T^k denote the subtree of T_d spanned by $\{x \in V \mid d(x, T) \leq k\}$. We set $H^{+k} = \langle \{H_{e^{k-1}} \mid e \in E\} \rangle$. Then $H^{+1} = H^+$ and

$$H^{+k} \trianglelefteq H^+ \trianglelefteq {}^+H \trianglelefteq H.$$

Classification of automorphisms. Automorphisms of T_d can be divided into three distinct types. Refer to [10, Section 6.2.2] for details.

For $g \in \text{Aut}(T_d)$, set $l(g) := \min_{x \in V} d(x, gx)$ and $V(g) := \{x \in V \mid d(x, gx) = l(g)\}$. If $l(g) = 0$ then g fixes a vertex. An automorphism of this kind is *elliptic*. Suppose now that $l(g) > 0$. If $V(g)$ is infinite then g is *hyperbolic*. Geometrically, it is a translation of length $l(g)$ along the line in T_d defined by $V(g)$. If $V(g)$ is finite then $l(g) = 1$ and g maps some edge $e \in E$ to \bar{e} and is termed an *inversion*.

Independence and simplicity. The base case of the simplicity criterion presented below is due to Tits [29] and applies to sufficiently rich subgroups of $\text{Aut}(T_d)$. The generalized version is due to Banks *et al.* [1]; see also [10].

Let C denote a path in T_d (finite, half-infinite or infinite). For every $x \in V(C)$ and $k \in \mathbb{N}_0$, the pointwise stabilizer H_{C^k} of C^k induces an action $H_{C^k}^{(x)} \leq \text{Aut}(\pi^{-1}(x))$ on $\pi^{-1}(x)$, the subtree spanned by those vertices of T whose closest vertex in C is x . We therefore obtain an injective homomorphism

$$\varphi_C^{(k)} : H_{C^k} \rightarrow \prod_{x \in V(C)} H_{C^k}^{(x)}.$$

A subgroup $H \leq \text{Aut}(T_d)$ satisfies *property* (P_k) ($k \in \mathbb{N}$) if $\varphi_C^{(k-1)}$ is an isomorphism for every path C in T_d . If $H \leq \text{Aut}(T_d)$ is closed, it suffices to check the above properties in the case where C is a single edge. For example, given a closed subgroup $H \leq \text{Aut}(T_d)$, property (P_k) is satisfied by its (P_k) -closure:

$$H^{(P_k)} = \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V(T_d) \text{ there exists } h \in H : g|_{B(x,k)} = h|_{B(x,k)}\}.$$

THEOREM 2.1 [1, Theorem 7.3]. *Let $H \leq \text{Aut}(T_d)$. Suppose H neither fixes an end nor stabilizes a proper subtree of T_d setwise, and that H satisfies property (P_k) . Then the group H^{+k} is either trivial or simple.*

2.3. Burger–Mozes theory. In [2], Burger and Mozes develop a structure theory of certain locally quasiprimitive automorphism groups of graphs which resembles the theory of semisimple Lie groups. Their fundamental definitions are meaningful in the setting of t.d.l.c. groups. Let H be a t.d.l.c. group. Define

$$H^{(\infty)} := \bigcap \{N \trianglelefteq H \mid N \text{ is closed and cocompact in } H\},$$

alternatively the intersection of all open finite-index subgroups of H , and

$$\text{QZ}(H) := \{h \in H \mid Z_H(h) \leq H \text{ is open}\},$$

the *quasicentre* of H . Both $H^{(\infty)}$ and $\text{QZ}(H)$ are topologically characteristic subgroups of H , that is, they are preserved by continuous automorphisms of H . Whereas $H^{(\infty)} \leq H$ is closed, the quasicentre need not be so.

Whereas for a general t.d.l.c. group H nothing much can be said about the size of $H^{(\infty)}$ and $\text{QZ}(H)$, Burger and Mozes show that good control can be obtained in the case of certain locally quasiprimitive automorphism groups of graphs. The following result summarizes their structure theory. It is a combination of Proposition 1.2.1, Corollary 1.5.1, Theorem 1.7.1 and Corollary 1.7.2 in [2].

THEOREM 2.2. *Let Γ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be closed, nondiscrete and locally quasiprimitive. Then*

- (i) $H^{(\infty)}$ is minimal closed normal cocompact in H ;
- (ii) $\text{QZ}(H)$ is maximal discrete normal, and noncocompact in H ; and
- (iii) $H^{(\infty)}/\text{QZ}(H^{(\infty)}) = H^{(\infty)}/(\text{QZ}(H) \cap H^{(\infty)})$ admits minimal, nontrivial closed normal subgroups finite in number, H -conjugate and topologically simple.

If Γ is a tree and moreover H is locally primitive then

- (iv) $H^{(\infty)}/\text{QZ}(H^{(\infty)})$ is a direct product of topologically simple groups.

2.3.1. Burger–Mozes universal groups. The first introduction of Burger–Mozes universal groups in [2, Section 3.2] was expanded in the introductory article [10], which we follow closely. Most results are generalized in Section 4.

Let Ω be a set of cardinality $d \in \mathbb{N}_{\geq 3}$ and let $T_d = (V, E)$ denote the d -regular tree. A labelling l of T_d is a map $l : E \rightarrow \Omega$ such that for every $x \in V$ the map

$l_x : E(x) \rightarrow \Omega$, $e \mapsto l(e)$ is a bijection, and $l(e) = l(\bar{e})$ for all $e \in E$. The *local action* $\sigma(g, x) \in \text{Sym}(\Omega)$ of an automorphism $g \in \text{Aut}(T_d)$ at a vertex $x \in V$ is defined via

$$\sigma : \text{Aut}(T_d) \times X \rightarrow \text{Sym}(\Omega), (g, x) \mapsto \sigma(g, x) := l_{gx} \circ g \circ l_x^{-1}.$$

DEFINITION 2.3. Let $F \leq \text{Sym}(\Omega)$ and l a labelling of T_d . Define

$$U^{(l)}(F) := \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V : \sigma(g, x) \in F\}.$$

The map σ satisfies a *cocycle identity*: for all $g, h \in \text{Aut}(T_d)$ and $x \in V$ we have $\sigma(gh, x) = \sigma(g, hx)\sigma(h, x)$. As a consequence, $U^{(l)}(F)$ is a subgroup of $\text{Aut}(T_d)$.

Passing to a different labelling amounts to passing to a conjugate of $U^{(l)}(F)$ inside $\text{Aut}(T_d)$. We therefore omit reference to an explicit labelling from here onwards.

The following proposition collects several basic properties of Burger–Mozes groups. We refer the reader to [10, Section 6.4] for proofs.

PROPOSITION 2.4. Let $F \leq \text{Sym}(\Omega)$. The group $U(F)$ is

- (i) closed in $\text{Aut}(T_d)$,
- (ii) vertex-transitive,
- (iii) compactly generated,
- (iv) locally permutation isomorphic to F ,
- (v) edge-transitive if and only if F is transitive, and
- (vi) discrete if and only if F is semiregular.

Part (iii) of Proposition 2.4 relies on the following result which we include for future reference. Given $x \in V$ and $\omega \in \Omega$, let $\iota_\omega^{(x)} \in U(\{\text{id}\})$ denote the unique label-respecting inversion of the edge $e_\omega \in E$ with $o(e_\omega) = x$ and $l(e_\omega) = \omega$.

LEMMA 2.5. Let $x \in V$. Then $U(\{\text{id}\}) = \langle \{\iota_\omega^{(x)} \mid \omega \in \Omega\} \rangle \cong \prod_{\omega \in \Omega} \langle \iota_\omega^{(x)} \rangle \cong \prod_{\omega \in \Omega} \mathbb{Z}/2\mathbb{Z}$.

PROOF. Every element of $U(\{\text{id}\})$ is determined by its image on x . Hence, it suffices to show that $\langle \{\iota_\omega^{(x)} \mid \omega \in \Omega\} \rangle$ is vertex-transitive and has the asserted structure. Indeed, let $y \in V \setminus \{x\}$, and let $\omega_1, \dots, \omega_n \in \Omega$ be the labels of the shortest path from x to y . Then $\iota_{\omega_1}^{(x)} \circ \dots \circ \iota_{\omega_n}^{(x)}$ maps x to y as every $\iota_\omega^{(x)}$ ($\omega \in \Omega$) is label-respecting. Setting $X_\omega := T_{l(e_\omega)}$, we have $\iota_\omega(X_{\omega'}) \subseteq X_\omega$ for all distinct $\omega, \omega' \in \Omega$. Hence, the assertion follows from the ping-pong lemma. □

The name *universal group* is due to the following maximality statement. Its proof (see [2, Proposition 3.2.2]) should be compared with the proof of Theorem 4.34.

PROPOSITION 2.6. 5 Let $H \leq \text{Aut}(T_d)$ be locally transitive and vertex-transitive. Then there is a labelling l of T_d such that $H \leq U^{(l)}(F)$ where $F \leq \text{Sym}(\Omega)$ is action isomorphic to the local action of H .

3. Structure theory of locally semiprimitive groups

We generalize the Burger–Mozes theory of locally quasiprimitive automorphism groups of graphs to the semiprimitive case. While this adjustment of Sections 1.1–1.5

in [2] is straightforward and was initiated in [30, Section II.7] and [3, Section 6.2], we provide a full account for the reader's convenience.

3.1. General facts. Let $\Gamma = (V, E)$ be a connected graph. We first present a few general facts about several classes of subgroups of $\text{Aut}(\Gamma)$ for future reference.

LEMMA 3.1. *Let $H \leq \text{Aut}(\Gamma)$ be locally transitive. Then ${}^+H$ is geometric edge transitive and of index at most 2 in H .*

PROOF. Since H is locally transitive, so is ${}^+H$, given that ${}^+H_x = H_x$ for all $x \in V$. Hence, it is geometric edge transitive. In particular, it has at most two vertex orbits, which implies the second assertion. \square

LEMMA 3.2. *Let $H \leq \text{Aut}(\Gamma)$ and let $\Gamma' = (V', E')$ be a connected subgraph of Γ . Suppose $R \subseteq H$ is such that for every $x' \in V'$ and $e \in E(x')$ there is $r \in R$ such that $re \in E'$. Then $\Lambda := \langle R \rangle$ satisfies $\bigcup_{\lambda \in \Lambda} \lambda \Gamma' = \Gamma$.*

PROOF. By assumption, $B(\Gamma', 1) \subseteq \bigcup_{\lambda \in \Lambda} \lambda \Gamma'$. Now suppose $B(\Gamma', n) \subseteq \bigcup_{\lambda \in \Lambda} \lambda \Gamma'$ for some $n \in \mathbb{N}$. Let $x' \in V(B(\Gamma', n))$. Pick $\lambda \in \Lambda$ such that $\lambda(x') \in V'$. Since λ induces a bijection between $E(x')$ and $E(\lambda(x'))$ we conclude that $B(\Gamma', n + 1) \subseteq \bigcup_{\lambda \in \Lambda} \lambda \Gamma'$. \square

Assume from now on that Γ is a locally finite, connected graph.

LEMMA 3.3. *Let $H \leq \text{Aut}(\Gamma)$. If $H \setminus \Gamma$ is finite then there is a finitely generated subgroup $\Lambda \leq H$ such that $\Lambda \setminus \Gamma$ is finite.*

PROOF. Let $\Gamma' = (V', E') \subseteq \Gamma$ be a connected subgraph that projects onto $H \setminus \Gamma$. For every $x' \in V'$ and $e \in E(x')$, pick $\lambda_{x',e} \in H$ such that $\lambda_{x',e}(e) \in E'$. Then the group $\Lambda := \langle \{\lambda_{x',e} \mid x' \in X, e \in E(x')\} \rangle$ satisfies the conclusion by Lemma 3.2. \square

LEMMA 3.4. *Let $\Lambda \leq \text{Aut}(\Gamma)$. If $\Lambda \setminus \Gamma$ is finite then $Z_{\text{Aut}(\Gamma)}(\Lambda)$ is discrete.*

PROOF. Let $F \subseteq E$ be finite such that $\bigcup_{\lambda \in \Lambda} \lambda F = E$ and $U := \Lambda_F \cap Z_{\text{Aut}(\Gamma)}(\Lambda)$, which is open in $Z_{\text{Aut}(\Gamma)}(\Lambda)$. Given that U and Λ commute, U acts trivially on $E = \bigcup_{\lambda \in \Lambda} \lambda F$. Hence, $U = \{\text{id}\}$ and $Z_{\text{Aut}(\Gamma)}(\Lambda)$ is discrete. \square

LEMMA 3.5. *Let $\Lambda_1, \Lambda_2 \leq \text{Aut}(\Gamma)$. If $\Lambda_1 \setminus \Gamma$ is finite and $[\Lambda_1, \Lambda_2] \leq \text{Aut}(\Gamma)$ is discrete then $\Lambda_2 \leq \text{Aut}(\Gamma)$ is discrete.*

PROOF. Using Lemma 3.3 pick $R \subseteq \Lambda_1$ such that $\langle R \rangle \setminus \Gamma$ is finite. As $[\Lambda_1, \Lambda_2] \leq \text{Aut}(\Gamma)$ is discrete, there is an open subgroup $U \leq \Lambda_2$ such that $[r, U] = \{e\}$ for all $r \in R$. That is, $U \leq Z_{\text{Aut}(\Gamma)}(\langle R \rangle)$. Hence, U is discrete by Lemma 3.4, and so is Λ_2 . \square

LEMMA 3.6. *Let $H \leq \text{Aut}(\Gamma)$ be nondiscrete. Then $\text{QZ}(H) \setminus \Gamma$ is infinite.*

PROOF. If $\text{QZ}(H) \setminus \Gamma$ is finite, there is a finitely generated subgroup $\Lambda \leq \text{QZ}(H)$ such that $\Lambda \setminus \Gamma$ is finite as well by Lemma 3.3. Hence, there is an open subgroup $U \leq H$ with $U \leq Z_{\text{Aut}(\Gamma)}(\Lambda)$. Hence, U and therefore H are discrete by Lemma 3.4. \square

LEMMA 3.7. *Let $\Lambda \leq \text{Aut}(\Gamma)$ be discrete. If $\Lambda \setminus \Gamma$ is finite then $N_{\text{Aut}(\Gamma)}(\Lambda)$ is discrete.*

PROOF. Apply Lemma 3.5 to $\Lambda_1 := \Lambda$ and $\Lambda_2 := N_{\text{Aut}(\Gamma)}(\Lambda)$. \square

3.2. Normal subgroups. Let $\Gamma = (V, E)$ denote a locally finite, connected graph. For closed subgroups $\Lambda \trianglelefteq H$ of $\text{Aut}(\Gamma)$ we define

$$\mathcal{N}_{\text{nf}}(H, \Lambda) = \{N \trianglelefteq H \mid \Lambda \leq N \trianglelefteq H, N \text{ is closed and does not act freely on } E\},$$

the set of closed normal subgroups of H that contain Λ and do not act freely on E . The set $\mathcal{N}_{\text{nf}}(H, \Lambda)$ is partially ordered by inclusion. We let $\mathcal{M}_{\text{nf}}(H, \Lambda) \subseteq \mathcal{N}_{\text{nf}}(H, \Lambda)$ denote the set of minimal elements in $\mathcal{N}_{\text{nf}}(H, \Lambda)$.

LEMMA 3.8. *Let $\Gamma = (V, E)$ be a locally finite, connected graph and $\Lambda \trianglelefteq H \leq \text{Aut}(\Gamma)$. If $H \setminus \Gamma$ is finite and H does not act freely on E then $\mathcal{M}_{\text{nf}}(H, \Lambda) \neq \emptyset$.*

PROOF. We argue using Zorn’s lemma. First note that $\mathcal{N}_{\text{nf}}(H, \Lambda)$ is nonempty as it contains H . Let $C \subseteq \mathcal{N}_{\text{nf}}(H, \Lambda)$ be a chain. Pick a finite set $F \subseteq E$ of representatives of $H \setminus E$. For every $N \in C$, the set $F_N := \{e \in F \mid N|_{e^1} \leq \text{Aut}(e^1)\}$ is nontrivial} is nonempty. Since F is finite and C is a chain it follows that $\bigcap_{N \in C} F_N$ is nonempty, that is, there exists $e \in F$ such that $N|_{e^1}$ is nontrivial for every $N \in C$. As before, we conclude that $M := \bigcap_{N \in C} N|_{e^1}$ is nontrivial. For $\alpha \in M \setminus \{\text{id}\}$, $N^\alpha := \{g \in N_e \mid g|_{e^1} = \alpha\}$ is a nonempty compact subset of H_e , and since C is a chain every finite subset of $\{N^\alpha \mid N \in C\}$ has nonempty intersection. Hence, $\bigcap_{N \in C} N^\alpha$ is nonempty and therefore $N_C := \bigcap_{N \in C} N$ is a closed normal subgroup of H containing Λ that does not act freely on E . Overall, $N_C \in \mathcal{M}_{\text{nf}}(H, \Lambda)$. □

The following lemma is contained in the author’s PhD thesis [30, Section II.7] and, independently, in Caprace and Le Boudec [3, Section 6.2].

LEMMA 3.9. *Let $\Gamma = (V, E)$ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be locally semiprimitive and $N \trianglelefteq H$. Define*

$$V_1 := \{x \in V \mid N_x \curvearrowright S(x, 1) \text{ is transitive and not semiregular}\},$$

$$V_2 := \{x \in V \mid N_x \curvearrowright S(x, 1) \text{ is semiregular}\}.$$

Then one of the following assertions holds.

- (i) $V = V_2$ and N acts freely on E .
- (ii) $V = V_1$ and N is geometric edge transitive.
- (iii) $V = V_1 \sqcup V_2$ is an H -invariant partition of V and $B(x, 1)$ is a fundamental domain for the action of N on Γ for any $x \in V_2$.

PROOF. Since H is locally semiprimitive and N is normal in H , we have $V = V_1 \sqcup V_2$. If N does not act freely on E then there exist an edge $e \in E$ with $N_e \neq \{\text{id}\}$ and an N_e -fixed vertex $x \in V$ for which $N_x \curvearrowright S(x, 1)$ is not semiregular, hence transitive. That is, $V_1 \neq \emptyset$. Now, either $V_2(N) = \emptyset$ in which case N is locally transitive and we are in case (ii), or $V_2(N) \neq \emptyset$. Being locally transitive, H acts transitively on the set of geometric edges and therefore has at most two vertex orbits. Given that both V_1 and V_2 are nonempty and H -invariant, they constitute exactly the said orbits. Since any pair of adjacent vertices (x, y) is a fundamental domain for the H -action on V , we conclude

that if $y \in V_2$ then $x \in V_1$. Thus every leaf of $B(y, 1)$ is in V_1 and we are in case (iii) by Lemma 3.2. \square

3.3. The subquotient $H^{(\infty)}/QZ(H^{(\infty)})$. In this section, we achieve control over $H^{(\infty)}$ and $QZ(H)$ as well as the normal subgroups of H in the semiprimitive case. We then describe the structure of the subquotient $H^{(\infty)}/QZ(H^{(\infty)})$. First, recall the following lemma from topological group theory.

LEMMA 3.10. *Let G be a topological group. If $H \trianglelefteq G$ is discrete then $H \subseteq QZ(G)$.*

PROOF. For $h \in H$, the map $\varphi_h : G \rightarrow H$, $g \mapsto ghg^{-1}$ is well defined because $H \trianglelefteq G$, and continuous. Hence, there is an open set $U \subseteq G$ containing $1 \in G$ and such that $\varphi_h(U) \subseteq \{h\}$, that is, $U \subseteq Z_G(h)$. \square

PROPOSITION 3.11. *Let $\Gamma = (V, E)$ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be closed, nondiscrete and locally semiprimitive. Then the following assertions hold.*

- (i) $H/H^{(\infty)}$ is compact.
- (ii) $QZ(H)$ acts freely on E , and is discrete noncocompact in H .
- (iii) For any closed normal subgroup $N \trianglelefteq H$, either N is nondiscrete cocompact and $N \supseteq H^{(\infty)}$, or N is discrete and $N \trianglelefteq QZ(H)$.
- (iv) $QZ(H^{(\infty)}) = QZ(H) \cap H^{(\infty)}$ acts freely on E without inversions.
- (v) For any open normal subgroup $N \trianglelefteq H^{(\infty)}$ we have $N = H^{(\infty)}$.
- (vi) $H^{(\infty)}$ is topologically perfect, that is, $H^{(\infty)} = [H^{(\infty)}, H^{(\infty)}]$.

PROOF. For (i), let $N \trianglelefteq H$ be closed and cocompact. Since H is nondiscrete, so is N in view of Lemma 3.7. Hence, $N \in \mathcal{N}_{\text{nf}}(H, \{\text{id}\})$. Conversely, if $N \in \mathcal{N}_{\text{nf}}(H, \{\text{id}\})$ then N is cocompact in H by Lemma 3.9. We conclude that $H^{(\infty)} = \bigcap \mathcal{N}_{\text{nf}}(H, \{\text{id}\})$. This intersection is in fact given by a single minimal element of $\mathcal{N}_{\text{nf}}(H, \{\text{id}\})$. Using Lemma 3.8, pick $M \in \mathcal{M}_{\text{nf}}(H, \{\text{id}\})$, and let $N \in \mathcal{N}_{\text{nf}}(H, \{\text{id}\})$. Suppose $N \not\supseteq M$. Because M is minimal, $N \cap M$ acts freely on E . In particular, $N \cap M$ is discrete. Since both N and M are normal in H , we also have $N \cap M \supseteq [N, M]$ and hence N and M are discrete by Lemma 3.5. Then so is $H \subseteq N_{\text{Aut}(\mathfrak{g})}(H)$ by Lemma 3.7. Overall, $H^{(\infty)} = M \in \mathcal{M}_{\text{nf}}(H, \{\text{id}\})$ and the assertion now follows from Lemma 3.9.

As to (ii), the group $QZ(H)$ is noncocompact by Lemma 3.6 and therefore acts freely on E by Lemma 3.9. In particular, it is discrete.

For (iii), let $N \trianglelefteq H$ be a closed normal subgroup. If N acts freely on E , then N is discrete and hence contained in $QZ(H)$ by Lemma 3.10. If N does not act freely on E then N is cocompact in H by Lemma 3.9 and therefore contains $H^{(\infty)}$.

Concerning (iv), the inclusion $QZ(H) \cap H^{(\infty)} \subseteq QZ(H^{(\infty)})$ is automatic. Further, $QZ(H^{(\infty)})$ is normal in H because it is topologically characteristic in $H^{(\infty)} \trianglelefteq H$. Therefore, if $QZ(H^{(\infty)}) \not\subseteq QZ(H)$, then $QZ(H^{(\infty)})$ is nondiscrete by part (iii) and does not act freely on E . Then $QZ(H^{(\infty)}) \setminus \Gamma$ is finite by Lemma 3.9, contradicting Lemma 3.6 applied to $H^{(\infty)}$ which is nondiscrete because $QZ(H^{(\infty)}) \leq H^{(\infty)}$ is. Consequently, $QZ(H^{(\infty)}) \leq QZ(H)$, which proves the assertion.

For part (v), note that $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \{\text{id}\})$ is nonempty by Lemma 3.8 as $H^{(\infty)}$ is cocompact in $\text{Aut}(\Gamma)$ by part (i) and nondiscrete by part (iii). Further, since $\text{QZ}(H^{(\infty)})$ acts freely on E , every $N \in \mathcal{N}_{\text{nf}}(H^{(\infty)}, \{\text{id}\})$ is nondiscrete by part (iii) as well. Given an open subgroup $U \trianglelefteq H^{(\infty)}$ and $N \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \{\text{id}\})$, the group $U \cap N$ is normal in $H^{(\infty)}$ and nondiscrete. In particular, $U \cap N$ does not act freely on E and hence $U \cap N = N$. Thus U contains the subgroup of $H^{(\infty)}$ generated by the elements of $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \{\text{id}\})$, which is closed, normal and nondiscrete. Hence, $U = H^{(\infty)}$.

As to (vi), the group $[H^{(\infty)}, H^{(\infty)}]$ is nondiscrete by part (i) and Lemma 3.5. Hence, so is $[H^{(\infty)}, H^{(\infty)}] \trianglelefteq H^{(\infty)}$. Now apply part (iii). □

PROPOSITION 3.12. *Let $\Gamma = (V, E)$ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be closed, nondiscrete and locally semiprimitive. Finally, let $\Lambda \trianglelefteq H$ such that $\Lambda \leq \text{QZ}(H^{(\infty)})$. Then the following assertions hold.*

- (i) (a) *The group H acts transitively on $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$.*
 (b) *The set $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is finite and nonempty.*
- (ii) *Let $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$.*
 - (a) *The group M/Λ is topologically perfect.*
 - (b) *The group $\text{QZ}(M)$ acts freely on E and $\text{QZ}(M) = \text{QZ}(H^{(\infty)}) \cap M$.*
 - (c) *The group $M/\text{QZ}(M)$ is topologically simple.*
- (iii) *For every $N \in \mathcal{N}_{\text{nf}}(H^{(\infty)}, \Lambda)$ there is $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ with $N \supseteq M$.*

PROOF. Since every discrete normal subgroup of $H^{(\infty)}$ is contained in $\text{QZ}(H^{(\infty)})$ by Lemma 3.10 (iii), and the latter acts freely on E by Proposition 3.11(iii), every element of $\mathcal{N}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is nondiscrete. We proceed with a number of claims.

- (1) For every $N \in \mathcal{N}_{\text{nf}}(H^{(\infty)}, \Lambda)$ we have $[H^{(\infty)}, N] \not\subseteq \text{QZ}(H^{(\infty)})$.

This follows from the above combined with Proposition 3.11(i) and Lemma 3.5.

In the following, given $S \subseteq \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$, we let $M_S := \langle M \mid M \in S \rangle \leq H^{(\infty)}$ denote the subgroup of $H^{(\infty)}$ generated by $\bigcup_{M \in S} M$.

- (2) The group H acts transitively on $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$.
 Let S be an orbit for the action of H on $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$, and suppose there is an element $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda) \setminus S$. For every $N \in S$, the subgroup $N \cap M$ is normal in $H^{(\infty)}$ and acts freely on E by minimality of M , hence is discrete. The same therefore holds for $[N, M] \subseteq N \cap M$. Thus $[N, M] \subseteq \text{QZ}(H^{(\infty)})$. As $\text{QZ}(H^{(\infty)})$ is discrete by Proposition 3.11 and therefore closed in $H^{(\infty)}$ we conclude $\overline{[M_S, M]} \subseteq \text{QZ}(H^{(\infty)})$. On the other hand, $\overline{M_S}$ is normal in H since S is an H -orbit. It is also closed in H , and nondiscrete by the above. Thus $\overline{M_S} = H^{(\infty)}$ by Proposition 3.11(iii), and $[H^{(\infty)}, M] \subseteq \text{QZ}(H^{(\infty)})$, which contradicts part (1).
- (3) For every $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ we have $\overline{[M, M]} \cdot \Lambda = M$.
 Note that $\overline{[M, M]} \cdot \Lambda$ is a group because Λ is normal in M . Suppose there is an element $M_0 \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ with $\overline{[M_0, M_0]} \cdot \Lambda \leq M_0$. Then $\overline{[M_0, M_0]} \cdot \Lambda$ acts freely on E by minimality of M_0 and is discrete. Since $[M_0, M_0]$ is also

normal in $H^{(\infty)}$, we obtain $[M_0, M_0] \subseteq \text{QZ}(H^{(\infty)})$. Part (2) now implies that $[M, M] \subseteq \text{QZ}(H^{(\infty)})$ for all $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$. Given that $[M, M'] \subseteq \text{QZ}(H^{(\infty)})$ for all distinct M, M' in $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ as well, we conclude that $[H^{(\infty)}, H^{(\infty)}] \subseteq \text{QZ}(H^{(\infty)})$, which contradicts part (1).

- (4) For every $N \in \mathcal{N}_{\text{nf}}(H^{(\infty)}, \Lambda)$ there is $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ with $N \supseteq M$. Let $S := \{M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda) \mid N \not\supseteq M\}$. Then $[\overline{M_S}, N] \subseteq \text{QZ}(H^{(\infty)})$ as above. On the other hand, for $T := \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$, the group $\overline{M_T} \subseteq H^{(\infty)}$ is closed, nondiscrete and normal in H , thus $\overline{M_T} = H^{(\infty)}$. Using (1), we conclude that $S \neq T$, which proves the assertion.
- (5) Let S, S' be disjoint subsets of $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$. Then $\overline{M_S} \cap \overline{M_{S'}} \subseteq \text{QZ}(H^{(\infty)})$. If not, we have $\overline{M_S} \cap \overline{M_{S'}} \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ and there is, by part (4), an element $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ with $M \subseteq \overline{M_S} \cap \overline{M_{S'}}$. However, this implies that $[M, M] \subseteq [\overline{M_S}, \overline{M_{S'}}] \subseteq \text{QZ}(H^{(\infty)})$, which contradicts part (3).
- (6) The set $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is finite and nonempty.

The set $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is nonempty by Lemma 3.8. Let $G = \bigcup \overline{M_S}$, where the union is taken over all finite subsets S of the set $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$. Then G is nondiscrete and normal in H . Hence, $\overline{G} = H^{(\infty)}$ by Proposition 3.11(ii). Since H is second-countable and locally compact, it is metrizable. Hence, $H^{(\infty)}$ is a separable metric space and the same holds for G . Let $L \subseteq G$ be a countable dense subgroup, and fix an exhaustion $F_1 \subseteq F_2 \subseteq \dots \subseteq F$ of F by finite sets. Let $(S_n)_{n \in \mathbb{N}}$ be an increasing sequence of finite subsets of $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ such that $F_n \subseteq \overline{M_{S_n}}$. In particular,

$$L \subseteq \overline{M_{\bigcup_{n \in \mathbb{N}} S_n}} \quad \text{and thus} \quad \overline{M_{\bigcup_{n \in \mathbb{N}} S_n}} = H^{(\infty)},$$

which by (5) and (1) implies $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda) = \bigcup_{n \in \mathbb{N}} S_n$. Thus $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is countable. Next, fix $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$. Then $N_H(M)$ is closed and of countable index in H , and thus has nonempty interior as H is a Baire space. Hence, $N_H(M)$ is open in H . Given that $N_H(M)$ contains $H^{(\infty)}$, we conclude that $N_H(M)$ is of finite index in H using Proposition 3.11(i). Since H acts transitively on $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ by (2) we conclude that $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is finite by the orbit–stabilizer theorem.

The above claims yield parts (i)(a), (i)(b), (ii)(a) and (iii) of Proposition 3.12. We now turn to parts (ii)(b) and (ii)(c).

- (ii)(b) Using part (6), let $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda) = \{M_1, \dots, M_r\}$ and define

$$\Omega := \text{QZ}(M_1) \cdots \cdots \text{QZ}(M_r).$$

Note that since $\text{QZ}(M_i)$ is characteristic in M_i , which is normal in $H^{(\infty)}$, the quasicentres in the above definition normalize each other, so Ω is a group. It is then normal in H . If Ω does not act freely on E then $\Omega \backslash \Gamma$ is finite by Lemma 3.9 and there exist $\lambda_1, \dots, \lambda_k \in \Omega$ by Lemma 3.3 such that for $\Omega' := \langle \lambda_1, \dots, \lambda_k \rangle$ the quotient $\Omega' \backslash \Gamma$ is finite. For every $i \in \{1, \dots, k\}$, write $\lambda_i = a_i b_i$ where $a_i \in \text{QZ}(M_1)$ and $b_i \in \text{QZ}(M_2) \cdots \cdots \text{QZ}(M_r)$. Let $U_1 \leq M_1$ be an open

subgroup such that $[a_i, U_1] = \{e\}$ for all $i \in \{1, \dots, k\}$. Since $[M_2 \cdots M_r, M_1] \subseteq \text{QZ}(H^{(\infty)})$, there is an open subgroup $U_2 \leq M_1$ such that $[b_i, U_2] = \{e\}$ for all $i \in \{1, \dots, k\}$. Hence, $U := U_1 \cap U_2 \leq M_1$ is contained in $Z_{\text{Aut}(\Gamma)}(\Omega')$, which by Lemma 3.4 implies that U and hence M_1 is discrete, a contradiction. Thus Ω acts freely on E , is discrete and therefore $\Omega \subseteq \text{QZ}(H^{(\infty)})$. That is, $\text{QZ}(M_i) \subseteq \text{QZ}(H^{(\infty)}) \cap M_i$. The opposite inclusion follows from the definitions.

- (ii)(c) Let $M \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ and let $N \trianglelefteq M$ be a closed subgroup containing $\text{QZ}(M)$. For every $M' \in \mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ with $M \neq M'$ we have

$$[M', M] \subseteq M' \subseteq M \subseteq \text{QZ}(H^{(\infty)}).$$

This implies $[M', N] \subseteq \text{QZ}(H^{(\infty)}) \cap M = \text{QZ}(M) \subseteq N$; that is, M' normalizes N . Since $N \trianglelefteq M$, this implies $N \trianglelefteq H^{(\infty)}$; and hence, by minimality of M , we have either that $N = M$ or else that N acts freely on E and $N \subseteq \text{QZ}(H^{(\infty)}) \cap M = \text{QZ}(M)$. □

COROLLARY 3.13. *Let $\Gamma = (V, E)$ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be closed, nondiscrete and locally semiprimitive. Minimal, nontrivial closed normal subgroups of $H^{(\infty)}/\text{QZ}(H^{(\infty)})$ exist. They are all H -conjugate, finite in number and topologically simple.*

PROOF. Apply Proposition 3.12 to $\Lambda = \text{QZ}(H^{(\infty)})$. □

We summarize the previous results in the following theorem, which is a verbatim copy of Burger and Mozes' Theorem 2.2, except that the local action need only be semiprimitive, not quasiprimitive.

THEOREM 3.14. *Let Γ be a locally finite, connected graph. Further, let $H \leq \text{Aut}(\Gamma)$ be closed, nondiscrete and locally semiprimitive. Then the following assertions hold.*

- (i) $H^{(\infty)}$ is minimal closed normal cocompact in H .
- (ii) $\text{QZ}(H)$ is maximal discrete normal, and noncocompact in H .
- (iii) $H^{(\infty)}/\text{QZ}(H^{(\infty)}) = H^{(\infty)}/(\text{QZ}(H) \cap H^{(\infty)})$ admits minimal, nontrivial closed normal subgroups finite in number, H -conjugate and topologically simple.

If Γ is a tree, and, in addition, H is locally primitive then

- (iv) $H^{(\infty)}/\text{QZ}(H^{(\infty)})$ is a direct product of topologically simple groups.

PROOF. Parts (i) and (ii) stem from parts (i), (ii) and (iii) of Proposition 3.11 in combination with Section 2.3. For part (iii), use part (iv) of Proposition 3.11 and Corollary 3.13. Finally, part (iv) is Corollary 1.7.2 in [2]. It follows from Theorem 1.7.1 in [2] as the commutator of any two distinct elements in $\mathcal{M}_{\text{nf}}(H^{(\infty)}, \Lambda)$ is contained in $\text{QZ}(H^{(\infty)})$. □

4. Universal groups

In this section, we develop a generalization of Burger–Mozes universal groups that arises through prescribing the local action on balls of a given radius $k \in \mathbb{N}$ around vertices. The Burger–Mozes construction corresponds to the case $k = 1$.

Whereas many properties of the original construction carry over to the new set-up, others require adjustments. Notably, there are compatibility and discreteness conditions on the local action F under which the associated universal group is locally action isomorphic to F and discrete, respectively.

We then exhibit examples and (non)rigidity phenomena of our construction. Finally, a universality statement holds under an additional assumption.

4.1. Definition and basic properties.

4.1.1. Definition. Let Ω be a set of cardinality $d \in \mathbb{N}_{\geq 3}$ and let $T_d = (V, E)$ denote the d -regular tree. A labelling l of T_d is a map $l : E \rightarrow \Omega$ such that for every $x \in V$ the map $l_x : E(x) \rightarrow \Omega, e \mapsto l(e)$ is a bijection, and $l(e) = l(\bar{e})$ for all $e \in E$.

For every $k \in \mathbb{N}$, fix a tree $B_{d,k}$ that is isomorphic to a ball of radius k around a vertex in T_d . Let b denote its centre and carry over the labelling of T_d to $B_{d,k}$ via the chosen isomorphism. Then for every $x \in V$ there is a unique, label-respecting isomorphism $l_x^k : B(x, k) \rightarrow B_{d,k}$. We define the k -local action $\sigma_k(g, x) \in \text{Aut}(B_{d,k})$ of an automorphism $g \in \text{Aut}(T_d)$ at a vertex $x \in V$ via

$$\sigma_k : \text{Aut}(T_d) \times V \rightarrow \text{Aut}(B_{d,k}), (g, x) \mapsto \sigma_k(g, x) := l_{g x}^k \circ g \circ (l_x^k)^{-1}.$$

DEFINITION 4.1. Let $F \leq \text{Aut}(B_{d,k})$ and l be a labelling of T_d . Define

$$U_k^{(l)}(F) := \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V : \sigma_k(g, x) \in F\}.$$

The following lemma states that the maps σ_k satisfy a cocycle identity which implies that $U_k^{(l)}(F)$ is a subgroup of $\text{Aut}(T_d)$ for every $F \leq \text{Aut}(B_{d,k})$.

LEMMA 4.2. Let $x \in V$ and $g, h \in \text{Aut}(T_d)$. Then $\sigma_k(gh, x) = \sigma_k(g, hx)\sigma_k(h, x)$.

PROOF. We compute

$$\begin{aligned} \sigma_k(gh, x) &= l_{(gh)x}^k \circ gh \circ (l_x^k)^{-1} = l_{(gh)x}^k \circ g \circ h \circ (l_x^k)^{-1} \\ &= l_{(gh)x}^k \circ g \circ (l_{hx}^k)^{-1} \circ l_{hx}^k \circ h \circ (l_x^k)^{-1} = \sigma_k(g, hx)\sigma_k(h, x). \quad \square \end{aligned}$$

4.1.2. Basic properties. Note that the group $U_1^{(l)}(F)$ of Definition 4.1 coincides with the Burger–Mozes universal group $U_{(l)}(F)$ introduced in [2, Section 3.2] under the natural isomorphism $\text{Aut}(B_{d,1}) \cong \text{Sym}(\Omega)$. Several basic properties of the latter group carry over to the generalized set-up. First of all, passing between different labellings of T_d amounts to conjugating in $\text{Aut}(T_d)$. Subsequently, we therefore omit the reference to an explicit labelling.

LEMMA 4.3. For every quadruple (l, l', x, x') of labellings l, l' of T_d and vertices $x, x' \in V$, there is a unique automorphism $g \in \text{Aut}(T_d)$ with $gx = x'$ and $l' = l \circ g$.

PROOF. Set $gx := x'$. Now assume inductively that g is uniquely determined on $B(x, n)$, $n \in \mathbb{N}_0$, and let $v \in S(x, n)$. Then g is also uniquely determined on $E(v)$ by the requirement $l' = l \circ g$, namely $g|_{E(v)} := l_{E(gv)}^{-1} \circ l'|_{E(v)}$. \square

PROPOSITION 4.4. *Let $F \leq \text{Aut}(B_{d,k})$. Further, let l and l' be labellings of T_d . Then the groups $U_k^{(l)}(F)$ and $U_k^{(l')}(F)$ are conjugate in $\text{Aut}(T_d)$.*

PROOF. Choose $x \in V$. Let $\tau \in \text{Aut}(T_d)$ denote the automorphism of T_d associated to (l, l', x, x) by Lemma 4.3. Then $U_k^{(l)}(F) = \tau U_k^{(l')}(F) \tau^{-1}$. \square

The following basic properties of $U_k(F)$ are as in Proposition 2.4.

PROPOSITION 4.5. *Let $F \leq \text{Aut}(B_{d,k})$. The group $U_k(F)$ is*

- (i) *closed in $\text{Aut}(T_d)$,*
- (ii) *vertex-transitive, and*
- (iii) *compactly generated.*

PROOF. For (i), note that if $g \notin U_k(F)$ then $\sigma_k(g, x) \notin F$ for some $x \in V$. In this case, the open neighbourhood $\{h \in \text{Aut}(T_d) \mid h|_{B(x,k)} = g|_{B(x,k)}\}$ of g in $\text{Aut}(T_d)$ is also contained in the complement of $U_k(F)$.

For (ii), let $x, x' \in V$ and let $g \in \text{Aut}(T_d)$ be the automorphism of T_d associated to (l, l', x, x') by Lemma 4.3. Then $g \in U_k(F)$ as $\sigma_k(g, v) = \text{id} \in F$ for all $v \in V$.

To prove (iii), fix $x \in V$. We show that $U_k(F)$ is generated by the join of the compact set $U_k(F)_x$ and the finite generating set of $U_1(\{\text{id}\}) = U_k(\{\text{id}\}) \leq U_k(F)$ guaranteed by Lemma 2.5. Indeed, for $g \in U_k(F)$ pick g' in the finitely generated, vertex-transitive subgroup $U_1(\{\text{id}\})$ of $U_k(F)$ such that $g'gx = x$. We then have $g'g \in U_k(F)_x$ and the assertion follows. \square

For completeness, we explicitly state the following proposition.

PROPOSITION 4.6. *Let $F \leq \text{Aut}(B_{d,k})$. Then $U_k(F)$ is a compactly generated, totally disconnected, locally compact, second countable group.*

PROOF. The group $U_k(F)$ is totally disconnected, locally compact, second countable as a closed subgroup of $\text{Aut}(T_d)$, and compactly generated by Proposition 4.5. \square

Finally, we record that the groups $U_k(F)$ are (P_k) -closed.

PROPOSITION 4.7. *Let $F \leq \text{Aut}(B_{d,k})$. Then $U_k(F)$ satisfies property (P_k) .*

PROOF. Let $e = (x, y) \in E$. Clearly, $U_k(F)^{e^k} \supseteq U_k(F)^{e^k, T_y} \cdot U_k(F)^{e^k, T_x}$. Conversely, consider $g \in U_k(F)^{e^k}$ and define $g_y \in \text{Aut}(T_d)$ and $g_x \in \text{Aut}(T_d)$ by

$$\sigma_k(g_y, v) = \begin{cases} \sigma_k(g, v) & v \in V(T_x) \\ \text{id} & v \in V(T_y) \end{cases} \quad \text{and} \quad \sigma_k(g_x, v) = \begin{cases} \text{id} & v \in V(T_x) \\ \sigma_k(g, v) & v \in V(T_y), \end{cases}$$

respectively. Then $g_y \in U_k(F)^{e^k, T_y}$, $g_x \in U_k(F)^{e^k, T_x}$ and $g = g_y \circ g_x$. \square

4.2. Compatibility and discreteness. We now generalize parts (iv) and (vi) of Burger and Mozes' Proposition 2.4. There are compatibility and discreteness conditions (C) and (D) on subgroups $F \leq \text{Aut}(B_{d,k})$ that hold if and only if the associated universal group is locally action isomorphic to F and discrete, respectively.

We introduce the following notation for vertices in the labelled tree (T_d, l) . Given $x \in V$ and $\xi = (\omega_1, \dots, \omega_n) \in \Omega^n (n \in \mathbb{N}_0)$, set $x_\xi := \gamma_{x,\xi}(n)$ where

$$\gamma_{x,\xi} : \text{Path}_n^{(\xi)} := \begin{array}{c} \omega_1 \quad \omega_2 \quad \dots \quad \dots \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \dots \text{---} \bullet \\ 0 \quad 1 \quad 2 \quad \dots \quad n \end{array} \rightarrow T_d$$

is the unique label-respecting morphism sending 0 to $x \in V$. If ξ is the empty word, set $x_\xi := x$. Whenever admissible, we also adopt this notation in the case of $B_{d,k}$ and its labelling. In particular, $S(x, n)$ is in natural bijection with the set $\Omega^{(n)} := \{(\omega_1, \dots, \omega_n) \in \Omega^n \mid \text{for all } k \in \{1, \dots, n-1\} : \omega_{k+1} \neq \omega_k\}$.

4.2.1. Compatibility. First, we ask whether $U_k(F)$ locally acts like F , that is, whether the actions $U_k(F)_x \curvearrowright B(x, k)$ and $F \curvearrowright B_{d,k}$ are isomorphic for every $x \in V$. Whereas this always holds for $k = 1$ by Proposition 2.4(iv), it need not be true for $k \geq 2$, the issue being (non)compatibility among elements of F . See Example 4.9. The condition developed in this section allows for computations. A more practical version from a theoretical viewpoint follows in Section 4.4.

Now, let $x \in V$ and suppose that $\alpha \in U_k(F)_x$ realizes $a \in F$ at x , that is,

$$\alpha|_{B(x,k)} = (l_x^k)^{-1} \circ a \circ l_x^k.$$

Then given the condition that $\sigma_k(\alpha, x_\omega)$ is in F for all $\omega \in \Omega$, we obtain the following necessary *compatibility condition* on F for $U_k(F)$ to act like F at $x \in V$:

for all $a \in F$, for all $\omega \in \Omega$: there exists $a_\omega \in F$:

$$(l_x^k)^{-1} \circ a \circ l_x^k|_{S_\omega} = (l_{\alpha x_\omega}^k)^{-1} \circ a_\omega \circ l_{x_\omega}^k|_{S_\omega}$$

where $S_\omega := B(x, k) \cap B(x_\omega, k) \subseteq T_d$. Set $T_\omega := l_x^k(S_\omega) \subseteq B_{d,k}$. Then the above condition can be rewritten as

for all $a \in F$, for all $\omega \in \Omega$: there exists $a_\omega \in F$:

$$a_\omega|_{T_\omega} = l_{\alpha x_\omega}^k \circ (l_x^k)^{-1} \circ a \circ l_x^k \circ (l_{x_\omega}^k)^{-1}|_{T_\omega}.$$

Now note the following observations. First, αx_ω depends only on a . Second, the subtree T_ω of $B_{d,k}$ does not depend on x . Third, $\iota_\omega := l_x^k|_{S_\omega}^{T_\omega} \circ (l_{x_\omega}^k)^{-1}|_{T_\omega}^{S_\omega}$ is the unique nontrivial, involutive and label-respecting automorphism of T_ω ; it is given by

$$\iota_\omega := l_x^k|_{S_\omega}^{T_\omega} \circ (l_{x_\omega}^k)^{-1}|_{T_\omega}^{S_\omega} : T_\omega \rightarrow S_\omega \rightarrow T_\omega, \quad b_\xi \mapsto x_{\omega\xi} \mapsto b_{\omega\xi}$$

for admissible words ξ . Hence, the above condition may be rewritten as

$$\text{for all } a \in F, \text{ for all } \omega \in \Omega : \text{there exists } a_\omega \in F : a_\omega|_{T_\omega} = \iota_{a\omega} \circ a \circ \iota_\omega. \tag{C}$$

In this situation we say that a_ω is compatible with a in direction ω .

PROPOSITION 4.8. *Let $F \leq \text{Aut}(B_{d,k})$. Then $U_k(F)$ is locally action isomorphic to F if and only if F satisfies (C).*

PROOF. By the above, condition (C) is necessary. To show that it is also sufficient, let $x \in V$ and $a \in F$. We aim to define an automorphism $\alpha \in U_k(F)_x$ which realizes a at x . This forces us to define

$$\alpha|_{B(x,k)} := (l_x^k)^{-1} \circ a \circ l_x^k.$$

Now, assume inductively that α is defined consistently on $B(x, n)$ in the sense that $\sigma_k(\alpha, y) \in F$ for all $y \in B(x, n)$ with $B(y, k) \subseteq B(x, n)$. In order to extend α to $B(x, n + 1)$, let $y \in S(x, n - k + 1)$ and let $\omega \in \Omega$ be the unique label such that $y_\omega \in S(x, n - k)$. Set $c := \sigma_k(\alpha, y_\omega)$. Applying condition (C) to the pair (c, ω) yields an element $c_\omega \in F$ such that

$$(l_{\alpha y_\omega}^k)^{-1} \circ c \circ l_{y_\omega}^k|_{S_\omega} = (l_{\alpha y}^k)^{-1} \circ c_\omega \circ l_y^k|_{S_\omega},$$

where $S_\omega := B(y, k) \cap B(y_\omega, k)$ and we have realized

$$\iota_\omega \text{ as } l_{y_\omega}^k|_{S_\omega}^{T_\omega} \circ (l_y^k)^{-1}|_{T_\omega}^{S_\omega} \quad \text{and} \quad \iota_{c\omega} \text{ as } l_{\alpha y}^k|_{\alpha S_\omega}^{T_{c\omega}} \circ (l_{\alpha y_\omega}^k)^{-1}|_{T_{c\omega}}^{\alpha S_\omega}.$$

Now extend α consistently to $B(x, n + 1)$ by setting $\alpha|_{B(x,k)} := (l_{\alpha x}^k)^{-1} \circ c_\omega \circ l_x^k$. □

EXAMPLE 4.9. Let $\Omega := \{1, 2, 3\}$ and $a \in \text{Aut}(B_{3,2})$ be the element that swaps the leaves b_{12} and b_{13} of $B_{3,2}$. Then $F := \langle a \rangle = \{\text{id}, a\}$ does not contain an element compatible with a in direction $1 \in \Omega$ and hence does not satisfy condition (C).

We show that it suffices to check condition (C) on the elements of a generating set. Let $F \leq \text{Aut}(B_{d,k})$ and $a, b \in F$. Set $c := ab$. Then

$$\begin{aligned} c_\omega|_{T_\omega} &= \iota_{c\omega} \circ a \circ b \circ \iota_\omega = (\iota_{c\omega} \circ a \circ \iota_{b\omega}) \circ (\iota_{b\omega} \circ b \circ \iota_\omega) \\ &= (\iota_{a(b\omega)} \circ a \circ \iota_{b\omega}) \circ (\iota_{b\omega} \circ b \circ \iota_\omega). \end{aligned} \tag{M}$$

Let $C_F(a, \omega)$ denote the compatibility set of elements in F that are compatible with $a \in F$ in direction $\omega \in \Omega$. Then (M) shows that $C_F(ab, \omega) \supseteq C_F(a, b\omega)C_F(b, \omega)$. It therefore suffices to check condition (C) on a generating set of F .

Given $S \subseteq \Omega$, we also define $C_F(a, S) := \bigcap_{\omega \in S} C_F(a, \omega)$, the set of elements in F that are compatible with $a \in F$ in all directions from S . We omit F in this notation when it is clear from the context.

As a consequence, we obtain the following description of the local action of $U_k(F)$ when F does not satisfy condition (C).

PROPOSITION 4.10. *Let $F \leq \text{Aut}(B_{d,k})$. Then F has a unique maximal subgroup $C(F)$ that satisfies (C). We have $C(C(F)) = C(F)$ and $U_k(F) = U_k(C(F))$.*

PROOF. By the above, $C(F) := \langle H \leq F \mid H \text{ satisfies (C)} \rangle \leq F$ satisfies condition (C). It is the unique maximal such subgroup of F by definition, and $C(C(F)) = C(F)$.

Furthermore, $U_k(C(F)) \leq U_k(F)$. Conversely, suppose $g \in U_k(F) \setminus U_k(C(F))$. Then there is $x \in V$ such that $\sigma_k(g, x) \in F \setminus C(F)$ and the group

$$C(F) \leq \langle C(F), \{\sigma_k(g, x) \mid x \in V\} \rangle \leq F$$

satisfies condition (C), too, as can be seen by setting $\sigma_k(g, x)_\omega := \sigma_k(g, x_\omega)$. This contradicts the maximality of $C(F)$. □

REMARK 4.11. Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C). The proof of Proposition 4.8 shows that elements of $U_k(F)$ are readily constructed. Given $x, y \in V(T_d)$ and $a \in F$, define $g : B(x, k) \rightarrow B(y, k)$ by setting $g(x) = y$ and $\sigma_k(g, x) = a$. Then, given elements $a_\omega \in F (\omega \in \Omega)$ such that $a_\omega \in C_F(a, \omega)$ for all $\omega \in \Omega$, there is a unique extension of g to $B(x, k + 1)$ so that $\sigma_k(g, x_\omega) = a_\omega$ for all $\omega \in \Omega$. Proceed iteratively.

4.2.2. Discreteness. The group $F \leq \text{Aut}(B_{d,k})$ also determines whether or not $U_k(F)$ is discrete. In fact, the following proposition generalizes Proposition 2.4(vi).

PROPOSITION 4.12. *Let $F \leq \text{Aut}(B_{d,k})$. Then $U_k(F)$ is discrete if F satisfies*

$$\text{for all } \omega \in \Omega : F_{T_\omega} = \{\text{id}\}. \tag{D}$$

Conversely, if $U_k(F)$ is discrete and F satisfies (C), then F satisfies (D).

Alternatively, $U_k(F)$ is discrete if and only if $C(F)$ satisfies (D). Example 4.9 shows that condition (C) is necessary for the second part of Proposition 4.12.

Finally, note that F satisfies (D) if and only if $C_F(\text{id}, \omega) = \{\text{id}\}$ for all $\omega \in \Omega$.

PROOF OF PROPOSITION 4.12. Fix $x \in V$. A subgroup $H \leq \text{Aut}(T_d)$ is nondiscrete if and only if for every $n \in \mathbb{N}$ there is $h \in H \setminus \{\text{id}\}$ such that $h|_{B(x,n)} = \text{id}$.

Suppose that $U_k(F)$ is nondiscrete. Then there are $n \in \mathbb{N}_{\geq k}$ and $\alpha \in U_k(F)$ such that $\alpha|_{B(x,n)} = \text{id}$ and $\alpha|_{B(x,n+1)} \neq \text{id}$. Hence, there is $y \in S(x, n - k + 1)$ with $a := \sigma_k(\alpha, y) \neq \text{id}$. In particular, $a \in F_{T_\omega} \setminus \{\text{id}\}$ where ω is the label of the unique edge $e \in E$ with $o(e) = y$ and $d(x, y) = d(x, t(e)) + 1$.

Conversely, suppose that F satisfies (C) and $F_{T_\omega} \neq \{\text{id}\}$ for some $\omega \in \Omega$. Then for every $n \in \mathbb{N}_{\geq k}$, we define an automorphism $\alpha \in U_k(F)$ with $\alpha|_{B(x,n)} = \text{id}$ and $\alpha|_{B(x,n+1)} \neq \text{id}$. If $\alpha|_{B(x,n)} = \text{id}$, then $\sigma_k(\alpha, y) \in F$ for all $y \in B(x, n - k)$. Choose $e \in E$ with $y := o(e) \in S(x, n - k + 1)$ and $t(e) \in S(x, n - k)$ such that $l(e) = \omega$. We extend α to $B(y, k)$ by setting $\alpha|_{B(y,k)} := I_y^k \circ s \circ (I_y^k)^{-1}$ where $s \in F_{T_\omega} \setminus \{\text{id}\}$. Finally, we extend α to T_d using (C). □

We define the following condition (CD) on $F \leq \text{Aut}(B_{d,k})$ as the conjunction of (C) and (D):

$$\text{for all } a \in F, \text{ for all } \omega \in \Omega : \text{there exists } a_\omega \in F : a_\omega|_{T_\omega} = \iota_{a_\omega} \circ a \circ \iota_{a_\omega}. \tag{CD}$$

The following description is immediate from the above. When F satisfies (CD), an element of $U_k(F)_x$ is determined by its action on $B(x, k)$. Hence, $U_k(F)_x \cong F$ for every

$x \in V$ and $U_k(F)_{(x,y)} \cong F_{(b,b_\omega)}$ for every $(x, y) \in E$ with $l(x, y) = \omega$. Furthermore, F admits a unique *involutive compatibility cocycle*, that is, a map $z : F \times \Omega \rightarrow F$, $(a, \omega) \mapsto a_\omega$ which for all $a, b \in F$ and $\omega \in \Omega$ satisfies

- (i) (compatibility) $z(a, \omega) \in C_F(a, \omega)$,
- (ii) (cocycle) $z(ab, \omega) = z(a, b_\omega)z(b, \omega)$, and
- (iii) (involutive) $z(z(a, \omega), \omega) = a$.

Note that z restricts to an automorphism z_ω of $F_{(b,b_\omega)}$ ($\omega \in \Omega$) of order at most 2.

4.3. Group structure. For $\tilde{F} \leq \text{Aut}(B_{d,k})$, let $F := \pi\tilde{F} \leq \text{Sym}(\Omega)$ denote the projection of \tilde{F} onto $\text{Aut}(B_{d,1}) \cong \text{Sym}(\Omega)$. As an illustration, we record that the group structure of $U_k(\tilde{F})$ is particularly clear when F is regular.

PROPOSITION 4.13. *Let $\tilde{F} \leq \text{Aut}(B_{d,k})$ satisfy (C). Suppose $F := \pi\tilde{F}$ is regular. Then $U_k(\tilde{F}) = U_1(F) \cong F * \mathbb{Z}/2\mathbb{Z}$.*

PROOF. Fix $x \in V$. Since F is transitive, the group $U_k(\tilde{F})$ is generated by $U_k(\tilde{F})_x$ and an involution ι inverting an edge with origin x . Given $\alpha \in U_k(\tilde{F})_x$, regularity of F implies that $\sigma_1(\alpha, y) = \sigma_1(\alpha, x) \in F$ for all $y \in V$. Now, the subgroups $H_1 := U_k(\tilde{F})_x \cong F$ and $H_2 := \langle \iota \rangle$ of $U_k(\tilde{F})$ generate a free product within $U_k(\tilde{F})$ by the ping-pong lemma. Put $X_1 := V(T_x)$ and $X_2 := V(T_{x_\omega})$. Any nontrivial element of H_1 maps X_2 into X_1 as $F_\omega = \{\text{id}\}$, and $\iota \in H_2$ maps X_1 into X_2 . □

More generally, Bass–Serre theory [25] identifies the universal groups $U_k(F)$ as amalgamated free products, taking into account that $U_k(F)$ acts with inversions.

PROPOSITION 4.14. *Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C) (and (D)). If πF is transitive then*

$$U_k(F) \cong U_k(F)_x *_{U_k(F)_{(x,y)}} U_k(F)_{(x,y)} \left(\cong F *_{F_{(b,b_\omega)}} (F_{(b,b_\omega)} \rtimes \mathbb{Z}/2\mathbb{Z}) \right)$$

for any edge $(x, y) \in E$, where $\omega = l(x, y)$ and $\mathbb{Z}/2\mathbb{Z}$ acts on $F_{(b,b_\omega)}$ as z_ω .

COROLLARY 4.15. *Let $F, F' \leq \text{Aut}(B_{d,k})$ satisfy (CD). If there are $\omega, \omega' \in \Omega$ and an isomorphism $\varphi : F \rightarrow F'$ such that $\varphi(F_{(b,b_\omega)}) = F'_{(b,b_{\omega'})}$, then $U_k(F) \cong U_k(F')$.*

Note that Corollary 4.15 applies to conjugate subgroups of $\text{Aut}(B_{d,k})$ that satisfy (CD). The following example shows that the assumption that both F and F' in Corollary 4.15 satisfy (CD) is indeed necessary.

EXAMPLE 4.16. Let $\Omega := \{1, 2, 3\}$ and $t \in \text{Aut}(B_{3,2})$ be the element that swaps the leaves x_{12} and x_{13} of $B_{3,2}$. Using the notation of Section 4.4.1 below, consider the group $\Gamma(A_3) \leq \text{Aut}(B_{3,2})$ which satisfies (C). In particular, $U_2(\Gamma(A_3)) \cong A_3 * \mathbb{Z}/2\mathbb{Z}$ by Proposition 4.13. On the other hand, set $F' := t\Gamma(A_3)t^{-1}$. Then $\pi F' = A_3$, while for a nontrivial element α of F' we have $\sigma_1(\alpha, b_\omega) \in S_3 \setminus A_3$ for some $\omega \in \Omega$. Therefore, $U_2(F') = U_1(\{\text{id}\})$ is isomorphic to $\mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$ by Lemma 2.5. In particular, $U_2(\Gamma(A_3))$ and $U_2(t\Gamma(A_3)t^{-1})$ are not isomorphic.

Conversely, the following proposition based on [23, Appendix A], which states that in certain cases the tree can be recovered from the topological group structure of a subgroup of $\text{Aut}(T_d)$, applies to appropriate universal groups.

PROPOSITION 4.17. *Let $H, H' \leq \text{Aut}(T_d)$ be closed and locally transitive with distinct point-stabilizers. Then H and H' are isomorphic topological groups if and only if they are conjugate in $\text{Aut}(T_d)$.*

PROOF. By [8], every compact subgroup of H is either contained in a vertex stabilizer $H_x (x \in V)$ or, in case $H \not\leq \text{Aut}(T_d)^+$, in a geometric edge stabilizer $H_{\{e, \bar{e}\}} (e \in E)$. Since H is locally transitive, the above are pairwise distinct.

The vertex stabilizers are precisely those maximal compact subgroups $K \leq H$ for which there is no maximal compact subgroup K' with $[K : K \cap K'] = 2$. Indeed, for $e \in E$ and $x \in \{o(e), t(e)\}$ we have $[H_{\{e, \bar{e}\}} : H_{\{e, \bar{e}\}} \cap H_x] = 2$, whereas $[H_x : H_x \cap H_y] \geq 3$ and $[H_x : H_x \cap H_{\{e, \bar{e}\}}] \geq 3$ for all distinct $x, y \in V$ and $e \in E$ by the orbit–stabilizer theorem because $d \geq 3$ and H is locally transitive.

Adjacency can be expressed in terms of indices as well. Let $x, y \in V$ be distinct. Then $(x, y) \in E$ if and only if $[H_x : H_x \cap H_y] \leq [H_x : H_x \cap H_z]$ for all $z \in V$. Indeed, if $(x, y) \in E$, then $[H_x : H_x \cap H_y] = d$ by the orbit–stabilizer theorem, given that H is locally transitive. If $z \in V$ is not adjacent to x then $[H_x : H_x \cap H_z] > d$ because point-stabilizers of every local action of H are distinct.

Now, let $\Phi : H \rightarrow H'$ be an isomorphism of topological groups. Then Φ induces a bijection between the maximal compact subgroups of H and H' , and preserves indices. Hence, there is an automorphism $\varphi \in \text{Aut}(T_d)$ such that $\Phi(H_x) = H'_{\varphi(x)}$ for all $x \in V$. Furthermore, since vertex stabilizers in H' are pairwise distinct and

$$H'_{\varphi h \varphi^{-1}(x)} = \Phi(H_{h \varphi^{-1}(x)}) = \Phi(h H_{\varphi^{-1}(x)} h^{-1}) = \Phi(h) H'_x \Phi(h^{-1}) = H'_{\Phi(h)x}$$

for all $x \in V$, we have $\varphi h \varphi^{-1} = \Phi(h)$ for all $h \in H$. □

The following corollary uses the notation $\Phi^k(F')$ from Section 4.4.2.

COROLLARY 4.18. *Let $F \leq \text{Aut}(B_{d,k})$ and $F' \leq \text{Aut}(B_{d,k'})$ satisfy (C). Assume $k \geq k'$ and $\pi F, \pi F' \leq \text{Sym}(\Omega)$ are transitive with distinct point-stabilizers. If $U_k(F)$ and $U_{k'}(F')$ are isomorphic topological groups then $F, \Phi^k(F') \leq \text{Aut}(B_{d,k})$ are conjugate.*

PROOF. By Proposition 4.17, the groups $U_k(F)$ and $U_k(F')$ are conjugate in $\text{Aut}(T_d)$; hence so are $U_k(F)_x$ and $U_{k'}(F')_x$ for every $x \in V$, and the assertion follows. □

EXAMPLE 4.19. Section 4.4.1 introduces the isomorphic, nonconjugate subgroups $\Pi(S_3, \text{sgn}, \{1\})$ and $\Pi(S_3, \text{sgn}, \{0, 1\})$ of $\text{Aut}(B_{3,2})$, both of which project onto S_3 and satisfy (C) but not (D). An explicit isomorphism satisfies the assumption of Corollary 4.15. However, by Corollary 4.18 the universal groups $U_2(\Pi(S_3, \text{sgn}, \{1\}))$ and $U_2(\Pi(S_3, \text{sgn}, \{0, 1\}))$ are nonisomorphic. Therefore, Corollary 4.15 does not generalize to the nondiscrete case.

QUESTION 4.20. Let $F, F' \leq \text{Aut}(B_{d,k})$ satisfy (C) and be conjugate. Are the associated universal groups $U_k(F)$ and $U_k(F')$ necessarily isomorphic?

In the following, we determine the Burger–Mozes subquotient $H^{(\infty)}/\text{QZ}(H^{(\infty)})$ of Theorem 3.14 for nondiscrete, locally semiprimitive universal groups.

PROPOSITION 4.21. *Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C). If, in addition, F satisfies (D) then $\text{QZ}(U_k(F)) = U_k(F)$. Otherwise, $\text{QZ}(U_k(F)) = \{\text{id}\}$.*

PROOF. If F satisfies (D) then $U_k(F)$ is discrete and hence $\text{QZ}(U_k(F)) = U_k(F)$. Conversely, if F satisfies (C) but not (D) then the stabilizer of any half-tree $T \subseteq T_d$ in $U_k(F)$ is nontrivial: we have $T \in \{T_x, T_y\}$ for some edge $e := (x, y) \in E$. Since $U_k(F)$ is nondiscrete by Proposition 4.12 and has property (P_k) by Proposition 4.7, the group $U_k(F)_{e^k} = U_k(F)_{e^k, T_y} \cdot U_k(F)_{e^k, T_x}$ is nontrivial. In particular, either $U_k(F)_{T_x}$ or $U_k(F)_{T_y}$ is nontrivial. In view of the existence of label-respecting inversions, both are nontrivial and hence so is $U_k(F)_T$. Therefore, $U_k(F)$ has Property H of Möller–Vonk [18, Definition 2.3] and [18, Proposition 2.6] implies that $U_k(F)$ has trivial quasicentre. \square

PROPOSITION 4.22. *Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C) but not (D). Suppose that πF is semiprimitive. Then $U_k(F)^{(\infty)}/\text{QZ}(U_k(F)^{(\infty)}) = U_k(F)^{(\infty)} = U_k(F)^{+k}$.*

PROOF. The subgroup $U_k(F)^{+k} \leq U_k(F)$ is open, hence closed, and normal in $U_k(F)$ by definition. Since $U_k(F)$ is nondiscrete by Proposition 4.12, so is $U_k(F)^{+k}$. Using Proposition 3.11(iii), we conclude that $U_k(F)^{+k} \geq U_k(F)^{(\infty)}$. Since $U_k(F)$ satisfies property (P_k) by Proposition 4.7, the group $U_k(F)^{+k}$ is simple due to Theorem 2.1. Thus $U_k(F)^{+k} = U_k(F)^{(\infty)}$. Given that $\text{QZ}(U_k(F)^{(\infty)}) = \text{QZ}(U_k(F)) \cap U_k(F)^{(\infty)}$ by Proposition 3.11(iv), the assertion follows from Proposition 4.21. \square

In the context of Proposition 4.22, the group $U_k(F)^{+k}$ is simple, compactly generated, nondiscrete, totally disconnected, locally compact, second countable. Compact generation follows from [15, Corollary 2.11], given that $U_k(F)^{+k}$ is cocompact in $U_k(F)$ by Proposition 3.11(i).

4.4. Examples. We now construct various classes of examples of subgroups of $\text{Aut}(B_{d,k})$ satisfying (C) or (CD), and prove a rigidity result for certain local actions.

First, we give a suitable realization of $\text{Aut}(B_{d,k})$ and conditions (C) and (D). Namely, we view an automorphism α of $B_{d,k}$ as the set $\{\sigma_{k-1}(\alpha, v) \mid v \in B(b, 1)\}$ as follows. Let $\text{Aut}(B_{d,1}) \cong \text{Sym}(\Omega)$ be the natural isomorphism. For $k \geq 2$, we iteratively identify $\text{Aut}(B_{d,k})$ with its image under the map

$$\text{Aut}(B_{d,k}) \rightarrow \text{Aut}(B_{d,k-1}) \times \prod_{\omega \in \Omega} \text{Aut}(B_{d,k-1}), \quad \alpha \mapsto (\sigma_{k-1}(\alpha, b), (\sigma_{k-1}(\alpha, b_\omega))_\omega)$$

where $\text{Aut}(B_{d,k-1})$ acts on $\prod_{\omega \in \Omega} \text{Aut}(B_{d,k-1})$ by permuting the factors according to its action on $S(b, 1) \cong \Omega$. That is, multiplication in $\text{Aut}(B_{d,k})$ is given by

$$(\alpha, (\alpha_\omega)_{\omega \in \Omega}) \circ (\beta, (\beta_\omega)_{\omega \in \Omega}) = (\alpha\beta, (\alpha_{\beta\omega}\beta_\omega)_{\omega \in \Omega}).$$

Consider the homomorphism $\pi_{k-1} : \text{Aut}(B_{d,k}) \rightarrow \text{Aut}(B_{d,k-1})$, $\alpha \mapsto \sigma_{k-1}(\alpha, b)$, the projections $\text{pr}_\omega : \text{Aut}(B_{d,k}) \rightarrow \text{Aut}(B_{d,k-1})$, $\alpha \mapsto \sigma_{k-1}(\alpha, b_\omega)$ ($\omega \in \Omega$), and

$$p_\omega = (\pi_{k-1}, \text{pr}_\omega) : \text{Aut}(B_{d,k}) \rightarrow \text{Aut}(B_{d,k-1}) \times \text{Aut}(B_{d,k-1}),$$

whose image we interpret as a relation on $\text{Aut}(B_{d,k-1})$. Conditions (C) and (D) for a subgroup $F \leq \text{Aut}(B_{d,k})$ now read as follows:

$$\text{for all } \omega \in \Omega : p_\omega(F) \text{ is symmetric;} \tag{C}$$

$$\text{for all } \omega \in \Omega : p_\omega|_F^{-1}(\text{id}, \text{id}) = \{\text{id}\}. \tag{D}$$

4.4.1. The case $k = 2$. We first consider the case $k = 2$ which is all-encompassing in certain situations; see Theorem 4.32. By the above, $\text{Aut}(B_{d,2})$ is realized as follows: $\text{Aut}(B_{d,2}) = \{(a, (a_\omega)_{\omega \in \Omega}) \mid a \in \text{Sym}(\Omega), \text{ for all } \omega \in \Omega : a_\omega \in \text{Sym}(\Omega) \text{ and } a_\omega \omega = a\omega\}$.

Consider the map $\gamma : \text{Sym}(\Omega) \rightarrow \text{Aut}(B_{d,2})$, $a \mapsto (a, (a, \dots, a)) \in \text{Aut}(B_{d,2})$, using the realization of $\text{Aut}(B_{d,2})$ from above. For every $F \leq \text{Sym}(\Omega)$, the image

$$\Gamma(F) := \text{im}(\gamma|_F) = \{(a, (a, \dots, a)) \mid a \in F\} \cong F$$

is a subgroup of $\text{Aut}(B_{d,2})$ which is isomorphic to F and satisfies both (C) and (D). The involutive compatibility cocycle is given by $\Gamma(F) \times \Omega \rightarrow \Gamma(F)$, $(\gamma(a), \omega) \mapsto \gamma(a)$. Note that $\Gamma(F) \cong F$ implements the diagonal action $F \curvearrowright \Omega^2$ on $S(b, 2) \cong \Omega^{(2)} \subset \Omega^2$.

We obtain $U_2(\Gamma(F)) = \{\alpha \in \text{Aut}(T_d) \mid \text{there exists } a \in F : \text{for all } x \in V : \sigma_1(\alpha, x) = a\} =: D(F)$, following the notation of [1]. Moreover, there is the following description of all subgroups $\tilde{F} \leq \text{Aut}(B_{d,2})$ with $\pi\tilde{F} = F$ that satisfy (C) and contain $\Gamma(F)$.

PROPOSITION 4.23. *Let $F \leq \text{Sym}(\Omega)$. Given $K \leq \prod_{\omega \in \Omega} F_\omega \cong \ker \pi \leq \text{Aut}(B_{d,2})$, there is $\tilde{F} \leq \text{Aut}(B_{d,2})$ satisfying (C) and fitting into the split exact sequence*

$$1 \longrightarrow K \xrightarrow{\iota} \tilde{F} \xrightleftharpoons[\gamma]{\pi} F \longrightarrow 1$$

if and only if K is preserved by the action $F \curvearrowright \prod_{\omega \in \Omega} F_\omega$, $a \cdot (a_\omega)_\omega := (aa_{a^{-1}\omega}a^{-1})_\omega$.

PROOF. If there is a split exact sequence as above then $K \trianglelefteq \tilde{F}$ is invariant under conjugation by $\Gamma(F) \leq \tilde{F}$, hence the assertion.

Conversely, if K is invariant under the given action, then

$$\tilde{F} := \{(a, (aa_\omega)_\omega) \mid a \in F, (a_\omega)_\omega \in K\}$$

fits into the sequence. First, note that \tilde{F} contains both K and $\Gamma(F)$. It is also a subgroup of $\text{Aut}(B_{d,2})$: for $(a, (aa_\omega)_\omega), (b, (bb_\omega)_\omega) \in \tilde{F}$ we have

$$(a, (aa_\omega)_\omega) \circ (b, (bb_\omega)_\omega) = (ab, (aa_{b\omega}bb_\omega)_\omega) = (ab, (ab \circ b^{-1}a_{b\omega}b \circ b_\omega)_\omega) \in \tilde{F}$$

by assumption. In particular, $\tilde{F} = \langle \Gamma(F), K \rangle$. It suffices to check condition (C) on these generators of \tilde{F} . As before, $\gamma(a) \in C(\gamma(a), \omega)$ for all $a \in F$ and $\omega \in \Omega$. Now let $k \in K$. Then $\gamma(\text{pr}_\omega k)k^{-1} \in C(k, \omega)$ for all $\omega \in \Omega$. □

EXAMPLE 4.24. We show that for certain dihedral groups there are only four groups of the type given in Proposition 4.23. Set $F := D_p \leq \text{Sym}(p)$ for some prime $p \geq 3$. Then $F_\omega \cong (\mathbb{F}_2, +)$. Hence, $U := \prod_{\omega \in \Omega} F_\omega$ is a p -dimensional vector space over \mathbb{F}_2 and the F -action on it permutes coordinates. When $2 \in (\mathbb{Z}/p\mathbb{Z})^*$ is primitive, there are only four F -invariant subspaces of U : the trivial subspace; the diagonal subspace $\langle(1, \dots, 1)\rangle$; the whole space; and $K := \ker \sigma \cong \mathbb{F}_2^{(p-1)}$ where $\sigma: U \rightarrow \mathbb{F}_2$ is given by $(v_1, \dots, v_p) \mapsto \sum_{i=1}^p v_i$. Note that K is F -invariant because the homomorphism σ is. Conjecturally, there are infinitely many primes for which $2 \in (\mathbb{Z}/p\mathbb{Z})^*$ is primitive. The list starts with 3, 5, 11, 13, ...; see [26, A001122].

Suppose that $W \leq U$ is F -invariant. It suffices to show that W contains K as soon as $W \cap \ker \sigma$ contains a nontrivial element w . To see this, we show that the orbit of w under the cyclic group $\langle \varrho \rangle = C_p \leq D_p$ generates a $(p - 1)$ -dimensional subspace of K which hence equals K . Indeed, the rank of the circulant matrix $C := (w, \varrho w, \varrho^2 w, \dots, \varrho^{(p-1)} w)$ equals $p - \deg(\gcd(x^p - 1, f(x)))$ where $f(x) \in \mathbb{F}_2[x]$ is the polynomial $f(x) = w_p x^{p-1} + \dots + w_2 x + w_1$; see, for example, [5, Corollary 1]. The polynomial $x^p - 1 \in \mathbb{F}_2[x]$ factors into the irreducibles $(x^{p-1} + x^{p-2} + \dots + x + 1)(x - 1)$ by the assumption on p . Since f has an even number of nonzero coefficients, we conclude that $\text{rank}(C) = p - 1$.

The following subgroups of $\text{Aut}(B_{d,2})$ are of the type given in Proposition 4.23. Let $F \leq \text{Sym}(\Omega)$ be transitive. Fix $\omega_0 \in \Omega$, let $C \leq Z(F_{\omega_0})$ and let $N \trianglelefteq F_{\omega_0}$ be normal. Furthermore, fix elements $f_\omega \in F$ ($\omega \in \Omega$) satisfying $f_\omega(\omega_0) = \omega$. We define

$$\Delta(F, C) := \{(a, (a \circ f_\omega a_0 f_\omega^{-1})_\omega) \mid a \in F, a_0 \in C\} \cong F \times C,$$

$$\Phi(F, N) := \{(a, (a \circ f_\omega a_0^{(\omega)} f_\omega^{-1})_\omega) \mid a \in F, \text{ for all } \omega \in \Omega : a_0^{(\omega)} \in N\} \cong F \ltimes N^d.$$

In the case of $\Delta(F, C)$ we have $K = \{(f_\omega a_0 f_\omega^{-1})_\omega \mid a_0 \in C\}$, whereas in the case of $\Phi(F, N)$ we have $K = \{(f_\omega a_0^{(\omega)} f_\omega^{-1})_\omega \mid \text{for all } \omega \in \Omega : a_0^{(\omega)} \in N\}$. In both cases, invariance under the action of F is readily verified, as is condition (D) for $\Delta(F, C)$.

The group $\Delta(F, F_{\omega_0})$ can be defined for nonabelian F_{ω_0} as well, namely,

$$\Delta(F) := \{(a, (f_{a\omega} f_\omega^{-1} \circ f_\omega a_0 f_\omega^{-1})_\omega) \mid a \in F, a_0 \in F_{\omega_0}\} \cong F \times F_{\omega_0}.$$

However, it need not contain $\Gamma(F)$. Note that $\Phi(F, N)$ does not depend on the choice of the elements $(f_\omega)_{\omega \in \Omega}$ as N is normal in F_{ω_0} , whereas $\Delta(F, C)$ and $\Delta(F)$ may. However, any group of the form $\{(a, (z(a, \omega) \alpha_\omega(a_0))_\omega) \mid a \in F, a_0 \in F_{\omega_0}\}$, where z is a compatibility cocycle of F and $\alpha_\omega : F_{\omega_0} \rightarrow F_\omega$ ($\omega \in \Omega$) are isomorphisms, that satisfies (C) and in which $\{(a, (z(a, \omega))_\omega) \mid a \in F\}$ and $\{(\text{id}, (\alpha_\omega(a_0))_\omega) \mid a_0 \in F_{\omega_0}\}$ commute, will be referred to as $\Delta(F)$ in view of Corollary 4.15.

The group $\Phi(F, F_{\omega_0})$ can be defined without assuming transitivity of F , namely,

$$\Phi(F) := \{(a, (a_\omega)_\omega) \mid a \in F, \text{ for all } \omega \in \Omega : a_\omega \in C_F(a, \omega)\} \cong F \times \prod_{\omega \in \Omega} F_\omega.$$

We conclude that $U_2(\Phi(F)) = U_1(F)$ for every $F \leq \text{Sym}(\Omega)$.

When $F \leq \text{Sym}(\Omega)$ preserves a partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω , we define

$$\Phi(F, \mathcal{P}) := \{(a, (a_\omega)_\omega) \mid a \in F, a_\omega \in C_F(a, \omega) \text{ constant w.r.t. } \mathcal{P}\} \cong F \times \prod_{i \in I} F_{\Omega_i}.$$

The group $\Phi(F, \mathcal{P})$ satisfies (C) as well and features prominently in Section 5.1.

The following kind of 2-local action generalizes the sign construction in [23]. Let $F \leq \text{Sym}(\Omega)$ and let $\rho : F \rightarrow A$ be a homomorphism to an abelian group A . Define

$$\begin{aligned} \Pi(F, \rho, \{1\}) &:= \left\{ (a, (a_\omega)_\omega) \in \Phi(F) \mid \prod_{\omega \in \Omega} \rho(a_\omega) = 1 \right\}, \\ \Pi(F, \rho, \{0, 1\}) &:= \left\{ (a, (a_\omega)_\omega) \in \Phi(F) \mid \rho(a) \prod_{\omega \in \Omega} \rho(a_\omega) = 1 \right\}. \end{aligned}$$

This construction is generalized to $k \geq 2$ in Section 4.4.2 where the third entry of Π is a set of radii over which the defining product is taken.

PROPOSITION 4.25. *Let $F \leq \text{Sym}(\Omega)$ and let $\rho : F \rightarrow A$ be a homomorphism to an abelian group A . Let $\widetilde{F} \in \{\Pi(F, \rho, \{1\}), \Pi(F, \rho, \{0, 1\})\}$. If $\rho(F_\omega) = A$ for all $\omega \in \Omega$ then $\pi\widetilde{F} = F$ and \widetilde{F} satisfies (C).*

PROOF. As $C_F(a, \omega) = aF_\omega$, and $\rho(F_\omega) = A$ for all $\omega \in \Omega$, an element $(a, (a_\omega)_\omega) \in \Phi(F)$ can be turned into an element of \widetilde{F} by changing a_ω for a single, arbitrary $\omega \in \Omega$. We conclude that $\pi\widetilde{F} = F$ and that \widetilde{F} satisfies (C). □

4.4.2. General case. We extend some constructions of Section 4.4.1 to arbitrary k . Given $F \leq \text{Aut}(B_{d,k})$ satisfying (C), define the subgroup

$$\Phi_k(F) := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in F, \text{ for all } \omega \in \Omega : \alpha_\omega \in C_F(\alpha, \omega)\} \leq \text{Aut}(B_{d,k+1}).$$

Then $\Phi_k(F)$ inherits condition (C) from F and we obtain $U_{k+1}(\Phi_k(F)) = U_k(F)$. Concerning the construction Γ we have the following proposition.

PROPOSITION 4.26. *Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C). Then there exists a group $\Gamma_k(F) \leq \text{Aut}(B_{d,k+1})$ satisfying (CD) such that $\pi_k|_{\Gamma_k(F)}$ is an isomorphism onto F if and only if F admits an involutive compatibility cocycle z .*

PROOF. If F admits an involutive compatibility cocycle z , define

$$\Gamma_k(F) := \{(\alpha, (z(\alpha, \omega))_\omega) \mid \alpha \in F\} \leq \text{Aut}(B_{d,k+1}).$$

Then $\gamma_z : F \rightarrow \Gamma_k(F)$, $\alpha \mapsto (\alpha, (z(\alpha, \omega))_\omega)$ is an isomorphism and the involutive compatibility cocycle of $\Gamma_k(F)$ is given by $\widetilde{z} : (\gamma_z(\alpha), \omega) \mapsto \gamma_z(z(\alpha, \omega))$. Conversely, if a group $\Gamma_k(F)$ with the asserted properties exists, set $z : (\alpha, \omega) \mapsto \text{pr}_\omega \pi_k^{-1} \alpha$. □

Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C) and let $l > k$. We set $\Gamma^l(F) := \Gamma_{l-1} \circ \dots \circ \Gamma_k(F)$ for an implicit sequence of involutive compatibility cocycles. Similarly, we define $\Phi^l(F) := \Phi_{l-1} \circ \dots \circ \Phi_k(F)$. Now, let $\widetilde{F} \leq \text{Aut}(B_{d,k})$. Assume $F := \pi\widetilde{F} \leq \text{Sym}(\Omega)$ preserves a

partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω . Define the group

$$\Phi_k(\widetilde{F}, \mathcal{P}) := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in \widetilde{F}, \alpha_\omega \in C_{\widetilde{F}}(\alpha, \omega) \text{ is constant w.r.t. } \mathcal{P}\}.$$

If $C_{\widetilde{F}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in \widetilde{F}$ and $i \in I$ then $\Phi_k(\widetilde{F}, \mathcal{P})$ satisfies (C), and if $C_{\widetilde{F}}(\text{id}, \Omega_i)$ is nontrivial for all $i \in I$ then $\Phi_k(\widetilde{F}, \mathcal{P})$ does not satisfy (D).

The following statement generalizes Proposition 4.23.

PROPOSITION 4.27. *Let $F \leq \text{Aut}(B_{d,k})$ satisfy (C). Suppose F admits an involutive compatibility cocycle z . Given $K \leq \Phi_k(F) \cap \ker(\pi_k)$, there is $\widetilde{F} \leq \text{Aut}(B_{d,k+1})$ satisfying (C) and fitting into the split exact sequence*

$$1 \longrightarrow K \xrightarrow{\iota} \widetilde{F} \xrightarrow[\gamma_z]{\pi} F \longrightarrow 1$$

if and only if $\Gamma_k(F)$ normalizes K , and for all $k \in K$ and $\omega \in \Omega$ there is $k_\omega \in K$ such that $\text{pr}_\omega k_\omega = z(\text{pr}_\omega k, \omega)^{-1}$.

PROOF. If there is a split exact sequence as above then $K \leq \widetilde{F}$ is invariant under conjugation by $\Gamma_k(F)$. Moreover, all elements of \widetilde{F} have the form $(\alpha, (z(\alpha, \omega)\alpha_\omega)_\omega)$ for some $\alpha \in F$ and $(\alpha_\omega)_\omega \in K$. This implies the second assertion on K .

Conversely, if K satisfies the assumptions, then

$$\widetilde{F} := \{(\alpha, (z(\alpha, \omega)\alpha_\omega)_\omega) \mid \alpha \in F, (\alpha_\omega)_\omega \in K\}$$

fits into the sequence. First, note that \widetilde{F} contains both K and $\Gamma_k(F)$. It is also a subgroup of $\text{Aut}(B_{d,k+1})$: for $(\alpha, (z(\alpha, \omega)\alpha_\omega)_\omega), (\beta, (z(\beta, \omega)\beta_\omega)_\omega) \in \widetilde{F}$ we have

$$\begin{aligned} (\alpha, (z(\alpha, \omega)\alpha_\omega)_\omega) \circ (\beta, (z(\beta, \omega)\beta_\omega)_\omega) &= (\alpha\beta, (z(\alpha, \beta\omega)\alpha_{\beta\omega}z(\beta, \omega)\beta_\omega)_\omega) \\ &= (\alpha\beta, (z(\alpha, \beta\omega)z(\beta, \omega) \circ z(\beta, \omega)^{-1}\alpha_{\beta\omega}z(\beta, \omega) \circ \beta_\omega)_\omega) \\ &= (\alpha\beta, (z(\alpha\beta, \omega)\alpha'_\omega\beta_\omega)_\omega) \in \widetilde{F} \end{aligned}$$

for some $(\alpha'_\omega)_\omega \in K$ because $\Gamma_k(F)$ normalizes K . In particular, $\widetilde{F} = \langle \Gamma_k(F), K \rangle$. We check condition (C) on these generators. As before, $\gamma_z(z(\alpha, \omega)) \in C(\gamma_z(\alpha), \omega)$ for all $\alpha \in F$ and $\omega \in \Omega$ because z is involutive. Now, let $k \in K$. We then have $\gamma_z(\text{pr}_\omega k)k_\omega \in C(k, \omega)$ for all $\omega \in \Omega$ by the assumption on k_ω . \square

In the split situation of Proposition 4.27 we also denote \widetilde{F} by $\Sigma_k(F, K)$. For instance, the group $\Pi(S_3, \text{sgn}, \{1\})$ of Proposition 4.25 satisfies (C), admits an involutive compatibility cocycle but does not satisfy (D); see Section 5.3.

Now, let $F \leq \text{Sym}(\Omega)$ and $\rho : F \rightarrow A$ a homomorphism to an abelian group A . Further, let $k \in \mathbb{N}$ and $X \subseteq \{0, \dots, k - 1\}$. Define

$$\Pi^k(F, \rho, X) := \left\{ \alpha \in \Phi^k(F) \mid \prod_{r \in X} \prod_{x \in S(b,r)} \rho(\sigma_1(\alpha, x)) = 1 \right\}.$$

PROPOSITION 4.28. *Let $F \leq \text{Sym}(\Omega)$ and let $\rho : F \rightarrow A$ be a homomorphism to an abelian group A . Further, let $k \in \mathbb{N}$ and $X \subseteq \{0, \dots, k - 1\}$ be nonempty and nonzero with $k - 1 \in X$. If $\rho(F_\omega) = A$ for all $\omega \in \Omega$ then $\pi(\Pi^k(F, \rho, X)) = F$ and $\Pi^k(F, \rho, X)$ has (C).*

PROOF. As $C_F(a, \omega) = aF_\omega$, and $\rho(F_\omega) = A$ for all $\omega \in \Omega$, an element $\alpha \in \Phi^k(F)$ can be turned into an element of $\Pi^k(F, \rho, X)$ by changing $\sigma_1(\alpha, x)$ for a single, arbitrary $x \in S(b, k - 1)$. When X is nonzero we conclude that $\pi(\Pi^k(F, \rho, X)) = F$ and that $\Pi^k(F, \rho, X)$ satisfies (C). □

4.4.3. A rigid case. For certain $F \leq \text{Sym}(\Omega)$ the groups $\Gamma(F)$, $\Delta(F)$ and $\Phi(F)$ already yield all possible $U_k(\tilde{F})$ with $\pi\tilde{F} = F$. The main argument is based on Sections 3.4 and 3.5 of [2]. We first record the following lemma whose proof is due to M. Giudici by personal communication.

LEMMA 4.29. *Let $F \leq \text{Sym}(\Omega)$ be 2-transitive and F_ω ($\omega \in \Omega$) simple nonabelian. Then every extension \tilde{F} of F_ω ($\omega \in \Omega$) by F is equivalent to $F_\omega \times F$.*

PROOF. Regarding F_ω as a normal subgroup of \tilde{F} , consider the conjugation map $\varphi : \tilde{F} \rightarrow \text{Aut}(F_\omega)$. We show that $K := \ker \varphi = Z_{\tilde{F}}(F_\omega) \trianglelefteq \tilde{F}$ complements F_ω in \tilde{F} . Since $Z(F_\omega) = \{\text{id}\}$, we have $F_\omega \cap K = \{\text{id}\}$. Hence, $F_\omega K \trianglelefteq \tilde{F}$. Next, consider $\tilde{F}/(F_\omega K) \cong \text{Out}(F_\omega)$. By the solution of Schreier’s conjecture, $\text{Out}(F_\omega)$ is solvable. Since $\tilde{F}/F_\omega \cong F$ is not solvable we conclude $K \neq \{\text{id}\}$. Now, by a theorem of Burnside, every 2-transitive permutation group F is either almost simple or of affine type; see [6, Theorem 4.1B and Section 4.8].

In the first case, F is actually simple: Let $N \trianglelefteq F$. Then $F_\omega \cap N \trianglelefteq F_\omega$. Hence, either $F_\omega \cap N = \{\text{id}\}$ or $F_\omega \cap N = F_\omega$. Since F is 2-transitive and therefore primitive, every normal subgroup acts transitively. Hence, in the first case, N is regular, which contradicts F being almost simple. Thus the second case holds and $N = NF_\omega = F$. Now $\tilde{F}/F_\omega K$ is a proper quotient of F and therefore trivial. We conclude that $\tilde{F} = F_\omega K \cong F_\omega \times K$ and $K \cong \tilde{F}/F_\omega \cong F$.

In the second case, $F = F_\omega \rtimes C_p^d$ for some $d \in \mathbb{N}$ and prime p . Given that K is nontrivial and $K \cong F_\omega K/F_\omega \trianglelefteq F$, it contains the unique minimal normal subgroup $C_p^d \trianglelefteq K \trianglelefteq F$. Since $F/C_p^d \cong F_\omega$ is nonabelian simple whereas the proper quotient $\tilde{F}/F_\omega K$ of F is solvable, $K \neq C_p^d$. But $F/C_p^d \cong F_\omega$ is simple, so $F_\omega K = F$. □

The following propositions are of independent interest and used in Theorem 4.32 below. We introduce the following notation. Let $\tilde{F} \leq \text{Aut}(B_{d,k})$ and $K \leq \tilde{F}_{b_\xi}$ for some $\xi = (\omega_1, \dots, \omega_{k-1}) \in \Omega^{(k-1)}$. We set $\pi_\xi K := \sigma_1(K, b_\xi) \leq \text{Sym}(\Omega)_{\omega_{k-1}}$.

PROPOSITION 4.30. *Let $\tilde{F} \leq \text{Aut}(B_{d,k})$ satisfy (C). Suppose $F := \pi\tilde{F}$ is transitive. Further, let $\omega \in \Omega$ and $\xi = (\omega_1, \dots, \omega_{k-1}) \in \Omega^{(k-1)}$ with $\omega_1 \neq \omega$. Then $\pi_\xi(\tilde{F}_{b_\xi} \cap \ker \pi)$ and $\pi_\xi \tilde{F}_{T_\omega}$ are subnormal in $F_{\omega_{k-1}}$ of depth at most $k - 1$ and k , respectively.*

PROOF. We argue by induction on $k \geq 2$. For $k = 2$, the assertion that $\pi_\xi(\tilde{F}_{b_\xi} \cap \ker \pi)$ is normal in F_{ω_1} is a consequence of condition (C). Now, suppose $\tilde{F} \leq \text{Aut}(B_{d,k+1})$ satisfies the assumptions, and let $\omega \in \Omega$ and $\xi = (\omega_1, \dots, \omega_k) \in \Omega^{(k)}$ be such that $\omega_1 \neq \omega$. Since \tilde{F} satisfies (C), we have $\text{pr}_{\omega_1}(\tilde{F}_{b_\xi} \cap \ker \pi) \trianglelefteq (\pi_k \tilde{F})_{b_{\xi'}} \cap \ker \pi$, where $\xi' := (\omega_2, \dots, \omega_{k-1})$ and the right-hand-side π implicitly has domain $\pi_k \tilde{F}$. Hence,

$$\pi_\xi(\tilde{F}_{b_\xi} \cap \ker \pi) = \pi_{\xi'}(\text{pr}_{\omega_1}(\tilde{F}_{b_\xi} \cap \ker \pi)) \trianglelefteq \pi_{\xi'}((\pi_k \tilde{F})_{b_{\xi'}} \cap \ker \pi) \trianglelefteq F_{\omega_{k-1}}$$

by the induction hypothesis. The second assertion follows as $\tilde{F}_{T_\omega} \trianglelefteq \tilde{F}_{b_\xi} \cap \ker \pi$. □

PROPOSITION 4.31. *Let $\widetilde{F} \leq \text{Aut}(B_{d,k})$ satisfy (C) but not (D). Suppose $F := \pi\widetilde{F}$ is transitive, and every nontrivial subnormal subgroup of F_ω ($\omega \in \Omega$) of depth at most $k - 1$ is transitive on $\Omega \setminus \{\omega\}$. Then $U_k(\widetilde{F})$ is locally k -transitive.*

PROOF. We argue by induction on k . For $k = 1$, the assertion follows from transitivity of F . Now, let $\widetilde{F} \leq \text{Aut}(B_{d,k+1})$ satisfy (C) but not (D). Then the same holds for $F^{(k)} := \pi_k \widetilde{F} \leq \text{Aut}(B_{d,k})$. Given $\widetilde{\xi}, \widetilde{\xi}' \in \Omega^{(k)}$, write $\widetilde{\xi} = (\xi, \omega)$ and $\widetilde{\xi}' = (\xi', \omega')$ where $\xi, \xi' \in \Omega^{(k-1)}$ and $\omega, \omega' \in \Omega$. By the induction hypothesis, the group $F^{(k)}$ acts transitively on $S(b, k)$. Hence, using (C), there is $g \in \widetilde{F}$ such that $g b_{\widetilde{\xi}} = b_{\widetilde{\xi}'}$. As \widetilde{F} does not satisfy (D) the said transitivity further implies that $\pi_{\widetilde{\xi}'}(\widetilde{F}_{b_{\widetilde{\xi}'}} \cap \ker \pi)$ is nontrivial. By Proposition 4.30, it is also subnormal of depth at most $k - 1$ in $F_{\omega'}$ and thus transitive. Hence, there is $g' \in \widetilde{F}_{b_{\widetilde{\xi}'}}$ with $g' g b_{\widetilde{\xi}} = b_{\widetilde{\xi}'}$. \square

The following theorem is closely related to [2, Proposition 3.3.1].

THEOREM 4.32. *Let $F \leq \text{Sym}(\Omega)$ be 2-transitive and F_ω ($\omega \in \Omega$) simple nonabelian. Further, let $\widetilde{F} \leq \text{Aut}(B_{d,k})$ with $\pi\widetilde{F} = F$ satisfy (C). Then $U_k(\widetilde{F})$ equals*

$$U_2(\Gamma_1(F)), \quad U_2(\Delta(F)) \quad \text{or} \quad U_2(\Phi(F)) = U_1(F).$$

PROOF. Since $U_1(F) = U_2(\Phi(F))$, we may assume $k \geq 2$. Given that $\widetilde{F} \leq \text{Aut}(B_{d,k})$ satisfies (C), so does the restriction $F^{(2)} := \pi_2 \widetilde{F} \leq \Phi(F) \leq \text{Aut}(B_{d,2})$. Consider the projection $\pi : F^{(2)} \rightarrow F$. We have $\ker \pi \leq \prod_{\omega \in \Omega} F_\omega$ and $\text{pr}_\omega \ker \pi \trianglelefteq F_\omega$ for all $\omega \in \Omega$ by Proposition 4.30. Since F_ω is simple, $\ker \pi \trianglelefteq F^{(2)}$ and F is transitive, this implies that either $\text{pr}_\omega \ker \pi = \{\text{id}\}$ for all $\omega \in \Omega$ or $\text{pr}_\omega \ker \pi = F_\omega$ for all $\omega \in \Omega$.

In the first case, $\pi : F^{(2)} \rightarrow F$ is an isomorphism. Hence, $F^{(2)}$ satisfies (CD) and $U_k(\widetilde{F}) = U_2(\Gamma_1(F))$ for an involutive compatibility cocycle of F by Proposition 4.26.

In the second case, fix $\omega_0 \in \Omega$. We have $\ker \pi \leq \prod_{\omega \in \Omega} F_\omega \cong F_{\omega_0}^d$ by transitivity of F . Since F_{ω_0} is simple nonabelian, [23, Lemma 2.3] implies that the group $\ker \pi$ is a product of subdiagonals preserved by the primitive action of F on the index set of $F_{\omega_0}^d$. Hence, either there is just one block and $\ker \pi = \{(\text{id}, (\alpha_\omega(a_0))_\omega)\}$ for some isomorphisms $\alpha_\omega : F_{\omega_0} \rightarrow F_\omega$, or all blocks are singletons and $\ker \pi = \prod_{\omega \in \Omega} F_\omega \cong F_{\omega_0}^d$. In the first case, there is a compatibility cocycle z of F such that $F \cong \{(a, (z(a, \omega))_\omega) \mid a \in F\} \leq F^{(2)}$ commutes with $\ker \pi \leq F^{(2)}$ by Lemma 4.29. Thus $F^{(2)} = \{a, (z(a, \omega)\alpha_\omega(a_0))_\omega \mid a \in F, a_0 \in F_{\omega_0}\}$. In particular, $F^{(2)}$ satisfies (CD). Hence, $U_k(\widetilde{F}) = U_2(\Delta(F))$.

When $\ker \pi \cong F_{\omega_0}^d$, we have $U_k(\widetilde{F}) = U_1(F)$ by [2, Proposition 3.3.1]. \square

If F does not have simple point-stabilizers or preserves a nontrivial partition, more universal groups are given by $U_2(\Phi(F, N))$ and $U_2(\Phi(F, \mathcal{P}))$; see Section 4.4.1. When F is 2-transitive and has abelian point-stabilizers, $F \cong \text{AGL}(1, q)$ for some prime power q by [14]. Hence, point-stabilizers in F are isomorphic to \mathbb{F}_q^* and simple if and only if $q - 1$ is a Mersenne prime. For any value of q , the projection $\rho : \text{AGL}(1, q) \rightarrow \mathbb{F}_q^*$ satisfies the assumptions of Proposition 4.28 and so the groups $U_k(\Pi^k(\text{AGL}(1, q), \rho, X))$ provide further examples. The following question remains.

QUESTION 4.33. Let $F \leq \text{Sym}(\Omega)$ be primitive and $F_\omega (\omega \in \Omega)$ simple nonabelian. Is there $\bar{F} \leq \text{Aut}(B_{d,k})$ with (C) and $\pi\bar{F} = F$ other than $\Gamma^k(F)$, $\Delta(F)$ and $\Phi^k(F)$?

4.5. Universality. The constructed groups $U_k(F)$ are universal in the sense of the following maximality statement, which should be compared to Proposition 2.6.

THEOREM 4.34. Let $H \leq \text{Aut}(T_d)$ be locally transitive and contain an involutive inversion. Then there is a labelling l of T_d such that

$$U_1^{(l)}(F^{(1)}) \geq U_2^{(l)}(F^{(2)}) \geq \dots \geq U_k^{(l)}(F^{(k)}) \geq \dots \geq H \geq U_1^{(l)}(\{\text{id}\})$$

where $F^{(k)} \leq \text{Aut}(B_{d,k})$ is action isomorphic to the k -local action of H .

PROOF. First, we construct a labelling l of T_d such that $H \geq U_1^{(l)}(\{\text{id}\})$. Fix $x \in V$ and choose a bijection $l_x : E(x) \rightarrow \Omega$. By the assumptions, there is an involutive inversion $\iota_\omega \in H$ of the edge $(x, x_\omega) \in E$ for every $\omega \in \Omega$. Using these inversions, we define the announced labelling inductively. Set $l_{E(x)} := l_x$ and assume that l is defined on $E(x, n)$. For $e \in E(x, n+1) \setminus E(x, n)$ put $l(e) := l(\iota_\omega(e))$ if x_ω is part of the unique reduced path from x to $o(e)$. Since the ι_ω ($\omega \in \Omega$) have order 2, we obtain $\sigma_1(\iota_\omega, y) = \text{id}$ for all $\omega \in \Omega$ and $y \in V$. Therefore, $\langle \{\iota_\omega \mid \omega \in \Omega\} \rangle = U_1^{(l)}(\{\text{id}\}) \leq H$, following the proof of Lemma 2.5.

Now, let $h \in H$ and $y \in V$. Further, let (x, x_1, \dots, x_n, y) and $(x, x'_1, \dots, x'_m, h(y))$ be the unique reduced paths from x to y and $h(y)$, respectively. Since $U_1^{(l)}(\{\text{id}\}) \leq H$, the group H contains the unique label-respecting inversion ι_e of every edge $e \in E$. We therefore have

$$s := \iota_{(x'_1, x)}^{-1} \cdots \iota_{(x'_m, x'_{m-1})}^{-1} \iota_{(h(y), x'_m)}^{-1} \circ h \circ \iota_{(y, x_n)} \cdots \iota_{(x_2, x_1)} \iota_{(x_1, x)} \in H.$$

Also, s stabilizes x . The cocycle identity implies for every $k \in \mathbb{N}$ that

$$\sigma_k(h, y) = \sigma_k(\iota_{(h(y), x'_m)} \cdots \iota_{(x'_1, x)} \circ s \circ \iota_{(x_1, x)}^{-1} \cdots \iota_{(y, x_n)}^{-1}, y) = \sigma_k(s, x) \in F^{(k)},$$

where $F^{(k)} \leq \text{Aut}(B_{d,k})$ is defined by $l_x^k \circ H_x|_{B(x,k)} \circ (l_x^k)^{-1}$. □

REMARK 4.35. Retain the notation of Theorem 4.34. By Proposition 2.6, there is a labelling l of T_d such that $U_1^{(l)}(F^{(1)}) \geq H$ regardless of the minimal order of an inversion in H . This labelling may be distinct from that of Theorem 4.34 which fails without assuming the existence of an involutive inversion. For example, a vertex-stabilizer of the group G_2^1 of Example 5.39 below is action isomorphic to $\Gamma(S_3)$ but $G_2^1 \not\geq U_2^{(l)}(\Gamma(S_3))$ for any labelling l because $(G_2^1)_{(b, b_\omega)} \cong \mathbb{Z}/4\mathbb{Z}$, whereas

$$U_2^{(l)}(\Gamma(S_3))_{(b, b_\omega)} \cong \Gamma(S_3)_{(b, b_\omega)} \rtimes \mathbb{Z}/2\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$$

by Proposition 4.14.

We complement Theorem 4.34 with the following criterion for certain subgroups of $\text{Aut}(T_d)$ to contain an involutive inversion.

PROPOSITION 4.36. Let $H \leq \text{Aut}(T_d)$ be locally transitive with odd-order point-stabilizers. If H contains a finite-order inversion then it contains an involutive one.

PROOF. Let $\iota \in H$ be a finite-order inversion of an edge $e \in E$ and $\text{ord}(\iota) = 2^k \cdot m$ for some odd $m \in \mathbb{N}$ and some $k \in \mathbb{N}$. It suffices to show that $k = 1$, in which case ι^m is an involutive inversion. Suppose $k \geq 1$. Then $\iota^{2^{k-1} \cdot m}$ is nontrivial and fixes the edge e . Because point-stabilizers in the local action of H have odd order, it follows that $(\iota^{2^{k-1} \cdot m})^2$ is nontrivial as well, but $(\iota^{2^{k-1} \cdot m})^2 = \iota^{\text{ord}(\iota)} = \text{id}$. \square

For example, Proposition 4.36 applies when H is discrete and vertex-transitive: Combined with local transitivity this implies the existence of a finite-order inversion.

We remark that primitive permutation groups with odd-order point-stabilizers were classified in [17]. For instance, they include $\text{PSL}(2, q) \curvearrowright \text{P}^1(\mathbb{F}_q)$ for any prime power q that satisfies $q \equiv 3 \pmod 4$.

4.6. A bipartite version. In this section we introduce a bipartite version of the universal groups developed above which plays a critical role in the proof of Theorem 5.2(iv)(b). As before, let $T_d = (V, E)$ denote the d -regular tree. Fix a regular bipartition $V = V_1 \sqcup V_2$ of T_d .

4.6.1. Definition and basic properties. The groups to be defined are subgroups of ${}^+\text{Aut}(T_d) \leq \text{Aut}(T_d)$, the maximal subgroup of $\text{Aut}(T_d)$ preserving the bipartition $V = V_1 \sqcup V_2$. Alternatively, it can be described as the subgroup generated by all point-stabilizers, or all edge-stabilizers.

DEFINITION 4.37. Let $F \leq \text{Aut}(B_{d,2k})$ and l be a labelling of T_d . Define

$$\text{BU}_{2k}^{(l)}(F) := \{\alpha \in {}^+\text{Aut}(T_d) \mid \text{for all } v \in V_1 : \sigma_{2k}(\alpha, v) \in F\}.$$

Note that $\text{BU}_{2k}^{(l)}(F)$ is a subgroup of ${}^+\text{Aut}(T_d)$ thanks to Lemma 4.2 and the assumption that it is a subset of ${}^+\text{Aut}(T_d)$. Further, Proposition 4.4 carries over to the groups $\text{BU}_{2k}^{(l)}(F)$. We therefore omit the reference to an explicit labelling in the following. Also, we recover the following basic properties.

PROPOSITION 4.38. Let $F \leq \text{Aut}(B_{d,2k})$. The group $\text{BU}_{2k}(F)$ is

- (i) *closed in $\text{Aut}(T_d)$*
- (ii) *transitive on both V_1 and V_2 , and*
- (iii) *compactly generated.*

Parts (i) and (ii) are proven as their analogues in Proposition 4.5, whereas part (iii) relies on part (ii) and the subsequent analogue of Lemma 2.5, for which we introduce the following notation: Given $x \in V$ and $\xi \in \Omega^{(2k)}$, let $t_\xi^{(x)} \in \text{BU}_2(\{\text{id}\})$ denote the unique label-respecting translation with $t_\xi^{(x)}(x) = x_\xi$. Given an element $\xi = (\omega_1, \dots, \omega_{2k}) \in \Omega^{(2k)}$, we set $\bar{\xi} := (\omega_{2k}, \dots, \omega_1) \in \Omega^{(2k)}$. Then $(t_\xi^{(x)})^{-1} = t_{\bar{\xi}}^{(x)}$, and if $\Omega_+^{(2k)} \subseteq \Omega^{(2k)}$ is such that for every $\xi \in \Omega^{(2k)}$ exactly one of $\{\xi, \bar{\xi}\}$ belongs to $\Omega_+^{(2k)}$, then $\Omega_+^{(2k)} = \Omega_+^{(2k)} \sqcup \bar{\Omega}_+^{(2k)}$ where $\bar{\Omega}_+^{(2k)} := \{\bar{\xi} \mid \xi \in \Omega_+^{(2k)}\}$.

LEMMA 4.39. Let $x \in V_1$. Then $\text{BU}_2(\{\text{id}\}) = \langle \{t_\xi^{(x)} \mid \xi \in \Omega^{(2)}\} \rangle \cong F_{\Omega_+^{(2)}}$, the free group on the set $\Omega_+^{(2)}$.

PROOF. Every element of $\text{BU}_{2k}(\{\text{id}\})$ is uniquely determined by its image on x . To see that $\text{BU}_2(\{\text{id}\}) = \langle \{t_\xi^{(x)} \mid \xi \in \Omega^{(2)}\} \rangle$ it thus suffices to show that $\{t_\xi^{(x)} \mid \xi \in \Omega^{(2)}\}$ is transitive on V_1 . Indeed, let $y \in V_1$. Then $y = x_\xi$ for some $\xi \in \Omega^{(2k)}$ where $2k = d(x, y)$. Write $\xi = (\xi_1, \dots, \xi_k) \in (\Omega^{(2)})^k$. Then $t_{\xi_1}^{(x)} \circ \dots \circ t_{\xi_k}^{(x)} = t_\xi^{(x)}$ as every $t_{\xi_i}^{(x)}$ ($i \in \{1, \dots, k\}$) is label respecting. Hence, $t_{\xi_1}^{(x)} \circ \dots \circ t_{\xi_k}^{(x)}(x) = y$ and

$$\langle \{t_\xi^{(x)} \mid \xi \in \Omega^{(2)}\} \rangle \rightarrow F_{\Omega^{(2)}}, \quad \begin{cases} t_\xi^{(x)} \mapsto \xi & \xi \in \Omega_+^{(2)} \\ t_\xi^{(x)} \mapsto \xi^{-1} & \xi \notin \Omega_+^{(2)} \end{cases}$$

yields a well-defined isomorphism. □

4.6.2. Compatibility and discreteness. In order to describe the compatibility and the discreteness condition in the bipartite setting, we first introduce a suitable realization of $\text{Aut}(B_{d,2k})$ ($k \in \mathbb{N}$), similar to that at the beginning of Section 4.4. Let $\text{Aut}(B_{d,1}) \cong \text{Sym}(\Omega)$ and $\text{Aut}(B_{d,2})$ be as before. For $k \geq 2$, we inductively identify $\text{Aut}(B_{d,2k})$ with its image under

$$\begin{aligned} \text{Aut}(B_{d,2k}) &\rightarrow \text{Aut}(B_{d,2(k-1)}) \times \prod_{\xi \in \Omega^{(2)}} \text{Aut}(B_{d,2(k-1)}) \\ \alpha &\mapsto (\sigma_{2(k-1)}(\alpha, b), (\sigma_{2(k-1)}(\alpha, b_\xi))_\xi) \end{aligned}$$

where $\text{Aut}(B_{d,2(k-1)})$ acts on $\Omega^{(2)}$ by permuting the factors according to its action on $S(b, 2) \cong \Omega^{(2)}$. In addition, consider the map $\text{pr}_\xi : \text{Aut}(B_{d,2k}) \rightarrow \text{Aut}(B_{d,2(k-1)})$, $\alpha \mapsto \sigma_{2(k-1)}(\alpha, b_\xi)$ for every $\xi \in \Omega^{(2)}$, as well as

$$p_\xi = (\pi_{2(k-1)}, \text{pr}_\xi) : \text{Aut}(B_{d,2k}) \rightarrow \text{Aut}(B_{d,2(k-1)}) \times \text{Aut}(B_{d,2(k-1)}).$$

For $k \geq 2$, conditions (C) and (D) for $F \leq \text{Aut}(B_{d,2k})$ now read as follows:

$$\begin{aligned} &\text{for all } \alpha \in F, \text{ for all } \xi \in \Omega^{(2)} : \text{there exists } \alpha_\xi \in F : \\ &\pi_{2(k-1)}(\alpha_\xi) = \text{pr}_\xi(\alpha), \text{pr}_{\bar{\xi}}(\alpha_\xi) = \pi_{2(k-1)}(\alpha); \end{aligned} \tag{C}$$

$$\text{for all } \xi \in \Omega^{(2)} : p_\xi|_F^{-1}(\text{id}, \text{id}) = \{\text{id}\}. \tag{D}$$

For $k = 1$ we have, using the maps pr_ω ($\omega \in \Omega$) as in Section 4.4,

$$\text{for all } \alpha \in F, \text{ for all } \xi = (\omega_1, \omega_2) \in \Omega^{(2)} : \text{there exists } \alpha_\xi \in F : \text{pr}_{\omega_2}(\alpha_\xi) = \text{pr}_{\omega_1}(\alpha), \tag{C}$$

$$\text{for all } \omega \in \Omega : \text{pr}_\omega|_F^{-1}(\text{id}) = \{\text{id}\}. \tag{D}$$

Analogues of Proposition 4.12 are proven using the discreteness conditions (D) above. We do not introduce new notation for any of the above as the context always implies which condition is to be considered. The definition of the compatibility sets $C_F(\alpha, S)$ for $F \leq \text{Aut}(B_{d,2k})$ and $S \subseteq \Omega^{(2)}$ carries over from Section 4.2 in a straightforward fashion.

4.6.3. Examples. Let $F \leq \text{Sym}(\Omega)$. Then the group $\Gamma(F) \leq \text{Aut}(B_{d,2})$ introduced in Section 4.4.1 satisfies conditions (C) and (D) for the case $k = 1$ above, and we have $\text{BU}_2(\Gamma(F)) = \text{U}_2(\Gamma(F)) \cap {}^+\text{Aut}(T_d)$.

Similarly, the group $\Phi(F) \leq \text{Aut}(B_{d,2})$ satisfies condition (C) for the case $k = 1$ as $\Gamma(F) \leq \Phi(F)$, and we have $\text{BU}_2(\Phi(F)) = \text{U}_1(F) \cap {}^+\text{Aut}(T_d)$.

The following example gives an analogue of the groups $\Phi(F, N)$. Notice, however, that in this case the second argument is a subgroup of F rather than F_{ω_0} and need not be normal, as the 1-local action at vertices in V_1 and V_2 need not be the same.

EXAMPLE 4.40. Let $F' \leq F \leq \text{Sym}(\Omega)$. Then

$$\text{B}\Phi(F, F') := \{(a, (a_\omega)_{\omega \in \Omega}) \mid a \in F, \text{ for all } \omega \in \Omega : a_\omega \in C_F(a, \omega) \cap F'\} \leq \text{Aut}(B_{d,2})$$

satisfies condition (C) for the case $k = 1$ above, given that $\Gamma(F') \leq \text{B}\Phi(F, F')$. If $F' \setminus \Omega = F \setminus \Omega$, the 1-local action of $\text{B}\Phi(F, F')$ at vertices in V_1 is indeed F , whereas it is F'^+ at vertices in V_2 . This construction is similar to $\mathcal{U}_{\mathcal{L}}(M, N)$ in [27].

The next example constitutes the base case in Section 5.1.5 below.

EXAMPLE 4.41. Let $F \leq \text{Sym}(\Omega)$. Suppose F preserves a nontrivial partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω . Then

$$\Omega_0^{(2)} := \{(\omega_1, \omega_2) \mid \text{there exists } i \in I : \omega_1, \omega_2 \in \Omega_i\} \subseteq \Omega^{(2)}$$

is preserved by the action of $\Phi(F)$ on $S(b, 2) \cong \Omega^{(2)}$. Let $\alpha = (a, (a_\omega)_\omega) \in \Phi(F)$ and $(\omega_1, \omega_2) \in \Omega_0^{(2)}$. Then $\alpha(\omega_1, \omega_2) = (a\omega_1, a_{\omega_1}\omega_2) = (a_{\omega_1}\omega_1, a_{\omega_1}\omega_2) \in \Omega_0^{(2)}$. Also, note that if $\xi = (\omega_1, \omega_2) \in \Omega_0^{(2)}$ then also $\bar{\xi} = (\omega_2, \omega_1) \in \Omega_0^{(2)}$.

The subgroup of $\Phi(F)$ consisting of those elements which are self-compatible in all directions from $\Omega_0^{(2)}$ is precisely given by

$$F^{(2)} := \{(a, (a_\omega)_\omega) \mid a \in F, a_\omega \in C_F(a, \omega) \text{ constant w.r.t. } \mathcal{P}\}$$

in view of condition (C) for the case $k = 1$ above.

Suppose now that $F \leq \text{Aut}(B_{d,2k})$ satisfies (C). Analogous to the group $\Phi_k(F)$ of Section 4.4.2, we define

$$\text{B}\Phi_{2k}(F) := \{(\alpha, (\alpha_\xi)_{\xi \in \Omega^{(2)}}) \mid \alpha \in F, \text{ for all } \xi \in \Omega^{(2)} : \alpha_\xi \in C_F(\alpha, \xi)\} \leq \text{Aut}(B_{d,2(k+1)}).$$

Then $\text{B}\Phi_{2k}(F) \leq \text{Aut}(B_{d,2(k+1)})$ satisfies (C) and $\text{BU}_{2(k+1)}(\text{B}\Phi_{2k}(F)) = \text{BU}_{2k}(F)$. Given $l > k$, we also set $\text{B}\Phi^{2l}(F) := \text{B}\Phi_{2(l-1)} \circ \cdots \circ \text{B}\Phi_{2k}(F)$; cf. Section 4.4.2.

5. Applications

In this section we give three applications of the framework of universal groups. First, we characterize the automorphism types that the quasicycle of a nondiscrete subgroup of $\text{Aut}(T_d)$ may feature in terms of the group's local action, and see that the Burger–Mozes theory does not extend to the transitive case. Second, we give an algebraic characterization of the (P_k) -closures of locally transitive subgroups of

$\text{Aut}(T_d)$ containing an involutive inversion, and thereby partially answer two questions by Banks, Elder and Willis. Third, we offer a new view on the Weiss conjecture.

5.1. Groups acting on trees with nontrivial quascentre. By Proposition 3.11(ii), a nondiscrete, locally semiprimitive subgroup of $\text{Aut}(T_d)$ does not contain any nontrivial quascentral edge-fixating elements. We extend this fact to the following local-to-global-type characterization of quascentral elements.

THEOREM 5.1. *Let $H \leq \text{Aut}(T_d)$ be nondiscrete. If H is locally:*

- (i) *Transitive, then $\text{QZ}(H)$ contains no inversion.*
- (ii) *Semiprimitive, then $\text{QZ}(H)$ contains no nontrivial edge-fixating element.*
- (iii) *Quasprimitive, then $\text{QZ}(H)$ contains no nontrivial elliptic element.*
- (iv) *k -transitive, ($k \in \mathbb{N}$) then $\text{QZ}(H)$ contains no hyperbolic element of length k .*

THEOREM 5.2. *There is a $d \in \mathbb{N}_{\geq 3}$ and a closed, nondiscrete, compactly generated subgroup of $\text{Aut}(T_d)$ which is locally:*

- (i) *Intransitive and contains a quascentral inversion.*
- (ii) *Transitive and contains a nontrivial quascentral edge-fixating element.*
- (iii) *Semiprimitive and contains a nontrivial quascentral elliptic element.*
- (iv) (a) *Intransitive and contains a quascentral hyperbolic element of length 1.*
 (b) *Quasprimitive and contains a quascentral hyperbolic element of length 2.*

PROOF OF THEOREM 5.1. Fix a labelling of T_d and let $H \leq \text{Aut}(T_d)$ be nondiscrete.

For (i), suppose $\iota \in \text{QZ}(H)$ inverts $(x, x_\omega) \in E$. Since H is locally transitive and $\text{QZ}(H) \trianglelefteq H$, there is an inversion $\iota_\omega \in \text{QZ}(H)$ of $(x, x_\omega) \in E$ for all $\omega \in \Omega$. By definition, the centralizer of ι_ω in H is open for all $\omega \in \Omega$. Hence, using the nondiscreteness of H , there is $n \in \mathbb{N}$ such that $H_{B(x,n)}$ commutes with ι_ω for all $\omega \in \Omega$ and $H_{B(x,n+1)} \neq \{\text{id}\}$. However, $H_{B(x,n)} = \iota_\omega H_{B(x,n)} \iota_\omega^{-1} = H_{B(x_\omega,n)}$ for all $\omega \in \Omega$; that is, $H_{B(x,n+1)} \subseteq H_{B(x,n)}$ in contradiction to the above.

Part (ii) is Proposition 3.11(ii) and part (iii) is [2, Proposition 1.2.1(ii)]. Here, the closedness assumption is unnecessary.

For part (iv), suppose $\tau \in \text{QZ}(H)$ is a translation of length k that maps $x \in V$ to $x_\xi \in V$ for some $\xi \in \Omega^{(k)}$. Since H is locally k -transitive and $\text{QZ}(H) \trianglelefteq H$, there is a translation $\tau_\xi \in \text{QZ}(H)$ that maps x to x_ξ for all $\xi \in \Omega^{(k)}$. By definition, the centralizer of τ_ξ in H is open for all $\xi \in \Omega^{(k)}$. Hence, using the nondiscreteness of H , there is $n \in \mathbb{N}$ such that $H_{B(x,n)}$ commutes with τ_ξ for all $\xi \in \Omega^{(k)}$ and $H_{B(x,n+1)} \neq \{\text{id}\}$. However, $H_{B(x,n)} = \tau_\xi H_{B(x,n)} \tau_\xi^{-1} = H_{B(x_\xi,n)}$ for all $\xi \in \Omega^{(k)}$; that is, $H_{B(x,n+k)} \subseteq H_{B(x,n)}$ in contradiction to the above. \square

We complement part (ii) of Theorem 5.1 with the following result which is inspired by [2, Proposition 3.1.2] and [24, Conjecture 2.63],

PROPOSITION 5.3. *Let $H \leq \text{Aut}(T_d)$ be nondiscrete and locally semiprimitive. If all orbits of $H \curvearrowright \partial T_d$ are uncountable then $\text{QZ}(H)$ is trivial.*

PROOF. By Theorem 5.1, the group $\text{QZ}(H)$ contains no inversions. Let $S \subseteq \partial T_d$ be the set of fixed points of hyperbolic elements in $\text{QZ}(H)$. Since $\text{QZ}(H) \trianglelefteq H$, the set S is H -invariant. Also, $\text{QZ}(H)$ is discrete by Theorem 5.1 and hence countable as H is second-countable. Thus S is countable and hence empty. We conclude that $\text{QZ}(H) \trianglelefteq H$ does not contain elliptic elements in view of [10, Lemma 6.4]. \square

The following strengthening of Theorem 5.2(ii) proved in Section 5.1.2 below shows that the Burger–Mozes theory does not generalize to the locally transitive case.

THEOREM 5.4. *There exist $d \in \mathbb{N}_{\geq 3}$ and a closed, nondiscrete, compactly generated, locally transitive subgroup of $\text{Aut}(T_d)$ with open, hence nondiscrete, quasicentre.*

We prove Theorem 5.2 by construction in the rest of this subsection. Whereas parts (i)–(iv)(a) all use groups of the form $\bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$ for appropriate local actions $F^{(k)} \leq \text{Aut}(B_{d,k})$, part (iv)(b) uses a group of the form $\bigcap_{k \in \mathbb{N}} \text{BU}(F^{(2k)})$. The various parts appear similar at first glance but vary in detail.

PROOF OF THEOREM 5.2(i). For certain intransitive $F \leq \text{Sym}(\Omega)$ we construct a closed, nondiscrete, compactly generated, vertex-transitive group $H(F) \leq \text{Aut}(T_d)$ which locally acts like F and contains a quasicentral involutive inversion.

Let $F \leq \text{Sym}(\Omega)$. Assume that the partition $F \backslash \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω into F -orbits has at least three elements and that $F_{\Omega_i} \neq \{\text{id}\}$ for all $i \in I$.

Fix an orbit Ω_0 of size at least 2 and $\omega_0 \in \Omega_0$. Define groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ for $k \in \mathbb{N}$ inductively by $F^{(1)} := F$ and

$$F^{(k+1)} := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in F^{(k)}, \alpha_\omega \in C_{F^{(k)}}(\alpha, \omega) \text{ constant w.r.t. } F \backslash \Omega, \alpha_{\omega_0} = \alpha\}. \quad \square$$

PROPOSITION 5.5. *The groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) defined above satisfy the following assertions.*

- (i) *Every $\alpha \in F^{(k)}$ is self-compatible in directions from Ω_0 .*
- (ii) *The compatibility set $C_{F^{(k)}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in F^{(k)}$ and $i \in I$. In particular, the group $F^{(k)}$ satisfies (C).*
- (iii) *The compatibility set $C_{F^{(k)}}(\text{id}, \Omega_i)$ is nontrivial for all $\Omega_i \neq \Omega_0$. In particular, the group $F^{(k)}$ does not satisfy (D).*

PROOF. We prove all three properties simultaneously by induction. For $k = 1$, assertions (i) and (ii) are trivial. Assertion (iii) translates to F_{Ω_i} being nontrivial for all $\Omega_i \neq \Omega_0$, which is an assumption. Now, assume that all properties hold for $F^{(k)}$. Then the definition of $F^{(k+1)}$ is meaningful because of (i) and it is a subgroup of $\text{Aut}(B_{d,k+1})$ because F preserves each Ω_i ($i \in I$). Assertion (i) is now evident. Assertion (ii) carries over from $F^{(k)}$ to $F^{(k+1)}$. So does (iii) since $|F \backslash \Omega| \geq 3$. \square

DEFINITION 5.6. Retain the above notation. Define $H(F) := \bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$.

Now, $H(F)$ is compactly generated, vertex-transitive and contains an involutive inversion because $U_1(\{\text{id}\}) \leq H(F)$. Also, $H(F)$ is closed as an intersection of closed

sets. The 1-local action of H is given by $F = F^{(1)}$ because $\Gamma^k(F) \leq F^{(k)}$ for all $k \in \mathbb{N}$, and therefore $D(F) \leq H(F)$.

LEMMA 5.7. *The group $H(F)$ is nondiscrete.*

PROOF. Let $x \in V$ and $n \in \mathbb{N}$. We construct a nontrivial element $h \in H(F)$ which fixes $B(x, n)$. Set $\alpha_n := \text{id} \in F^{(n)}$. By parts (i) and (iii) of Proposition 5.5 as well as the definition of $F^{(n+1)}$, there is a nontrivial element $\alpha_{n+1} \in F^{(n+1)}$ with $\pi_n \alpha_{n+1} = \alpha_n$. Applying parts (i) and (ii) of Proposition 5.5 repeatedly, we obtain nontrivial elements $\alpha_k \in F^{(k)}$ for all $k \geq n + 1$ with $\pi_k \alpha_{k+1} = \alpha_k$. Set $\alpha_k := \text{id} \in F^{(k)}$ for all $k \leq n$ and define $h \in \text{Aut}(T_d)_x$ by fixing x and setting $\sigma_k(h, x) := \alpha_k \in F^{(k)}$. Since $F^{(l)} \leq \Phi^l(F^{(k)})$ for all $k \leq l$, we conclude that $h \in \bigcap_{k \in \mathbb{N}} U_k(F^{(k)}) = H(F)$. \square

PROPOSITION 5.8. *The quasicycle of $H(F)$ contains an involutive inversion.*

PROOF. Let $x \in V$. We show that the group $\text{QZ}(H(F))$ contains the label-respecting inversion ι_ω of $(x, x_\omega) \in E$ for all $\omega \in \Omega_0$. To see this, let $h \in H(F)_{B(x, 1)}$ and $\omega \in \Omega_0$. Then $h u_\omega(x) = x_\omega = \iota_\omega h(x)$ and $\sigma_k(h u_\omega, x) = \sigma_k(h, \iota_\omega x) \sigma_k(\iota_\omega, x) = \sigma_k(h, x_\omega) = \sigma_k(\iota_\omega, h x)$ $\sigma_k(h, x) = \sigma_k(\iota_\omega h, x)$ for all $k \in \mathbb{N}$ since $h \in U_{k+1}(F^{(k+1)})$. That is, ι_ω commutes with $H(F)_{B(b, 1)}$. \square

PROOF OF THEOREM 5.2(ii). For certain transitive $F \leq \text{Sym}(\Omega)$ we construct a closed, nondiscrete, compactly generated, vertex-transitive group $H(F) \leq \text{Aut}(T_d)$ that locally acts like F and has open quasicycle.

Let $F \leq \text{Sym}(\Omega)$ be transitive. Assume that F preserves a nontrivial partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω and that $F_{\Omega_i} \neq \{\text{id}\}$ for all $i \in I$. Further, suppose that F^+ is abelian and preserves \mathcal{P} setwise. \square

EXAMPLE 5.9. Let $F' \leq \text{Sym}(\Omega')$ be regular abelian and $P \leq \text{Sym}(\Lambda)$ regular. Then $F := F' \wr P \leq \text{Sym}(\Omega' \times \Lambda)$ satisfies the above properties as $F^+ = \prod_{\lambda \in \Lambda} F'$.

Define groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ for $k \in \mathbb{N}$ inductively by $F^{(1)} := F$ and

$$F^{(k+1)} := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in F^{(k)}, \alpha_\omega \in C_{F^{(k)}}(\alpha, \omega) \text{ constant w.r.t. } \mathcal{P}\}.$$

PROPOSITION 5.10. *The groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) defined above satisfy the following assertions.*

- (i) *The compatibility set $C_{F^{(k)}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in F^{(k)}$ and $i \in I$. In particular, the group $F^{(k)}$ satisfies (C).*
- (ii) *The compatibility set $C_{F^{(k)}}(\text{id}, \Omega_i)$ is nontrivial for all $i \in I$. In particular, the group $F^{(k)}$ does not satisfy (D).*
- (iii) *The group $F^{(k)} \cap \Phi^k(F^+)$ is abelian.*

PROOF. We prove all three properties simultaneously by induction. For $k = 1$, assertion (i) is trivial whereas (iii) is an assumption. Assertion (ii) translates to F_{Ω_i} being nontrivial for all $i \in I$, which is an assumption. Now, assume all properties hold for $F^{(k)}$. Then the definition of $F^{(k+1)}$ is meaningful because of (i) and it is a subgroup of

$\text{Aut}(B_{d,k})$ because F preserves \mathcal{P} . Assertion (ii) carries over from $F^{(k)}$ to $F^{(k+1)}$. Finally, (iii) follows inductively because F^+ preserves \mathcal{P} setwise. \square

DEFINITION 5.11. Retain the above notation. Define $H(F) := \bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$.

Now, $H(F)$ is compactly generated, vertex-transitive and contains an involutive inversion because $U_1(\{\text{id}\}) \leq H(F)$. Also, $H(F)$ is closed as an intersection of closed sets. The 1-local action of H is given by $F = F^{(1)}$ because $\Gamma_k(F) \leq F^{(k)}$ for all $k \in \mathbb{N}$ and therefore $D(F) \leq H(F)$.

LEMMA 5.12. *The group $H(F)$ is nondiscrete.*

PROOF. Let $x \in V$ and $n \in \mathbb{N}$. We construct a nontrivial element $h \in H(F)$ that fixes $B(x, n)$. Consider $\alpha_n := \text{id} \in F^{(n)}$. By part (ii) of Proposition 5.10 as well as the definition of $F^{(n+1)}$, there is a nontrivial element $\alpha_{n+1} \in F^{(n+1)}$ with $\pi_n \alpha_{n+1} = \alpha_n$. Applying part (i) of Proposition 5.10 repeatedly, we obtain nontrivial elements $\alpha_k \in F^{(k)}$ for all $k \geq n + 1$ with $\pi_k \alpha_{k+1} = \alpha_k$. Set $\alpha_k := \text{id} \in F^{(k)}$ for all $k \leq n$ and define $h \in \text{Aut}(T_d)_x$ by fixing x and setting $\sigma_k(h, x) := \alpha_k \in F^{(k)}$. Since $F^{(l)} \leq \Phi^l(F^{(k)})$ for all $k \leq l$, we conclude that $h \in \bigcap_{k \in \mathbb{N}} U_k(F^{(k)}) = H(F)$. \square

PROPOSITION 5.13. *The group $H(F)$ has open quasicentre.*

PROOF. The group $H(F)_{B(x,1)}$ is a subgroup of the group $H(F^+)_{x,1}$ which is abelian by part (iii) of Proposition 5.10. Hence, $H(F)_{B(x,1)} \leq \text{QZ}(H(F))$. \square

REMARK 5.14. Without assuming local transitivity one can achieve abelian point-stabilizers, following the construction of the previous section. This cannot happen for nondiscrete locally transitive groups $H \leq \text{Aut}(T_d)$ that are vertex-transitive as the following argument shows. By Proposition 2.6, the group H is contained in $U(F)$ where $F \leq \text{Sym}(\Omega)$ is the local action of H . If H_x is abelian, then so is F . Since any transitive abelian permutation group is regular we conclude that $U(F)$ and hence H are discrete. In this sense, the construction of this section is efficient.

PROOF OF THEOREM 5.2(iii). For certain semiprimitive $F \leq \text{Sym}(\Omega)$ we construct a closed, nondiscrete, compactly generated, vertex-transitive group $H(F) \leq \text{Aut}(T_d)$ that locally acts like F and contains a nontrivial quasicentral elliptic element.

Let $F \leq \text{Sym}(\Omega)$ be semiprimitive. Suppose F preserves a nontrivial partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω and that $F_{\Omega_i} \neq \{\text{id}\}$ for all $i \in I$. Further, suppose that F contains a nontrivial central element τ which preserves \mathcal{P} setwise. \square

EXAMPLE 5.15. Consider $\text{SL}(2, 3) \curvearrowright \mathbb{F}_3^2 \setminus \{0\} = \{\pm e_1, \pm e_2, \pm(e_1 + e_2), \pm(e_1 - e_2)\}$ where e_1, e_2 are the standard basis vectors. We have $Z(\text{SL}(2, 3)) = \{\pm \text{Id}\}$. The blocks of size 2 are as listed above, given that $\text{SL}(2, 3)_{e_1} \leq_2 \pm \text{SL}(2, 3)_{e_1}$.

Define groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ for $k \in \mathbb{N}$ inductively by $F^{(1)} := F$ and

$$F^{(k+1)} := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in F^{(k)}, \alpha_\omega \in C_{F^{(k)}}(\alpha, \omega) \text{ constant w.r.t. } \mathcal{P}\}.$$

PROPOSITION 5.16. *The groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) defined above satisfy the following assertions.*

- (i) *The compatibility set $C_{F^{(k)}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in F^{(k)}$ and $i \in I$. In particular, the group $F^{(k)}$ satisfies (C).*
- (ii) *The compatibility set $C_{F^{(k)}}(\text{id}, \Omega_i)$ is nontrivial for all $i \in I$. In particular, the group $F^{(k)}$ does not satisfy (D).*
- (iii) *The element $\gamma_k(\tau) \in \text{Aut}(B_{d,k})$ is central in $F^{(k)}$.*

PROOF. We prove all three properties simultaneously by induction. For $k = 1$, assertion (i) is trivial whereas (iii) is an assumption. Assertion (ii) translates to F_{Ω_i} being nontrivial for all $i \in I$, which is an assumption. Now, assume all properties hold for $F^{(k)}$. Then the definition of $F^{(k+1)}$ is meaningful because of (i) and it is a subgroup of $\text{Aut}(B_{d,k+1})$ because F preserves \mathcal{P} . Assertion (ii) carries over from $F^{(k)}$ to $F^{(k+1)}$. Finally, (iii) follows inductively because τ and hence τ^{-1} preserves \mathcal{P} setwise: for $\tilde{\alpha} = (\alpha, (\alpha_\omega)_\omega) \in F^{(k+1)}$ we have

$$\gamma^{k+1}(\tau)\tilde{\alpha}\gamma^{k+1}(\tau)^{-1} = (\gamma^k(\tau)\alpha\gamma^k(\tau)^{-1}, (\gamma^k(\tau)\alpha_{\tau^{-1}(\omega)}\gamma^k(\tau)^{-1})_\omega). \quad \square$$

DEFINITION 5.17. Retain the above notation. Define $H(F) := \bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$.

Now, $H(F)$ is compactly generated, vertex-transitive and contains an involutive inversion because $U_1(\{\text{id}\}) \leq H(F)$. Also, $H(F)$ is closed as an intersection of closed sets. The 1-local action of H is given by $F = F^{(1)}$ because $\Gamma^k(F) \leq F^{(k)}$ for all $k \in \mathbb{N}$ and therefore $D(F) \leq H(F)$.

LEMMA 5.18. *The group $H(F)$ is nondiscrete.*

PROOF. Let $x \in V$ and $n \in \mathbb{N}$. We construct a nontrivial element $h \in H(F)$ that fixes $B(x, n)$. Consider $\alpha_n := \text{id} \in F^{(n)}$. By part (ii) of Proposition 5.16 and the definition of $F^{(n+1)}$, there is a nontrivial $\alpha_{n+1} \in F^{(n+1)}$ with $\pi_n\alpha_{n+1} = \alpha_n$. Applying part (i) of Proposition 5.16 repeatedly, we obtain nontrivial elements $\alpha_k \in F^{(k)}$ for all $k \geq n + 1$ with $\pi_k\alpha_{k+1} = \alpha_k$. Set $\alpha_k := \text{id} \in F^{(k)}$ for all $k \leq n$ and define $h \in \text{Aut}(T_d)_x$ by fixing x and setting $\sigma_k(h, x) := \alpha_k \in F^{(k)}$. Since $F^{(l)} \leq \Phi^l(F^{(k)})$ for all $k \leq l$, we conclude that $h \in \bigcap_{k \in \mathbb{N}} U_k(F^{(k)}) = H(F)$. □

PROPOSITION 5.19. *The quasicycle of $H(F)$ contains a nontrivial elliptic element.*

PROOF. By Proposition 5.16, the element $d(\tau)$ that fixes x and whose 1-local action is τ everywhere commutes with $H(F)_x$. Hence, $d(\tau) \in \text{QZ}(H(F))$. □

REMARK 5.20. The argument of this section does not work in the quasiprimitive case because a quasiprimitive group $F \leq \text{Sym}(\Omega)$ with nontrivial centre is abelian and regular. If $Z(F) \trianglelefteq F$ is nontrivial then it is transitive, and it suffices to show that F^+ is trivial. Suppose $a \in F_\omega$ moves $\omega' \in \Omega$. Pick $z \in Z(F)$ with $z(\omega) = \omega'$. Then $za(\omega) = \omega' \neq az(\omega)$, contradicting the assumption that $z \in Z(F)$.

PROOF OF THEOREM 5.2(iv)(a). For certain intransitive $F \leq \text{Sym}(\Omega)$ we construct a closed, nondiscrete, compactly generated, vertex-transitive group $H(F) \leq \text{Aut}(T_d)$ that locally acts like F and contains a quasicontral hyperbolic element of length 1.

Let $F \leq \text{Sym}(\Omega)$. Assume that the partition $F \backslash \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω has at least three elements and that $Z(F) \neq \{\text{id}\}$. Choose a nontrivial element $\tau \in Z(F)$ and $\omega_0 \in \Omega_0 \in F \backslash \Omega$ with $\tau(\omega_0) \neq \omega_0$. Further, suppose that $F_{\Omega_i} \neq \{\text{id}\}$ for all $\Omega_i \neq \Omega_0$.

Define groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ for $k \in \mathbb{N}$ inductively by $F^{(1)} := F$ and

$$F^{(k+1)} := \{(\alpha, (\alpha_\omega)_\omega) \mid \alpha \in F^{(k)}, \alpha_\omega \in C_{F^{(k)}}(\alpha, \omega) \text{ constant w.r.t. } F \backslash \Omega, \alpha_{\omega_0} = \alpha\}. \quad \square$$

PROPOSITION 5.21. *The groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) defined above satisfy the following assertions.*

- (i) *Every $\alpha \in F^{(k)}$ is self-compatible in directions from Ω_0 .*
- (ii) *The compatibility set $C_{F^{(k)}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in F^{(k)}$ and $i \in I$. In particular, the group $F^{(k)}$ satisfies (C).*
- (iii) *The compatibility set $C_{F^{(k)}}(\text{id}, \Omega_i)$ is nontrivial for all $i \in I \setminus \{0\}$. In particular, the group $F^{(k)}$ does not satisfy (D).*
- (iv) *The element $\gamma_k(\tau) \in \text{Aut}(B_{d,k})$ is central in $F^{(k)}$.*

PROOF. We prove all four properties simultaneously by induction. For $k = 1$, assertions (i) and (ii) are trivial. Assertion (iii) translates to F_{Ω_i} being nontrivial for all $i \in I \setminus \{0\}$, which is an assumption, as is (iv). Now, assume that all properties hold for $F^{(k)}$. Then the definition of $F^{(k+1)}$ is meaningful because of (i) and it is a subgroup of $\text{Aut}(B_{d,k})$ because F preserves $F \backslash \Omega$. Assertion (i) is now evident. Assertions (ii) and (iii) carry over from $F^{(k)}$ to $F^{(k+1)}$. Finally, (iv) follows inductively because τ and hence τ^{-1} preserves $F \backslash \Omega$ setwise: for $\tilde{\alpha} = (\alpha, (\alpha_\omega)_\omega) \in F^{(k+1)}$ we have

$$\gamma^{k+1}(\tau)\tilde{\alpha}\gamma^{k+1}(\tau)^{-1} = (\gamma^k(\tau)\alpha\gamma^k(\tau)^{-1}, (\gamma^k(\tau)\alpha_{\tau^{-1}(\omega)}\gamma^k(\tau)^{-1})_\omega). \quad \square$$

DEFINITION 5.22. Retain the above notation. Define $H(F) := \bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$.

Now, $H(F)$ is compactly generated, vertex-transitive and contains an involutive inversion because $U_1(\{\text{id}\}) \leq H(F)$. Also, $H(F)$ is closed as the intersection of all its (P_k) -closures. The 1-local action of H is given by $F = F^{(1)}$ as $\Gamma^k(F) \leq F^{(k)}$ for all $k \in \mathbb{N}$ and therefore $D(F) \leq H$.

LEMMA 5.23. *The group $H(F)$ is nondiscrete.*

PROOF. Let $x \in V$ and $n \in \mathbb{N}$. We construct a nontrivial element $h \in H(F)$ which fixes $B(x, n)$. Consider $\alpha_n := \text{id} \in F^{(n)}$. By parts (i) and (iii) of Proposition 5.21 as well as the definition of $F^{(n+1)}$, there is a nontrivial element $\alpha_{n+1} \in F^{(n+1)}$ with $\pi_n \alpha_{n+1} = \alpha_n$. Applying parts (i) and (ii) of Proposition 5.21 repeatedly, we obtain nontrivial elements $\alpha_k \in F^{(k)}$ for all $k \geq n + 1$ with $\pi_k \alpha_{k+1} = \alpha_k$. Set $\alpha_k := \text{id} \in F^{(k)}$ for all $k \leq n$, and define $h \in \text{Aut}(T_d)_x$ by fixing x and setting $\sigma_k(h, x) := \alpha_k \in F^{(k)}$. Since $F^{(l)} \leq \Phi^l(F^{(k)})$ for all $k \leq l$ we conclude that $h \in \bigcap_{k \in \mathbb{N}} U_k(F^{(k)}) = H(F)$. \square

PROPOSITION 5.24. *The quasicontral of $H(F)$ contains a translation of length 1.*

PROOF. Fix $x \in V$ and let τ be as above. Consider the line L through x with labels

$$\dots, \tau^{-2}\omega_0, \tau^{-1}\omega_0, \omega_0, \tau\omega_0, \tau^2\omega_0, \dots$$

Define $t \in D(F)$ by $t(x) = x_{\omega_0}$ and $\sigma_1(t, y) = \tau$ for all $y \in V$. Then t is a translation of length 1 along L . Furthermore, t commutes with $H(F)_{B(x,1)}$. Indeed, let $g \in H(F)_{B(x,1)}$. Then $(gt)(x) = t(x) = (tg)(x)$ and

$$\sigma_k(gt, x) = \sigma_k(g, tx)\sigma_k(t, x) = \sigma_k(t, x)\sigma_k(g, x) = \sigma_k(t, gx)\sigma_k(g, x) = \sigma_k(tg, x)$$

for all $k \in \mathbb{N}$ because $\sigma_k(t, x) = \gamma^k(\tau) \in Z(F^{(k)})$ and $g \in U_{k+1}(F^{(k+1)})_{B(x,1)}$. □

PROOF OF THEOREM 5.2(iv)(b). For certain quasiprimitive $F \leq \text{Sym}(\Omega)$ we construct a closed, nondiscrete, compactly generated group $H(F) \leq \text{Aut}(T_d)$ that locally acts like F and contains a quasicentral hyperbolic element of length 2.

Let $F \leq \text{Sym}(\Omega)$ be quasiprimitive. Suppose F preserves a nontrivial partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$. Further, suppose that $F_{\Omega_i} \neq \{\text{id}\}$ and that $F_{\omega_i} \curvearrowright \Omega_i \setminus \{\omega_i\}$ is transitive for all $i \in I$ and $\omega_i \in \Omega_i$. □

EXAMPLE 5.25. Consider the action $A_5 \curvearrowright A_5/C_5$. It has blocks of size $[D_5 : C_5] = 2$ and nontrivial block stabilizers as $C_5 \cap \tau C_5 \tau^{-1} = C_5$ for all $\tau \in D_5$ given that $C_5 \trianglelefteq D_5$.

Retain the notation of Example 4.41. Define groups $F^{(2k)} \leq \text{Aut}(B_{d,2k})$ for $k \in \mathbb{N}$ inductively by $F^{(2)} = \{(a, (a_\omega)_\omega) \mid a \in F, a_\omega \in C_F(a, \omega) \text{ constant w.r.t. } \mathcal{P}\}$ and

$$F^{(2(k+1))} := \{(\alpha, (\alpha_\xi)_\xi) \mid \alpha \in F^{(2k)}, \alpha_\xi \in C_{F^{(2k)}}(\alpha, \xi), \text{ for all } \xi \in \Omega_0^{(2)} : \alpha_\xi = \alpha\}.$$

PROPOSITION 5.26. *The groups $F^{(2k)} \leq \text{Aut}(B_{d,2k})$ ($k \in \mathbb{N}$) defined above satisfy the following assertions.*

- (i) Every $\alpha \in F^{(2k)}$ is self-compatible in all directions from $\Omega_0^{(2)}$.
- (ii) The compatibility set $C_{F^{(2k)}}(\alpha, \xi)$ is nonempty for all $\alpha \in F^{(2k)}$ and $\xi \in \Omega^{(2)}$. In particular, the group $F^{(2k)}$ satisfies (C).
- (iii) The compatibility set $C_{F^{(2k)}}(\text{id}, \xi)$ is nontrivial for all $\xi \in \Omega^{(2)}$. In particular, the group $F^{(2k)}$ does not satisfy (D).

PROOF. We prove all three properties simultaneously by induction. For $k = 1$, assertion (i) holds by construction of $F^{(2)}$, as do (ii) and (iii). Now assume that all properties hold for $F^{(2k)}$. Then the definition of $F^{(2(k+1))}$ is meaningful because of (i) and it is a subgroup because $F^{(2)}$ preserves $\Omega_0^{(2)}$. Also, $F^{(2(k+1))}$ satisfies (i) because $\Omega_0^{(2)}$ is inversion-closed. Assertions (ii) and (iii) carry over from $F^{(2k)}$. □

DEFINITION 5.27. Retain the above notation. Define $H(F) := \bigcap_{k \in \mathbb{N}} \text{BU}_{2k}(F^{(2k)})$.

Now, $H(F)$ is closed as an intersection of closed sets and compactly generated by $H(F)_x$ for some $x \in V_1$ and a finite generating set of $\text{BU}_2(\{\text{id}\})^+$; see Lemma 4.39. For vertices in V_1 , the 1-local action is F because $\Gamma^{2k}(F) \leq F^{(2k)}$. For vertices in V_2 the 1-local action is $F^+ = F$ as $\Gamma^2(F) \leq F^{(2)}$.

LEMMA 5.28. *The group $H(F)$ is nondiscrete.*

PROOF. Let $x \in V_1$ and $n \in \mathbb{N}$. We construct a nontrivial element $h \in H(F)$ that fixes $B(x, 2n)$. Consider $\alpha_{2n} := \text{id} \in F^{(2n)}$: By parts (i) and (iii) of Proposition 5.5 and the definition of $F^{(2(n+1))}$, there is a nontrivial element $\alpha_{2(n+1)} \in F^{(2(n+1))}$ with $\pi_{2n}\alpha_{2(n+1)} = \alpha_{2n}$. Applying parts (i) and (ii) of Proposition 5.26 repeatedly, we obtain nontrivial elements $\alpha_{2k} \in F^{(2k)}$ for all $k \geq n + 1$ with $\pi_{2k}\alpha_{2(k+1)} = \alpha_{2k}$. Set $\alpha_{2k} := \text{id} \in F^{(2k)}$ for all $k \leq n$ and define $h \in \text{Aut}(T_d)_x$ by fixing x and setting $\sigma_{2k}(h, x) := \alpha_{2k} \in F^{(2k)}$. Since $F^{(2l)} \leq \text{B}\Phi^{2l}(F^{(2k)})$ for all $k \leq l$, we conclude that $h \in \bigcap_{k \in \mathbb{N}} \text{BU}_{2k}(F^{(2k)}) = H(F)$. \square

PROPOSITION 5.29. *The quasicentre of $H(F)$ contains a translation of length 2.*

PROOF. Fix $x \in V_1$ and $\xi = (\omega_1, \omega_2) \in \Omega_0^{(2)}$. Consider the line L through b with labels

$$\dots, \omega_1, \omega_2, \omega_1, \omega_2, \dots$$

Define $t \in D(F)$ by $t(x) = x_\xi$ and $\sigma_1(t, y) = \text{id}$ for all $y \in V$. Then t is a translation of length 2 along L . Furthermore, t commutes with $H(F)_{B(x,2)}$. Indeed, let $g \in H(F)_{B(x,2)}$. Then $gt(x) = t(x) = tg(x)$ and, for all $k \in \mathbb{N}$,

$$\begin{aligned} \sigma_{2k}(gt, x) &= \sigma_{2k}(g, tx)\sigma_{2k}(t, x) = \sigma_{2k}(g, x_\xi) \\ &= \sigma_{2k}(g, x) = \sigma_{2k}(t, gx)\sigma_{2k}(g, x) = \sigma_{2k}(tg, x) \end{aligned}$$

as $\sigma_l(t, y) = \text{id}$ for all $l \in \mathbb{N}$ and $y \in V(T_d)$, and $g \in \text{BU}_{2(k+1)}(F^{(2(k+1))})_{B(b,2)}$. \square

REMARK 5.30. We argue that the construction of this section does not carry over to any primitive $F \leq \text{Sym}(\Omega)$ and $\Gamma(F) \leq F^{(2)} \leq \Phi(F)$.

First, note that $\Phi(F) \setminus \Omega^{(2)} = \Gamma(F) \setminus \Omega^{(2)}$. For $\alpha := (a, (a_\omega)_{\omega \in \Omega}) \in \Phi(F)$ and $(\omega_1, \omega_2) \in \Omega^{(2)}$ we have $\alpha(\omega_1, \omega_2) = (a\omega_1, a_{\omega_1}\omega_2) \in \{(a\omega_1, aF_{\omega_1}\omega_2)\} \subseteq \Gamma(F)(\omega_1, \omega_2)$. We now observe the following obstruction to nondiscreteness. Given any orbit $\Omega_0^{(2)} \in \Phi(F) \setminus \Omega^{(2)} = F^{(2)} \setminus \Omega^{(2)}$, the subgroup of $\Phi(F)$ consisting of elements that are self-compatible in all directions from $\Omega_0^{(2)}$ is precisely $\Gamma(F)$.

Indeed, every element of $\Gamma(F)$ is self-compatible in all directions from $\Omega^{(2)} \supseteq \Omega_0^{(2)}$. Conversely, let $(a, (a_\omega)_\omega) \in \Phi(F)$ be self-compatible in all directions from $\Omega_0^{(2)}$. Consider the equivalence relation on Ω defined by $\omega_1 \sim \omega_2$ if and only if $a_{\omega_1} = a_{\omega_2}$. Since $a_{\omega_1} = a_{\omega_2}$ whenever $\xi := (\omega_1, \omega_2) \in \Omega_0^{(2)}$, this relation is F -invariant. Since $\Gamma(F) \leq \Phi(F)$ we have $\gamma(a)(\omega_1, \omega_2) = (a\omega_1, a\omega_2) \in \Omega_0^{(2)}$ for all $a \in F$ whenever $(\omega_1, \omega_2) \in \Omega_0^{(2)}$. Since F is primitive, it is the universal relation, so $(a, (a_\omega)_\omega) \in \Gamma(F)$.

5.2. Banks–Elder–Willis (P_k) -closures. Theorem 4.34 yields a description of the (P_k) -closures of locally transitive subgroups of $\text{Aut}(T_d)$ that contain an involutive inversion, and therefore a characterization of the locally transitive universal groups. Recall that the (P_k) -closure of a subgroup $H \leq \text{Aut}(T_d)$ is

$$H^{(P_k)} = \{g \in \text{Aut}(T_d) \mid \text{for all } x \in V \text{ there exists } h \in H : g|_{B(x,k)} = h|_{B(x,k)}\}.$$

Combined with Corollary 4.18 the following partially answers the question for an algebraic description of a group’s (P_k) -closure in the last paragraph of [1].

THEOREM 5.31. *Let $H \leq \text{Aut}(T_d)$ be locally transitive and contain an involutive inversion. Then $H^{(P_k)} = \bigcup_k^{(l)}(F^{(k)})$ for some labelling l of T_d and $F^{(k)} \leq \text{Aut}(B_{d,k})$.*

PROOF. Let l and $F^{(k)} \leq \text{Aut}(B_{d,k})$ be as in Theorem 4.34. Then $H^{(P_k)} = \bigcup_k^{(l)}(F^{(k)})$.

Let $g \in \bigcup_k(F^{(k)})$ and $x \in V$. Since $\bigcup_1^{(l)}(\{\text{id}\}) \leq H$ there is $h' \in \bigcup_1^{(l)}(\{\text{id}\}) \leq H$ with $h'(x) = g(x)$, and since H is k -locally action isomorphic to $F^{(k)}$ there is $h'' \in H_x$ such that $\sigma_k(h'', x) = \sigma_k(g, x)$. Then $h := h'h'' \in H$ satisfies $g|_{B(x,k)} = h|_{B(x,k)}$.

Conversely, let $g \in H^{(P_k)}$. Then all k -local actions of g stem from elements of H . Given that $H \leq \bigcup_k(F^{(k)})$ by Theorem 4.34, we conclude that $g \in \bigcup_k(F^{(k)})$. □

COROLLARY 5.32. *Let $H \leq \text{Aut}(T_d)$ be closed, locally transitive and contain an involutive inversion. Then $H = \bigcup_k^{(l)}(F^{(k)})$ for some labelling l of T_d and an action $F^{(k)} \leq \text{Aut}(B_{d,k})$ if and only if H satisfies property (P_k) .*

PROOF. If $H = \bigcup_k^{(l)}(F^{(k)})$ then H satisfies property (P_k) by Proposition 4.7. Conversely, if H satisfies Property (P_k) then $H = \overline{H} = H^{(P_k)}$ by [1, Theorem 5.4] and the assertion follows from Theorem 5.31. □

Banks, Elder and Willis use certain subgroups of $\text{Aut}(T_d)$ with pairwise distinct (P_k) -closures to construct infinitely many, pairwise nonconjugate, nondiscrete simple subgroups of $\text{Aut}(T_d)$ via Theorem 2.1 and [1, Theorem 8.2]. For example, the group $\text{PGL}(2, \mathbb{Q}_p) \leq \text{Aut}(T_{p+1})$ qualifies by the argument in [1, Section 4.1]. Whereas $\text{PGL}(2, \mathbb{Q}_p)$ has trivial quasicycle given that it is simple, certain groups with nontrivial quasicycle always have infinitely many distinct (P_k) -closures.

PROPOSITION 5.33. *Let $H \leq \text{Aut}(T_d)$ be closed, nondiscrete, locally transitive and contain an involutive inversion. If also H has nontrivial quasicycle then H has infinitely many distinct (P_k) -closures.*

PROOF. We have $H^{(P_k)} = \bigcup_k(F^{(k)})$ by Theorem 5.31. Therefore, $H = \bigcap_{k \in \mathbb{N}} \bigcup_k(F^{(k)})$ by [1, Proposition 3.4(iii)]. If H had only finitely many distinct (P_k) -closures, the sequence $(H^{(P_k)})_{k \in \mathbb{N}}$ of subgroups of $\text{Aut}(T_d)$ would be eventually constant and equal to, say, $H^{(n)} = \bigcup_n(F^{(n)}) \geq H$. However, since H is nondiscrete, so is $\bigcup_n(F^{(n)})$ which thus has trivial quasicycle by Proposition 4.21. □

Banks, Elder and Willis ask whether the infinitely many, pairwise nonconjugate, nondiscrete simple subgroups of $\text{Aut}(T_d)$ they construct are also pairwise nonisomorphic as topological groups. By Proposition 4.17, this is the case if the said simple groups are locally transitive with distinct point-stabilizers, which can be determined from the original group's k -local actions thanks to Theorem 5.31.

THEOREM 5.34. *Let $H \leq \text{Aut}(T_d)$ be nondiscrete, locally permutation isomorphic to $F \leq \text{Sym}(\Omega)$ and contain an involutive inversion. Suppose that F is transitive and that every nontrivial subnormal subgroup of F_ω ($\omega \in \Omega$) is transitive on $\Omega \setminus \{\omega\}$. If $H^{(P_k)} \neq H^{(P_l)}$ for some $k, l \in \mathbb{N}$ then $(H^{(P_k)})^{+k}$ and $(H^{(P_l)})^{+l}$ are nonisomorphic.*

PROOF. In view of [1, Theorem 8.2], the groups $(H^{(P_k)})^{+k}$ and $(H^{(P_l)})^{+l}$ are nonconjugate. We show that they satisfy the assumptions of Proposition 4.17 which then implies

the assertion. It suffices to consider $H^{(P_k)}$. By Theorem 5.31, we have $H^{(P_k)} = U_k(F^{(k)})$ for some $F^{(k)} \leq \text{Aut}(B_{d,k})$. By virtue of Proposition 4.10, we may assume that $F^{(k)}$ satisfies (C). Since H is nondiscrete, so is $H^{(P_k)} = U_k(F^{(k)})$. Therefore, $F^{(k)}$ does not satisfy (D); see Proposition 4.12. Hence, in view of the local action of H and Proposition 4.31, the group $\pi_\xi F_{T_\omega}^{(k)}$ is nontrivial and thus transitive by Proposition 4.30 for all $\xi = (\omega_1, \dots, \omega_{k-1}) \in \Omega^{(k-1)}$ and $\omega \in \Omega \setminus \{\omega_1\}$. Now, let $x \in V(T_d)$. For every $\omega \in \Omega$ pick $\xi = (\omega_1, \dots, \omega_{k-2}, \omega) \in \Omega^{(k-1)}$. Let $y \in V(T_d)$ be such that $x = y_\xi$. Since $\pi_\xi F_{T_{\omega'}}^{(k)}$ is transitive for every $\omega' \in \Omega \setminus \{\omega_1\}$ we conclude that $(H^{(P_k)})^{+k}$ is locally 2-transitive at x . So Proposition 4.17 applies. \square

EXAMPLE 5.35. Theorem 5.34 applies to $\text{PGL}(2, \mathbb{Q}_p) \leq \text{Aut}(T_{p+1})$ for any prime p by Lemma 5.36 below. In fact, the local action is given by $\text{PGL}(2, \mathbb{F}_p) \curvearrowright \mathbb{P}^1(\mathbb{F}_p)$, point-stabilizers of which act like $\text{AGL}(1, p) = \mathbb{F}_p^* \ltimes \mathbb{F}_p \curvearrowright \mathbb{F}_p$. Retaining the notation of [1, Section 4.1], an involutive inversion in $\text{PGL}(2, \mathbb{Q}_p)$ is given by

$$\sigma := \begin{bmatrix} 0 & 1 \\ p & 0 \end{bmatrix} \quad \text{with } \sigma^2 = \begin{bmatrix} p & 0 \\ 0 & p \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Indeed, σ swaps the vertices v and \mathbf{L}_p .

LEMMA 5.36. *Let $F \leq \text{Sym}(\Omega)$ be 2-transitive. If $|\Omega| - 1$ is prime then every nontrivial subnormal subgroup of F_ω ($\omega \in \Omega$) acts transitively on $\Omega \setminus \{\omega\}$.*

PROOF. Since F_ω acts transitively on $\Omega \setminus \{\omega\}$, which has prime order, F_ω is primitive. So every nontrivial normal subgroup of F_ω acts transitively on $\Omega \setminus \{\omega\}$. Iterate. \square

EXAMPLE 5.37. The proof of Theorem 5.34 shows that the assumptions on F can be replaced with asking that $(H^{(P_k)})^{+k}$ be locally transitive with distinct point-stabilizers, which may be feasible to check in a given example.

For instance, let $F \leq \text{Sym}(\Omega)$ be transitive with distinct point-stabilizers. Assume that F preserves a nontrivial partition $\mathcal{P} : \Omega = \bigsqcup_{i \in I} \Omega_i$ of Ω and that it is generated by its block stabilizers, that is, $F = \langle \{F_{\Omega_i} \mid i \in I\} \rangle$.

Let $p : \Omega \rightarrow I$ be such that $\omega \in \Omega_{p\omega}$ for all $\omega \in \Omega$. Inductively define groups $F^{(k)} \leq \text{Aut}(B_{d,k})$ by $F^{(1)} := F$ and $F^{(k+1)} := \Phi_k(F^{(k)}, \mathcal{P})$, and check that

- (i) $C_{F^{(k)}}(\alpha, \Omega_i)$ is nonempty for all $\alpha \in F^{(k)}$ and $i \in I$,
- (ii) $C_{F^{(k)}}(\text{id}, \Omega_i)$ is nontrivial for all $i \in I$,
- (iii) $F^{(k+1)} \leq \Phi(F^{(k)})$, and
- (iv) $\pi_\xi F_{T_\omega}^{(k)} = F_{\Omega_{p\omega_{k-1}}}$ for all $\omega \in \Omega$ and $\xi = (\omega_1, \dots, \omega_{k-1}) \in \Omega^{(k-1)}$ with $\omega_1 \notin \Omega_{p\omega}$.

In particular, $F^{(k)}$ satisfies (C) but not (D) for all $k \in \mathbb{N}$. Set $H := \bigcap_{k \in \mathbb{N}} U_k(F^{(k)})$. By the above, H is nondiscrete and contains both $D(F)$ and $U_1(\{\text{id}\})$. Hence, Theorem 5.31 applies and we have $H^{(P_k)} = U_k(F^{(k)})$. From item (iii), we conclude that the $H^{(P_k)}$ ($k \in \mathbb{N}$) are pairwise distinct. Given that $(H^{(P_k)})^{+k}$ locally acts like F due to item (iv), the $(H^{(P_k)})^{+k}$ ($k \in \mathbb{N}$) are hence pairwise nonisomorphic.

5.3. A view on the Weiss conjecture. The Weiss conjecture states that there are only finitely many conjugacy classes of discrete, vertex-transitive, locally primitive subgroups of $\text{Aut}(T_d)$ for a given $d \in \mathbb{N}_{\geq 3}$. We now study the universal group construction in the discrete case and offer a new view on this conjecture. Under the additional assumption that each group contains an involutive inversion, it suffices to show that for every primitive $F \leq \text{Sym}(\Omega)$ there are only finitely many $\widetilde{F} \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) with $\pi\widetilde{F} = F$ and that satisfy (CD) in a minimal fashion; see Definition 5.42 and the discussion thereafter.

The following consequence of Theorem 5.31 identifies certain groups relevant to the Weiss conjecture as universal groups for local actions satisfying condition (CD).

COROLLARY 5.38. *Let $H \leq \text{Aut}(T_d)$ be discrete, locally transitive and contain an involutive inversion. Then $H = \bigcup_k^{(l)} (F^{(k)})$ for some $k \in \mathbb{N}$, a labelling l of T_d and $F^{(k)} \leq \text{Aut}(B_{d,k})$ satisfying (CD) that is isomorphic to the k -local action of H .*

PROOF. Discreteness of H implies property (P_k) for every $k \in \mathbb{N}$ such that stabilizers in H of balls of radius k in T_d are trivial. Then apply Theorem 5.31. \square

Therefore, the study of the class of groups given in Corollary 5.38 reduces to the study of subgroups $F \leq \text{Aut}(B_{d,k})$ ($k \in \mathbb{N}$) that satisfy (CD) and for which πF is transitive. By Corollary 4.15, any two conjugate such groups yield isomorphic universal groups. In this sense, it suffices to examine conjugacy classes of subgroups of $\text{Aut}(B_{d,k})$. This can be done computationally using the description of conditions (C) and (D) developed in Section 4.2, using, for example, [9].

EXAMPLE 5.39. Consider the case $d=3$. By [7, 31, 32], there are, up to conjugacy, seven discrete, vertex-transitive and locally transitive subgroups of $\text{Aut}(T_3)$. We denote them by $G_1, G_2, G_2^1, G_3, G_4, G_4^1$ and G_5 . The subscript n determines the isomorphism class of the vertex stabilizer, whose order is $3 \cdot 2^{n-1}$. A group contains an involutive inversion if and only if it has no superscript. The minimal order of an inversion in G_2^1 and G_4^1 is 4. See also [4]. By Corollary 5.38, the groups G_n ($n \in \{1, \dots, 5\}$) are of the form $\bigcup_k (F)$. We recover their local actions in Table 1. The list is complete for $k=2$, and for $k=3$ in the case of (CD).

The column labelled ‘i.c.c.’ records whether F admits an involutive compatibility cocycle. This can be determined in [9] and is automatic in the case of (CD). The group $\Pi(S_3, \text{sgn}, \{1\})$ of Proposition 4.25 admits an involutive compatibility cocycle z which we describe as follows. Suppose $\Omega := \{1, 2, 3\}$. Let $t_i \in \text{Sym}(\Omega)$ be the transposition that fixes i , and let $\tau_i \in \Pi(S_3, \text{sgn}, \{1\})$ be the element whose 1-local action is t_i everywhere except at b_i . Then $\Pi(S_3, \text{sgn}, \{1\}) = \langle \tau_1, \tau_2, \tau_3 \rangle$. Further, let $\kappa_i \in \Pi(S_3, \text{sgn}, \{1\}) \cap \ker \pi$ be the nontrivial element with $\sigma_1(\kappa_i, b_i) = e$. We then have $z(\tau_i, i) = \kappa_{i-1}$ and $z(\tau_i, j) = \tau_i \kappa_j$ for all distinct $i, j \in \Omega$, with cyclic notation.

The kernel K_2 is the diagonal subgroup of $\mathbb{Z}/2 \mathbb{Z}^{3 \cdot (3-1)} \cong \ker \pi_2 \leq \text{Aut}(B_{3,3})$. Using the above, we conclude that $G_1 = \bigcup_1(A_3)$, $G_2 = \bigcup_2(\Gamma(S_3))$, $G_3 = \bigcup_2(\Delta(S_3))$, $G_4 = \bigcup_3(\Gamma_2(\Pi(S_3, \text{sgn}, \{1\})))$ and $G_5 = \bigcup_3(\Sigma_2(\Pi(S_3, \text{sgn}, \{1\}), K_2))$.

TABLE 1. Conjugacy class representatives of subgroups F of $\text{Aut}(B_{3,2})$ and $\text{Aut}(B_{3,3})$ that satisfy (C) and project onto a transitive subgroup of S_3 .

Description of F	k	πF	$ F $	(C)	(D)	i.c.c.
$\Phi(A_3)$	2	A_3	3	yes	yes	yes
$\Gamma(S_3)$	2	S_3	6	yes	yes	yes
$\Delta(S_3)$	2	S_3	12	yes	yes	yes
$\Pi(S_3, \text{sgn}, \{0, 1\})$	2	S_3	24	yes	no	no
$\Pi(S_3, \text{sgn}, \{1\})$	2	S_3	24	yes	no	yes
$\Phi(S_3)$	2	S_3	48	yes	no	no
Description of F	k	$\pi_2 F$	$ F $	(C)	(D)	i.c.c.
$\Gamma_2(\Pi(S_3, \text{sgn}, \{1\}))$	3	$\Pi(S_3, \text{sgn}, \{1\})$	24	yes	yes	yes
$\Sigma_2(\Pi(S_3, \text{sgn}, \{1\}), K_2)$	3	$\Pi(S_3, \text{sgn}, \{1\})$	48	yes	yes	yes

QUESTION 5.40. Can the groups G_2^1 and G_4^1 be described as universal groups with prescribed local actions on edge neighbourhoods that prevent involutive inversions?

The long-standing Weiss conjecture [33] states that there are only finitely many conjugacy classes of discrete, vertex-transitive, locally primitive subgroups of $\text{Aut}(T_d)$ for a given $d \in \mathbb{N}_{\geq 3}$. Potočnic *et al.* [21] show that a permutation group $F \leq \text{Sym}(\Omega)$, for which there are only finitely many conjugacy classes of discrete, vertex-transitive subgroups of $\text{Aut}(T_d)$ that locally act like F , is necessarily semiprimitive, and conjecture the converse. Promising partial results were obtained in the same article as well as by Giudici and Morgan in [11].

Corollary 5.38 suggests restricting to discrete, locally semiprimitive subgroups of $\text{Aut}(T_d)$ containing an involutive inversion.

CONJECTURE 5.41. Let $F \leq \text{Sym}(\Omega)$ be semiprimitive. Then there are only finitely many conjugacy classes of discrete subgroups of $\text{Aut}(T_d)$ that locally act like F and contain an involutive inversion.

For a transitive permutation group $F \leq \text{Sym}(\Omega)$, let \mathcal{H}_F denote the collection of subgroups of $\text{Aut}(T_d)$ that are discrete, locally act like F and contain an involutive inversion. Then the following definition is meaningful by Corollary 5.38.

DEFINITION 5.42. Let $F \leq \text{Sym}(\Omega)$ be transitive. Define

$$\dim_{\text{CD}}(F) := \max_{H \in \mathcal{H}_F} \min\{k \in \mathbb{N} \mid \text{there exists } F^{(k)} \in \text{Aut}(B_{d,k}) \text{ with (CD) : } H = U_k(F^{(k)})\}$$

if the maximum exists and $\dim_{\text{CD}}(F) = \infty$ otherwise.

Given Definition 5.42, Conjecture 5.41 is equivalent to asserting that $\dim_{\text{CD}}(F)$ is finite whenever $F \leq \text{Sym}(\Omega)$ is semiprimitive. The remainder of this subsection is devoted to determining \dim_{CD} for certain classes of transitive permutation groups.

PROPOSITION 5.43. *Let $F \leq \text{Sym}(\Omega)$ be transitive. Then $\dim_{\text{CD}}(F) = 1$ if and only if F is regular.*

PROOF. If F is regular, then $\dim_{\text{CD}}(F) = 1$ by Proposition 4.13. Conversely, if $\dim_{\text{CD}}(F) = 1$ then $U_2(\Delta(F)) = U_1(F) = U_2(\Gamma(F))$. Hence, $\Gamma(F) \cong \Delta(F)$, which implies that F_ω is trivial for all $\omega \in \Omega$. That is, F is regular. \square

The next proposition provides a large class of primitive groups of dimension 2. It relies on the following relations between various characteristic subgroups of a finite group. Recall that the socle of a finite group is the subgroup generated by its minimal normal subgroups, which form a direct product.

LEMMA 5.44. *Let G be a finite group. Then the following assertions are equivalent.*

- (i) *The socle $\text{soc}(G)$ has no abelian factor.*
- (ii) *The solvable radical $O_\infty(G)$ is trivial.*
- (iii) *The nilpotent radical $\text{Fit}(G)$ is trivial.*

PROOF. If $\text{soc}(G)$ has no abelian factor then $O_\infty(G)$ is trivial: a nontrivial solvable normal subgroup of G would contain a minimal solvable normal subgroup of G which is necessarily abelian. Next, (ii) implies (iii) as every nilpotent group is solvable. Finally, if $\text{soc}(G)$ has an abelian factor then G contains a (minimal) normal abelian, hence nilpotent subgroup. Thus (iii) implies (i). \square

PROPOSITION 5.45. *Let $F \leq \text{Sym}(\Omega)$ be primitive, nonregular and assume that F_ω has trivial nilpotent radical for all $\omega \in \Omega$. Then $\dim_{\text{CD}}(F) = 2$.*

PROOF. Suppose that $F^{(2)} \leq \text{Aut}(B_{d,2})$ satisfies (C) and that the sequence

$$1 \longrightarrow \ker \pi \longrightarrow F^{(2)} \xrightarrow{\pi} F \longrightarrow 1$$

is exact. Fix $\omega_0 \in \Omega$. Then $\ker \pi \leq \prod_{\omega \in \Omega} F_\omega \cong F_{\omega_0}^d$. Since $F^{(2)}$ satisfies (C), we have $\text{pr}_\omega(\ker \pi) \trianglelefteq F_{\omega_0}$ for all $\omega \in \Omega$, and since F is transitive these projections all coincide with the same $N \trianglelefteq F_{\omega_0}$. Now consider $F_{T_\omega}^{(2)} = \ker \text{pr}_\omega|_{\ker \pi} \trianglelefteq \ker \pi$ for some $\omega \in \Omega$. Either $F_{T_\omega}^{(2)}$ is trivial, in which case $F^{(2)}$ has (CD), or $F_{T_\omega}^{(2)}$ is nontrivial. In the latter case, suppose $N_{\omega,\omega'} := \text{pr}_{\omega'} F_{T_\omega}^{(2)}$ is nontrivial for some $\omega' \in \Omega$. Then $N_{\omega,\omega'}$ is subnormal in F_{ω_0} as $N_{\omega,\omega'} \trianglelefteq N \trianglelefteq F_{\omega_0}$ and therefore has trivial nilpotent radical. The Thompson–Wielandt theorem [28, 34] (cf. [2, Theorem 2.1.1]) now implies that there is no $F^{(k)} \leq \text{Aut}(B_{d,k})$, $k \geq 3$, satisfying $\pi_2 F^{(k)} = F^{(2)}$ and (CD). Thus $\dim_{\text{CD}}(F) \leq 2$. Equality holds by Proposition 5.43. \square

Proposition 5.45 applies to $\text{Alt}(d)$ and $\text{Sym}(d)$, $d \geq 6$, whose point-stabilizers have nonabelian simple socle $\text{Alt}(d - 1)$. It also applies to primitive groups of O’Nan–Scott type (TW) and (HS), whose point-stabilizers have trivial solvable radical [6, Theorem 4.7B] and simple nonabelian socle [16], respectively.

EXAMPLE 5.46. By Example 5.39, we have $\dim_{\text{CD}}(S_3) \geq 3$. The article [7] shows that in fact $\dim_{\text{CD}}(S_3) = 3$.

To contrast the primitive case, we show that imprimitive wreath products have dimension at least 3, illustrating the use of involutive compatibility cocycles. Recall that for $F \leq \text{Sym}(\Omega)$ and $P \leq \text{Sym}(\Lambda)$ the wreath product $F \wr P := F^{|\Lambda|} \rtimes P$ admits a natural imprimitive action on $\Omega \times \Lambda$ with the partition $\bigsqcup_{\lambda \in \Lambda} \Omega \times \{\lambda\}$, namely $((a_\lambda)_\lambda, \sigma) \cdot (\omega, \lambda') := (a_{\sigma(\lambda')}\omega, \sigma\lambda')$.

PROPOSITION 5.47. *Let Ω and Λ be finite sets of size at least 2. Furthermore, let $F \leq \text{Sym}(\Omega)$ and $P \leq \text{Sym}(\Lambda)$ be transitive. Then $\dim_{\mathbb{C}D}(F \wr P) \geq 3$.*

PROOF. We define a subgroup $W(F, P) \leq \text{Aut}(B_{|\Omega \times \Lambda|, 2})$ that projects onto $F \wr P$, satisfies (C), does not satisfy (D) but admits an involutive compatibility cocycle. This suffices by Lemma 4.26. For $\lambda \in \Lambda$, let ι_λ denote the λ th embedding of F into $F \wr P = (\prod_{\lambda \in \Lambda} F) \rtimes P$. Recall the map γ from Section 4.4.1 and consider

$$\begin{aligned} \gamma_\lambda : F &\rightarrow \text{Aut}(B_{|\Omega \times \Lambda|, 2}), a \mapsto (\iota_\lambda(a), ((\iota_\lambda(a))_{(\omega, \lambda)}, (\text{id})_{(\omega, \lambda' \neq \lambda)})), \\ \gamma_\lambda^{(2)} : F &\rightarrow \text{Aut}(B_{|\Omega \times \Lambda|, 2}), a \mapsto (\text{id}, ((\text{id})_{(\omega, \lambda)}, (\iota_\lambda(a))_{(\omega, \lambda' \neq \lambda)})). \end{aligned}$$

Furthermore, let ι denote the embedding of P into $F \wr P$. We define

$$W(F, P) := \langle \gamma_\lambda(a), \gamma_\lambda^{(2)}(a), \gamma(\iota(\varrho)) \mid \lambda \in \Lambda, a \in F, \varrho \in P \rangle.$$

By construction, $W(F, P)$ does not satisfy (D). To see that $W(F, P)$ admits an involutive compatibility cocycle, we first determine its group structure. Consider the subgroups $V := \langle \gamma_\lambda(a) \mid \lambda \in \Lambda, a \in F \rangle$ and $\bar{V} := \langle \gamma_\lambda^{(2)}(a) \mid \lambda \in \Lambda, a \in F \rangle$. Then $W(F, P) = \langle V, \bar{V}, \Gamma(\iota(P)) \rangle$. Observe that $V \cong F^{|\Lambda|}$ and $\bar{V} \cong F^{|\Lambda|}$ commute, intersect trivially, and that $\Gamma(\iota(P))$ permutes the factors of each product. Hence,

$$W(F, P) \cong (V \times \bar{V}) \rtimes P \cong (F^{|\Lambda|} \times F^{|\Lambda|}) \rtimes P.$$

An involutive compatibility cocycle z of $W(F, P)$ may now be defined by setting

$$z(\gamma_\lambda(a), (\omega, \lambda')) := \begin{cases} \gamma_\lambda(a) & \lambda = \lambda' \\ \gamma_\lambda^{(2)}(a) & \lambda \neq \lambda' \end{cases}, z(\gamma_\lambda^{(2)}(a), (\omega, \lambda')) := \begin{cases} \gamma_\lambda^{(2)}(a) & \lambda = \lambda' \\ \gamma_\lambda(a) & \lambda \neq \lambda' \end{cases}$$

for all $\lambda \in \Lambda, a \in F$, and $z(\gamma(\iota(\varrho)), (\omega, \lambda)) := \gamma(\iota(\varrho))$ for all $\varrho \in P$. In fact, the map z extends to an involutive compatibility cocycle of $V \times \bar{V} \leq W(F, P)$ which in turn extends to an involutive compatibility cocycle of $W(F, P)$. □

Acknowledgements

The author is indebted to Marc Burger and George Willis for their support and the suggestion to define generalized universal groups. Thanks are also due to Luke Morgan and Michael Giudici for sharing their insight on permutation groups, and Michael Giudici for providing a proof of Lemma 4.29. Two anonymous referees' comments were also much appreciated.

A good part of this research was carried out during visits to The University of Newcastle, Australia, for the hospitality of which the author is thankful.

References

- [1] C. Banks, M. Elder and G. Willis, ‘Simple groups of automorphisms of trees determined by their actions on finite subtrees’, *J. Group Theory* **18**(2) (2015), 235–261.
- [2] M. Burger and S. Mozes, ‘Groups acting on trees: from local to global structure’, *Publ. Math. Inst. Hautes Études Sci.* **92**(1) (2000), 113–150.
- [3] P.-E. Caprace and A. Le Boudec, ‘Bounding the covolume of lattices in products’, *Compos. Math.* **155**(12) (2019), 2296–2333.
- [4] M. Conder and P. Lorimer, ‘Automorphism groups of symmetric graphs of valency 3’, *J. Combin. Theory Ser. B* **47**(1) (1989), 60–72.
- [5] D. Daykin, ‘On the rank of the matrix $f(A)$ and the enumeration of certain matrices over a finite field’, *J. Lond. Math. Soc. (2)* **1**(1) (1960), 36–42.
- [6] J. Dixon and B. Mortimer, *Permutation Groups*, Graduate Texts in Mathematics, 163 (Springer, New York, 1996).
- [7] D. Ž. Djoković and G. Miller, ‘Regular groups of automorphisms of cubic graphs’, *J. Combin. Theory Ser. B* **29**(2) (1980), 195–230.
- [8] A. Figà-Talamanca and C. Nebbia, *Harmonic Analysis and Representation Theory for Groups Acting on Homogenous Trees*, London Mathematical Society Lecture Note Series, 162 (Cambridge University Press, Cambridge, 1991).
- [9] GAP Group, GAP—Groups, Algorithms, and Programming, Version 4.8.7.
- [10] A. Garrido, Y. Glasner and S. Tornier, ‘Automorphism groups of trees: generalities and prescribed local actions’, in: *New Directions in Locally Compact Groups* (eds. P.-E. Caprace and M. Monod) (Cambridge University Press, Cambridge, 2018), Ch. 6, pp. 92–116.
- [11] M. Giudici and L. Morgan, ‘A class of semiprimitive groups that are graph-restrictive’, *Bull. Lond. Math. Soc.* **46**(6) (2014), 1226–1236.
- [12] M. Giudici and L. Morgan, ‘A theory of semiprimitive groups’, *J. Algebra* **503** (2018), 146–185.
- [13] A. Gleason, ‘Groups without small subgroups’, *Ann. of Math. (2)* **56** (1952), 193–212.
- [14] G. Károlyi, J. Kovács and P. Pálffy, ‘Doubly transitive permutation groups with abelian stabilizers’, *Aequationes Math.* **39**(2–3) (1990), 161–166.
- [15] B. Krön and R. Möller, ‘Analogues of Cayley graphs for topological groups’, *Math. Z.* **258**(3) (2008), 637.
- [16] M. Liebeck, C. Praeger and J. Saxl, ‘On the O’Nan–Scott theorem for finite primitive permutation groups’, *J. Aust. Math. Soc. Ser. A* **44**(03) (1988), 389–396.
- [17] M. Liebeck and J. Saxl, ‘On point stabilizers in primitive permutation groups’, *Comm. Algebra* **19**(10) (1991), 2777–2786.
- [18] R. G. Möller and J. Vonk, ‘Normal subgroups of groups acting on trees and automorphism groups of graphs’, *J. Group Theory* **15**(6) (2012), 831–850.
- [19] N. Monod, *Continuous Bounded Cohomology of Locally Compact Groups* (Springer, Berlin–Heidelberg, 2001).
- [20] D. Montgomery and L. Zippin, ‘Small subgroups of finite-dimensional groups’, *Proc. Natl. Acad. Sci. USA* **38**(5) (1952), 440–442.
- [21] P. Potočník, P. Spiga and G. Verret, ‘On graph-restrictive permutation groups’, *J. Combin. Theory Ser. B* **102**(3) (2012), 820–831.
- [22] C. Praeger, ‘Finite quasiprimitive graphs’, in: *Surveys in Combinatorics, 1997* (ed: R. A. Bailey) (Cambridge University Press, Cambridge, 1997), 65–85.
- [23] N. Radu, ‘A classification theorem for boundary 2-transitive automorphism groups of trees’, *Invent. Math.* **209**(1) (2017), 1–60.
- [24] D. Rattaggi, ‘Computations in groups acting on a product of trees: normal subgroup structures and quaternion lattices’, PhD Thesis, ETH Zurich, 2004.
- [25] J.-P. Serre, *Trees* (Springer, Berlin–Heidelberg–New York, 2003).
- [26] N. Sloane, ‘The on-line encyclopedia of integer sequences’, <https://oeis.org>.

- [27] S. Smith, 'A product for permutation groups and topological groups', *Duke Math. J.* **166**(15) (2017), 2965–2999.
- [28] J. Thompson, 'Bounds for orders of maximal subgroups', *J. Algebra* **14**(2) (1970), 135–138.
- [29] J. Tits, 'Sur le groupe des automorphismes d'un arbre', in: *Essays on Topology and Related Topics* (eds: A. Haefliger and R. Narasimhan) (Springer, Berlin–Heidelberg, 1970), 188–211.
- [30] S. Tornier, 'Groups acting on trees and contributions to Willis theory', PhD Thesis, ETH Zurich, 2018.
- [31] W. Tutte, 'A family of cubical graphs', *Math. Proc. Cambridge Philos. Soc.* **43**(4) (1947), 459–474.
- [32] W. Tutte, 'On the symmetry of cubic graphs', *Canad. J. Math.* **11** (1959), 621–624.
- [33] R. Weiss, 's-Transitive graphs', in: *Algebraic Methods in Graph Theory*, Colloq. Math. Soc. János Bolyai, 25 (North Holland, Amsterdam, 1978), 827–847.
- [34] H. Wielandt, *Subnormal Subgroups and Permutation Groups* (The Ohio State University, 1971).
- [35] H. Yamabe, 'A generalization of a theorem of Gleason', *Annals of Mathematics* **58** (1953), 351–365.

STEPHAN TORNIER, The University of Newcastle – SIPS,
University Drive, Callaghan, NSW 2308, Australia
e-mail: stephan.tornier@newcastle.edu.au