

# NEW HIGH-RESOLUTION OBSERVATIONS OF MOLECULAR GAS TOWARDS THE CENTER OF SPIRAL GALAXIES

*Observations and numerical simulations*

S.GARCIA-BURILLO

*Observatorio Astronómico Nacional*

*Campus Universitario, Apdo 1143, 28800 Alcalá de Henares, SPAIN*


We discuss the major results from a study of the molecular gas distribution and dynamics in the nuclear disks (ND) of a limited sample of nearby spirals: NGC891, NGC5907, NGC4565, NGC4013, NGC4321, M82, NGC3626 and NGC3593. The main objective is to search for observational evidences of secular evolution along the Hubble sequence in spirals and to study the specific role of bars in the process.

The 1–0 and 2–1 lines of  $^{12}\text{CO}$ , observed with the 30m single-dish and Bure interferometer, are the best tracer of neutral gas in ND. High-resolution and sensitivity requirements are fulfilled to study ND, typically of sizes  $r \sim 100 - 500pc$ .

Major axis position-velocity diagrams of edge-on spirals in the sample allows to trace the CO rotation curves and possible deviations. ND stand out as a distinct mass component, in addition to bulge, disk and halo. ND masses range from  $1-5 \times 10^9 M_{\odot}$  and their signature is imprinted on the steepest CO rotation curves (with no HI counterpart). Rapidly rotating ND are studied in the edge-on's NGC891 (García-Burillo and Guélin, 1995), NGC5907 (García-Burillo *et al.* 1997) and NGC4013 (Gómez de Castro and García-Burillo, 1997). ND host nuclear bars: evidences are discussed and modelled using bar-driven orbits. Bar signatures include: existence of forbidden velocities, strong streaming motions, lack of axisymmetry and out of the plane gas orbits connected to peanut bulges (see NGC4013).

Observations and models underline *secular evolution of rotation curves in spirals*: bars drive inwards gas by gravitational torques, changing mass distribution during galaxy lifetime. ND witness the intensity of these gas mass accretion episodes. In its trip towards the nucleus gas shifts from high energy  $x_1$  orbits to lower energy  $x_2$  orbits. When the fraction of mass in the ND reaches a minimum threshold a dynamically decoupled bar instability

develops. *Bars within bars* are a powerful mechanism to drive gas to the inner 100 pc of spirals.



*Figure 1.* We display the particle orbits for molecular clouds in the region where the bar instability develops, as they are seen, firstly, **a(top)**: from the frame rotating at  $\Omega=23 \text{ km s}^{-1} \text{ kpc}^{-1}$ , for the solution of a single slow pattern and **b(bottom)**: from the frame rotating at  $\Omega=23 \text{ km s}^{-1} \text{ kpc}^{-1}$  for the best-fit solution of a double pattern.

As an example we have studied the feasibility of a two independent patterns scenario in the disk of the double barred face-on spiral M100 (García-Burillo *et al.* 1997, submitted). Using the Bure array we mapped the  $^{12}\text{CO}(1-0)$  emission in M100's ND. Molecular gas is distributed in a two spiral arm structure starting from the end points of a nuclear bar ( $r = 600 \text{ pc}$ ) up to  $r = 1.2 \text{ kpc}$ , and a central non-resolved source ( $r < 100 \text{ pc}$ ). The kinematics of the gas indicates the existence of a steep rotation curve ( $v_{rot}=180 \text{ km s}^{-1}$  at  $r \sim 100 \text{ pc}$ ) and strong streaming motions characteristic of a trailing spiral wave inside corotation.

Interpretation of the CO observations is made in the light of a numerical model of the clouds hydrodynamics. Ballistic simulations analyse the gas response to a gravitational potential derived from a K plate. We found the best fit solution consisting of a fast pattern ( $\Omega_f=160 \text{ km s}^{-1} \text{ kpc}^{-1}$ ) for the nuclear bar (with corotation at  $R_{COR}^F=1.2 \text{ kpc}$ ) decoupled from the slow pattern of the outer bar+spiral ( $\Omega_f=23 \text{ km s}^{-1} \text{ kpc}^{-1}$ ) (with corotation at  $R_{COR}^S=8-9 \text{ kpc}$ ). Corotation of the fast pattern is overlapped with the oILR of the slow pattern, allowing an efficient fuelling of molecular gas towards the nucleus.

## References

- García-Burillo S., Guélin M., 1995, *A&A*, 299, 657  
 García-Burillo S., Guélin M., Neininger, N., 1997, *A&A*, 319, 450  
 Gómez de Castro, A.I., García-Burillo S., 1997, *A&A*, 322, 381  
 García-Burillo S., Sempere, M.J., Combes, F., Neri, R., 1997, *A&A*, submitted