a major risk factor for general hospital-based mortality with excess mortality due to acute T2DM, as well as other diabetes-related complications. Our study gives support for an aggressive multi-disciplinary approach to identify and treat T2DM to prevent diabetic, respiratory and vascular complications in all individuals with bipolar disorder.

- Forty L, Ulanova A, Jones L, Jones I, Gordon-Smith K, Fraser C, et al. Comorbid medical illness in bipolar disorder. Br J Psychiatry 2014; 205: 465–72.
- 2 Korszun A, Moskvina V, Brewster S, Craddock N, Ferrero F, Gill M, et al. Familiality of symptom dimensions in depression. Arch Gen Psychiatry 2004; 61: 468–74.
- 3 Schoepf D, Heun R. Bipolar disorder and comorbidity: Increased prevalence and increased relevance of comorbidity for hospital-based mortality during a 12.5-year observation period in general hospital admissions. J Affective Disorders 2014; 169: 170–8.

Dieter Schoepf, Department of Psychiatry and Psychotherapy, University of Bonn, Germany, **Reinhard Heun**, Department of Psychiatry, Radbourne Unit, Royal Derby Hospital, UK. Email: Dieter.Schoepf@ukb.uni-bonn.de

doi: 10.1192/bjp.206.6.522a

Authors' reply: We agree with Schoepf & Heun about the importance of control-sample characteristics in case—control comparisons and the need to consider the possibility of biased or unrepresentative findings. That is why, in our study, we used age as a covariate to take account of the younger age of controls compared with our mood disorder cases. We also discussed the potential limitations on generalisability caused by using a voluntary sample from a single ethnic group.

Schoepf & Heun describe a different methodology (with its own strengths and limitations) to examine a different issue, namely the influence of comorbidity on mortality. It is encouraging that the findings from the two different approaches are consistent in highlighting the clinical importance of comorbidity of physical illness with bipolar disorder. Psychiatrists and mental health services must ensure that patients with bipolar disorder have access to appropriate, high-quality physical, as well as psychiatric, care.

Liz Forty, Nick Craddock (on behalf of the authors), Institute of Psychological Medicine and Clinical Neurosciences, Cardiff University, UK. Email: craddockn@cardiff.ac.uk

doi: 10.1192/bjp.206.6.523

Advances



Volume 21, Issue 3
May 2015
http://apt.rcpsych.org

CONTENTS

New from CPD Online

Cochrane corner: First rank symptoms for schizophrenia?

Round the corner: First-rank symptoms: a first-rank diagnostic test?

Alex J. Mitchell

Immune abnormalities across psychiatric disorders: clinical relevance

Valeria Mondelli, Paola Dazzan & Carmine M. Pariante

What a psychiatrist needs to know about copy number variants George Kirov, Elliott Rees & James Walters

Reflection: Challenges and opportunities in psychiatry today Ketan Dipak Jethwa

The role of attachment in medically unexplained symptoms and long-term illness

Gwen Adshead & Elspeth Guthrie

The use of internet-mediated cross-sectional studies in mental health research

Alexandra Pitman, David P. J. Osborn & Michael B. King

Mindreading: Tolstoy's oak tree metaphor: depression, recovery and psychiatric 'spiritual ecology'

Jeremy Holmes

Best interests, mental capacity legislation and the UN Convention on the Rights of Persons with Disabilities

Brendan Kelly

Obsessive-compulsive disorder in young people

Tim I. Williams & Roz Shafran

Eating disorders in children and adolescents

Dasha Nicholls & Elizabeth Barrett

For further information on subscribing to Advances please contact:

Subscriptions Dept, Maney Publishing, Suite 1c, Joseph's Well, Hanover Walk, LEEDS LS3 1AB, UK. Tel: +44 (0)113 386 8168. Fax: +44 (0)113 386 8178. Email: subscriptions@maneypublishing.com
Order online: www.maneyonline.com/loi/apt

For US orders: Subscriptions Dept, Maney Publishing (USA) Inc, 1515 Market Street, Suite 1200, Philadelphia, PA 19102, USA. Tel: +1 (215) 854 6402. Fax: +1 (215) 569 0216. Email: NorthAmerica@maneypublishing.com