

# The origin of the dusty envelope around Betelgeuse

Xavier Haubois<sup>1</sup>, Barnaby Norris<sup>2</sup>, Peter G. Tuthill<sup>2</sup>,  
Christophe Pinte<sup>3</sup>, Pierre Kervella<sup>4</sup>, Julien Girard<sup>1</sup>, Guy Perrin<sup>4</sup>,  
Sylvestre Lacour<sup>4</sup>, Andrea Chiavassa<sup>5</sup> and S. T. Ridgway<sup>6</sup>

<sup>1</sup>European Organisation for Astronomical Research in the Southern Hemisphere, Casilla  
19001, Santiago 19, Chile; email: [xhaubois@eso.org](mailto:xhaubois@eso.org)

<sup>2</sup>Sydney Institute for Astronomy, School of Physics, University of Sydney, NSW 2006,  
Australia

<sup>3</sup>UJF-Grenoble 1 / CNRS-INSU, Institut de Planétologie et d'Astrophysique de Grenoble,  
UMR 5274, 38041 Grenoble, France

<sup>4</sup>LESIA, (UMR 8109), Observatoire de Paris, PSL, CNRS, UPMC, Univ. Paris-Diderot, 5  
place Jules Janssen, 92195 Meudon, France

<sup>5</sup>Laboratoire Lagrange, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS,  
Boulevard de l'Observatoire, CS 34229, 06304 Nice Cedex 4, France

<sup>6</sup>National Optical Astronomy Observatory, P.O. Box 26732, Tucson, AZ 85726-6732, USA

**Abstract.** The origin of red supergiant mass loss still remains to be unveiled. Characterising the formation loci and the dust distribution in the first stellar radii above the surface is key to understand the initiation of the mass loss phenomenon. Polarimetric interferometry observations in the near-infrared allowed us to detect an inner dust atmosphere located only 0.5 stellar radius above the photosphere of Betelgeuse. We modelled these observations and compare them with visible polarimetric measurements to discuss the dust distribution properties.

**Keywords.** Convection-techniques: interferometric- stars: fundamental parameters- infrared: stars- stars: individual: Betelgeuse

---

## Summary

Depending on their characteristics and composition, dust grains could interact with the stellar radiation from red supergiants and trigger mass loss via a dust wind. We observed Betelgeuse using the NACO/SAMPol instrument and detected a polarizing structure at 1.5 stellar radius that we modelled as a thin dust shell. The dust-shell-to-stellar-disk flux ratio can be modelled as the fraction of the stellar light that is scattered on the grains of the dust shell. We fit the wavelength variation of this fraction using a Mie scattering routine. Parameters of this model are the grain radius and density, which is translated in a dust shell mass. We derived similar grain sizes of about 300 nm and masses of about  $10^{-10} M_{\odot}$  for three dust species (enstatite  $MgSiO_3$ , forsterite  $Mg_2SiO_4$  and alumina  $Al_2O_3$ ). Extrapolating to the visible wavelengths, we can compare our radiative transfer modelling to the SPHERE/ZIMPOL data reported in Kervella *et al.* (2016). A preliminary analysis shows that the forsterite and enstatite models match the data better than the alumina dust except in the V-band filter. More spectral filters in the visible should be obtained to infirm/confirm this hint on the dust composition.

## Reference

Kervella, P., Lagadec, E., Montargès, M., *et al.* 2016, *A&A*, 585, A28