

APERTURE SYNTHESIS MAPS OF NH₂D AND CH₃OD LINES TOWARD ORION-KL: THE ORIGIN OF NH₃ AND CH₃OH

S. SAITO¹, H. MIKAMI², S. YAMAMOTO², Y. MURATA³, AND R. KAWABE⁴

¹Institute for Molecular Science, ²Department of Astrophysics, Nagoya University, ³The Institute of Space and Astronautical Science, ⁴Nobeyama Radio Observatory

ABSTRACT The $1_{11}-1_{01+}$ transition of NH₂D and the 2_1-2_0 E transition of CH₃OD were mapped toward Orion-KL with the Nobeyama Millimeter Array. The synthesized beamwidth is 4" to 5". NH₂D and CH₃OD are mainly distributed over the peak-intensity regions of NH₃ and CH₃OH in Orion A, respectively. These results suggest that "most" of the gas-phase ammonia and methanol in the region of Orion-KL originate from dust grains.

INTRODUCTION

The high abundance of deuterated isotopes of interstellar molecules has been a topic of extreme interest in interstellar chemistry. Large deuterium enhancement has been found in the relatively hot (T=50-150 K) region of Orion-KL: [HDCO]/[H₂CO]=0.01-0.03 (Loren and Wootten, 1985), [NH₂D]/[NH₃]=0.03 (Walmsley et al. 1987), [HDO]/[H₂O]=0.001 (Petuchowski and Bennett, 1988), [CH₃OD]/[CH₃OH]=0.01-0.06 (Mauersberger et al. 1988). The abundance ratios of the deuterated species to the parent species are more than two orders of magnitude larger than the interstellar D/H ratio. The large deuterium fractionation suggests that the molecules are produced at temperatures of 10 K or less (Dalgarno and Lepp, 1984).

So as to determine the detailed distribution of deuterated species in the hot region, we mapped the lines of NH₂D and CH₃OD toward Orion-KL with the Nobeyama Millimeter Array.

RESULTS AND DISCUSSION

Observations were made during January to April, 1992 (3 days). The observed lines are the $1_{11}-1_{01+}$ transition of NH₂D at 110.153599 GHz and the 2_1-2_0 E transition of CH₃OD at 110.262640 GHz. The synthesized beam sizes were 5" x 4" for NH₂D and 4" x 4" for CH₃OD. The integrated intensity maps are shown in Fig. 1. A comparison between the distribution of the NH₃, J, K=1, 1 line (Murata et al. 1990) and that of NH₂D shows that the distribution of NH₂D corresponds to the main peak region of NH₃ at 5.4" south by southwest of IRC2, the hot core region. The distribution of CH₃OD also corresponds to the main peak region of CH₃OH (Plambeck and Wright, 1988) midway between IRC4 and IRC5, the north side of the compact ridge.

Plambeck and Wright (1987), and Walmsley et al. (1987) suggest that the highly deuterium-fractionated species are formed on dust grains which memorize the cold conditions of the molecular cloud before the young star, IRC2, switched on, and we are observing nonsteady-state conditions of the evaporated gases in the region. The present result is consistent with their model and, furthermore, implies that "most" of the gas-phase NH_3 and CH_3OH in the Orion-KL region originate from dust grains. Finally, it must be noted that the distributions of $(\text{CH}_3)_2\text{O}$ and HCOOCH_3 (Mikami et al. 1992) also coincide with the main peak-intensity region of CH_3OH and, as a result, $(\text{CH}_3)_2\text{O}$ and HCOOCH_3 in the Orion-KL region may also originate from the same dust grains from which methanol is evaporated.

In conclusion, it has been demonstrated that a detailed study on the distributions of deuterated species is a method to get an insight into the mechanism for interstellar molecular production whether it be via gas-phase ion-molecule or dust-grain related reactions.

REFERENCES

- Dalgarno, A. and Lepp, S. 1984, *Ap. J.*, **287**, L47.
 Loren, R. B. and Wootten, A. 1985, *Ap. J.*, **299**, 947.
 Mauersberger, R., Henkel, C., Jacq, T., and Walmsley, C. M. 1988, *Astr. Ap.*, **194**, L1.
 Mikami, H., et al. 1992, to be published.
 Murata, Y., Kawabe, R., Ishiguro, M., Morita, K., Kasuga, T., Takano, T., and Hasegawa, T. 1990, *Ap. J.*, **359**, 125.
 Petuchowski, S. J. and Bennett, C. L. 1988, *Ap. J.*, **326**, 376.
 Plambeck, R. L. and Wright, M. C. H. 1987, *Ap. J.*, **317**, L101.
 Plambeck, R. L. and Wright, M. C. H. 1988, *Ap. J.*, **330**, L61.
 Walmsley, C. M., Hermsen, W., Henkel, C., Mauersberger, R., and Wilson, T. L. 1987, *Astr. Ap.*, **172**, 311.




Fig. 1. The total integrated maps of $\text{NH}_2\text{D}(1_{11}-1_{01+})$ with the peak flux of $0.66 \text{ Jy beam}^{-1}$ (solid contour) and $\text{CH}_3\text{OD}(2_{1-2_0 \text{ E}})$ with $0.58 \text{ Jy beam}^{-1}$ (dashed contour). The lowest contour and the contour interval are 2σ .