

quite different. The 19th and early 20th century psychiatrist was faced with two common conditions which caused a great deal of work – general paresis (GPI) and epilepsy – stimulating clinical observation and research. It was noted that: (a) some psychotics improved when they had a fever; (b) some psychotics improved after a spontaneous epileptic fit; and (c) at post-mortem, epileptic brains and schizophrenic brains differed greatly, suggesting some antagonism between the schizophrenic and the epileptic process. The first point led Wagner-Jauregg, a Viennese contemporary of Freud, on a long search for artificial fever therapy; in 1917 he found that malarial infection would cure GPI, and he received a Nobel prize in 1927. The second and third led Meduna in Budapest in 1935 to induce fits with metrazol, with therapeutic success. But his method was unpleasant for patients and difficult to control. Cerletti had been studying experimental epilepsy in dogs using an electrical stimulus; with Bini, he adapted the stimulus for man and so produced a painless and easily controllable variant of Meduna's treatment. ECT is part of the history of epileptic studies, and its understanding and that of epilepsy march together.

JOHN CRAMMER

*Institute of Psychiatry  
De Crespigny Park  
London SE5 8AF*

#### Is Castration Too "Barbarous" for Rapists?

SIR: I fear that Salzman (*Journal*, August 1988, 153, 270) is suffering from the illusion that motivation for recurrent sexual offending and rape is purely sexual. Often the apparently sexually motivated acts, which he attributes to "psychosexual malignancy", are in fact expressions of a deeper, more complex and less obvious psychopathology. Indeed, such pathology may still be expressed as serious aggression after libido has been artificially reduced.

Furthermore, even individuals whose main problem is deemed to be hypersexuality, and perhaps therefore those he believes most likely to respond to surgical castration, are probably those least likely to agree to such treatment. Indeed, individuals who might agree to voluntary sacrificial surgical castration to justify their liberation from detention may be those least helped by it and most in need of rather wider and more subtle treatments.

However, when libidinal suppression is required, the currently available, equally dependable, but reversible 'chemical castration' using hormonal implants already has an accepted role as an adjunct to the overall treatment of certain sexual offenders. Indeed, there is carefully controlled provision for

such treatment under Section 57 of the Mental Health Act 1983.

Surely the use of presently available treatments rather than radical, but not magical, surgical castration will result in the continuation of a more considered overall approach to our patients and also less iatrogenic psychological morbidity in those whose ongoing mental stability is, after all, critical to both their success and the safety of others beyond conditions of detention.

CLIVE J. MEUX

*Bethlem Royal Hospital  
Monks Orchard Road  
Beckenham  
Kent BR3 3BX*

#### Females and Caring

SIR: I read with interest the recent review article by Morris (*Journal*, August 1988, 153, 147–156) concerning factors affecting the emotional wellbeing of the caregivers of dementia sufferers. The different approach and strategies that men have to caring is described by Zarit *et al* (1986).

However, the evidence at present reveals that females receive less statutory help than males when caring for an elderly relative. A study of carers found that 4% of mothers, 20% of wives and 24% of daughters received home help support, while 95% of caring sons and 68% of husbands received this service (Equal Opportunities Commission, 1982). It could be said that these figures simply reflect the fact that men are more willing to organise and accept help, but my concern is that they are a reflection of society's basic assumption that women can cope with caring whatever the burden. Do we as professionals become more aware of the burdens of caring when the carer is male?

MIRANDA CONWAY  
TONY CANTWELL

*Department of Psychiatry  
Whiston Hospital*

#### References

- EQUAL OPPORTUNITIES COMMISSION (1982) *Caring for the Elderly and Handicapped*. London: Equal Opportunities Commission.  
ZARIT, S. H., TODD, P. A. & ZARIT, J. M. (1986) Subjective burden of husbands and wives as caregivers: a longitudinal study. *Gerontologist*, 26, 260–266.

#### Delusional Depression in Nineteenth Century Scotland

SIR: It is encouraging to find two serious studies of the history of psychiatry in the August edition of the *British Journal of Psychiatry*. It is surely a sign of the present health of psychiatry as a speciality that it is