

Extensive spectroscopic and photometric study of HD 25558, a long orbital-period binary with two SPB components

Á. Sódor^{1,2}, P. De Cat¹, D. J. Wright^{1,3}, C. Neiner⁴, M. Briquet⁵,
 R. J. Dukes⁶, F. C. Fekel⁷, G. W. Henry⁷, M. H. Williamson⁷,
 M. W. Muterspaugh⁷, E. Brunsden⁸, K. R. Pollard⁸, P. L. Cottrell⁸,
 F. Maisonneuve⁸, P. M. Kilmartin⁸, J. M. Matthews⁹, T. Kallinger¹⁰,
 P. G. Beck¹¹, E. Kambe¹², C. A. Engelbrecht¹³, R. J. Czanik¹⁴,
 S. Yang¹⁵, O. Hashimoto¹⁶, S. Honda^{16,17}, J.-N. Fu¹⁸,
 B. Castanheira¹⁹, H. Lehmann²⁰, N. Behara²¹, H. Van Winckel¹¹,
 S. Scaringi¹¹, J. Menu¹¹, A. Lobel¹, P. Lampens¹, and P. Mathias²²

¹Royal Observatory of Belgium, Brussel, Belgium

²Konkoly Observatory, Hungarian Academy of Sciences, Budapest, Hungary

³School of Physics, University of New South Wales, Sydney, Australia

⁴LESIA, CNRS, Observatoire de Paris, UPMC, Université Paris Diderot, Meudon, France

⁵Institut d'Astrophysique et de Géophysique, Université de Liège, Liège, Belgium

⁶Department of Physics and Astronomy, The College of Charleston, Charleston, SC, USA

⁷Center of Excellence in Information Systems, Tennessee State University, Nashville, TN, USA

⁸Department of Physics and Astronomy, University of Canterbury, Christchurch, New Zealand

⁹Dept. of Physics and Astronomy, University of British Columbia, Vancouver, BC, Canada

¹⁰Institute for Astronomy, University of Vienna, Vienna, Austria

¹¹Instituut voor Sterrenkunde, K. U. Leuven, Leuven, Belgium

¹²Okayama Astrophysical Observatory, National Astronomical Observatory, Okayama, Japan

¹³Department of Physics, University of Johannesburg, Johannesburg, South Africa

¹⁴Dept. of Phys., Potchefstroom Campus, North-West University, Potchefstroom, South Africa

¹⁵Department of Physics and Astronomy, University of Victoria, Victoria, BC, Canada

¹⁶Gunma Astronomical Observatory, Takayama-mura, Agatsuma, Gunma, Japan

¹⁷Kwasan Observatory, Kyoto University, Kyoto, Japan

¹⁸Department of Astronomy, Beijing Normal University, Beijing, China

¹⁹Department of Astronomy, The University of Texas at Austin, Austin, TX, USA

²⁰Thüringer Landessternwarte Tautenburg, Tautenburg, Germany

²¹Université Libre de Bruxelles, Bruxelles, Belgium

²²Université de Toulouse, UPS-OMP, IRAP, CNRS, Toulouse, France

Abstract. We carried out an extensive photometric and spectroscopic investigation of the SPB binary, HD 25558 (see Fig. 1 for the time and geographic distribution of the observations). The ~ 2000 spectra obtained at 13 observatories during 5 observing seasons, the ground-based multi-colour light curves and the photometric data from the *MOST* satellite revealed that this object is a double-lined spectroscopic binary with a very long orbital period of about 9 years. We determined the physical parameters of the components, and have found that both lie within the SPB instability strip. Accordingly, both components show line-profile variations consistent with stellar pulsations. Altogether, 11 independent frequencies and one harmonic frequency were identified in the data. The observational data do not allow the inference of a reliable orbital solution, thus, disentangling cannot be performed on the spectra. Since the lines of the two components are never completely separated, the analysis is very complicated. Nevertheless, pixel-by-pixel variability analysis of the cross-correlated line profiles was successful, and we were able to attribute all the frequencies to the primary or secondary component. Spectroscopic and photometric mode-identification was also performed for several of these frequencies of both binary components. The spectroscopic mode-identification results suggest that the inclination

and rotation of the two components are rather different. While the primary is a slow rotator with ~ 6 d rotation period, seen at $\sim 60^\circ$ inclination, the secondary rotates fast with ~ 1.2 d rotation period, and is seen at $\sim 20^\circ$ inclination. Our spectropolarimetric measurements revealed that the secondary component has a magnetic field with at least a few hundred Gauss strength, while no magnetic field was detected in the primary.

The detailed analysis and results of this study will be published elsewhere.

Keywords. line: profiles, stars: binaries: general, stars: oscillations (including pulsations), stars: rotation, stars: variables: other




Figure 1. Time and geographic distribution of the spectroscopic and photometric observations of HD 25558. The main campaign was organised in the 2010/11 season.

Acknowledgements. Á.S. acknowledges support of the Belgian Federal Science Policy (project M0/33/029) and of the Eötvös Scholarship from the Hungarian Scholarship Board Office. M.B. is F.R.S.-FNRS Postdoctoral Researcher, Belgium. T.K. acknowledges financial support from the Austrian Science Fund (FWF P23608). The observational data we used in this study were obtained with the following telescopes and instruments: 2-m AST (T13) and 75-cm APT (T5) at Fairborn Observatory, McLellan tel.+HERCULES at Mt John Observatory, Otto Struve tel. at McDonald Observatory, 1.9-m tel.+GIRAFFE and 50-cm tel. at SAAO, Mercator tel.+HERMES at La Palma, 2.2-m tel. at Xinglong Observatory, 1.5-m tel.+GAOES at GAO, Alfred Jensch tel. at Karl Schwarzschild Observatory, 1.9-m tel.+HIDES at OAO, 1.2-m tel.+McKellar at DAO, Bernard Lyot tel.+NARVAL at Obs. Pic du Midi, CFHT+ESPADOnS at Hawaii, Euler tel.+CORALIE at La Silla, the Canadian *MOST* satellite.