

An evidence-based framework to culturally adapt cognitive behaviour therapy

Shanaya Rathod^{1*}, Peter Phiri¹ and Farooq Naeem²

¹*Southern Health NHS Foundation Trust, UK*

²*Centre for Addiction and Mental Health, Toronto, Canada and University of Toronto, Toronto, ON, Canada*

Abstract. Cognitive behaviour therapy (CBT) in its current form might not be applicable in non-Western cultures. Differences between Western and non-Western cultures have been reported widely. Most psychotherapeutic interventions have been developed in the West and are underpinned by the values that might be specific only to Western culture. To adapt CBT, we need to understand whether the concepts associated with CBT may conflict with cultural beliefs, identifying barriers to the success of the therapy using the views of experts by experience, public and practitioners. This paper discusses the process, foci and framework of cultural adaptation of CBT. We describe an evidence-based framework for adapting CBT for individuals from non-Western cultures that will benefit clinicians who practise CBT and individuals from different cultural backgrounds.

Key words: cultural adaptation, culture, evidence-based practice, CBT

Introduction

Cognitive behaviour therapy (CBT) is the most widely recommended psychological intervention for mental health conditions in the United Kingdom [e.g. National Institute of Health and Care Excellence (NICE)], in America [e.g. American Psychiatric Association (APA) Steering Committee on Practice Guidelines] and many other countries.

However, the issues around the use of CBT in different countries and cultural groups are twofold. Firstly, literature confirms that the rates of recruitment, retention and engagement of ethnic minority participants in most randomized controlled trials (RCTs) of CBT (Rathod *et al.*, 2005) are low in high and middle income countries (HMIC).¹ Therefore, there is limited evidence of its effectiveness in minority communities in these countries.

*Author for correspondence: Professor Shanaya Rathod, Clinical Trials Facility, Tom Rudd Unit, Moorgreen Hospital, Botley Road, West End, Southampton SO30 3JB. Visiting Professor, University of Portsmouth. (email: shanayarathod@nhs.net)

¹The World Bank has used income categories to classify countries into high, middle and low, with the middle income category further subdivided into lower middle or upper middle income. For the purposes of this paper, we use the terms high and middle income (to include the upper middle income) countries (HMIC) and low and middle income (to include lower middle income countries) countries (LMIC).

Secondly, the majority of trials of the effectiveness of CBT have been conducted in HMIC. Therefore, the evidence base of the effectiveness of CBT in LMIC¹ populations is still developing. Further literature comes from qualitative studies (Scorzelli and Scorzelli, 1994) that demonstrate that principles underlying cognitive therapy can conflict with the values and beliefs of different cultural groups in the LMIC.

The last two decades have seen a trend towards cultural adaptation of interventions including CBT for ethnic minority people in the HMIC and the LMIC (Naeem *et al.*, 2010; Rathod *et al.*, 2013; Rathod *et al.*, 2015; Rathod *et al.*, 2017a,b). Meta-analyses also confirm that culturally adapted interventions are effective, but it is not known what components of the adaptation have maximum impact in engaging people and improving outcomes (Rathod *et al.*, 2017a). It is understandable that some adjustments for minority communities in HMIC may differ from the adjustments required for the majority populations in LMIC and this may present challenges.

This paper discusses the need for adaptations of CBT despite the challenges, highlighting available adaptation frameworks, and then we describe the evidence-based adaptation framework for CBT developed by our group. We do not propose to present a systematic review or critique of all available adaptation frameworks as it does not fit the remit of this paper. We aim to orientate the CBT practitioner to the available options and generate interest in understanding adaptation techniques relevant to practising personalized CBT with individuals from different cultures.

Why adapt cognitive behaviour therapy?

Cognitive behaviour therapy was developed by Beck as a structured, present-oriented therapy for depression. It was designed to solve current problems by targeting and modifying unhelpful thinking patterns and belief systems using the cognitive model (Beck, 1964). Automatic thoughts and schemas were explained by Clarke and colleagues (Clark *et al.*, 1999). Over time, benefits of CBT have been realized for many conditions like anxiety disorders, borderline personality disorder and psychosis through case series, RCTs and meta-analyses.

In CBT, the therapist explores the person's beliefs and experiences to elicit cognitive biases, emotions and behaviours to develop a formulation and treatment plan. Culture influences an individual's life experiences, belief systems and assumptions. As a result, it impacts on the psychopathology of mental illness. Therefore, it follows that culture shapes beliefs about health and illness, pathways into care, and trust in services.

Cultural differences in life philosophies (Okasha and Okasha, 2000) are well documented. The risk of continuing to practise CBT in different cultural groups without adaptation can be disengagement and poor outcomes of therapy (Rathod *et al.*, 2005). Additionally, it may affect the therapeutic relationship with the therapist which is central to any therapeutic intervention, again, leading to poor outcomes. The therapist may feel relatively incompetent (Rathod *et al.*, 2015) as a consequence.

CBT is an evidence-based, pragmatic, collaborative therapy that emphasizes the critical role of cognitions, emotions and behaviours and therefore lends itself to adaptation while maintaining fidelity to the core CBT techniques (Rathod *et al.*, 2015). Often, people from different cultural backgrounds struggle to trust the health systems, mainly mental health




Figure 1. Role of culture in CBT model. Reprinted with permission from Rathod *et al.* (2015).

services (Schwarzbaum, 2004), especially in developed countries. The fundamental principles of CBT based on engagement and working together, can help diffuse the mistrust if the therapy appeals to the individual’s cultural values.

A further argument is that CBT addresses relationships and not only cognitions and beliefs. The cognitive model of CBT suggests that the focus of therapy is on an individual’s relationship (1) with self, (2) with others and (3) with the future. In many Eastern cultures family and community play a central role and are often sources of support, although changing values in many non-Western cultures might mean the nature of relationships is changing. Therefore CBT lends itself to successful adaptation, using the family and community networks as collaborators in the process where appropriate.

Frameworks for cultural adaptations of interventions

Any attempts to adapt therapy to culture have challenges. There could be a tendency to over-generalize and stereotype to cultures and subcultural groups. Maintaining fidelity to the core intervention while adapting to culture is an important consideration. Despite the challenges, local adaptations are occurring globally, based on therapists' experiences and local population needs with some benefits. Examples include studies by Carter and colleagues (2003), Hinton and colleagues (2004), Kubany and colleagues (2003), Patel and colleagues (2007), Rahman *et al.* (2008), Rojas *et al.* (2007) and Nagel *et al.* (2009).

Numerous adaptation frameworks and guidance have been developed over time to aid this process. The first adaptation framework was published by Bernal and colleagues (1995), using Latino participants and based on the ecological validity model for cultural adaptation. This framework consists of eight elements: language, persons, metaphors, content, concepts, goals, methods, and context.

Bernal's model was further developed and adapted by Domenech Rodríguez and Weiling (2004), and then by Hwang and colleagues (2006) to extend its usability in different groups. Adaptation frameworks were developed by Tseng (1999) describing the influence of culture on therapies. Resnicow and colleagues (2000) described two levels of cultural adaptation: surface structure adaptations and deep structure adaptations, while Leong and Lee (2006) developed an integrative and multi-dimensional cultural accommodation model.

Lau (2006), and later Barrera and Castro (2006), developed adaptations consisting of systematic phases and a heuristic framework with specific emphasis on developing mechanisms that would allow for both measuring and evaluating the impact of cultural adaptations using specific outcome measures.

The models described above have similarities in the dimensions of adaptation, but differ in the level of detail, including the different cultural groups for which they were described. They have limitations in relation to cultures (Rathod *et al.*, 2015) other than Asian and Hispanic groups. Hays and Iwamasa (2006) attempted to address this. However, none of the previous adaptation frameworks has been tested through RCTs.

An evidence-based methodology to cultural adaptation of CBT

Our group has developed an adaptation framework that can be applied to CBT in different cultural groups. The adapted intervention has been evaluated through randomized trials (Naeem *et al.*, 2010; Naeem *et al.*, 2015a,b; Rathod *et al.*, 2013). Currently, this methodology and framework is being used to adapt CBT in Morocco (Fatema-Zahra El Rhermoul *et al.*, 2018), the Middle East and China (Li *et al.*, 2017).

The adaptation framework has been developed using a staged process. The first step has involved information gathering from literature and used the qualitative methodology, underpinned by the ethnographic approach. The process has ensured a co-production model with experts by experience, carers, lay members of minority communities and clinicians to develop the adaptation framework that is relevant. Emergent themes have informed development of the guidance used to deliver culturally adapted CBT that is personalized to an individual's culture. The therapy material or manual is then translated and culturally adapted in light of the guidance developed from the qualitative analyses and existent literature. The

Table 1. Staged process of developing adaptation framework (Naeem et al., 2016)

Stage 1: Review of previous literature and discussions with field experts, followed by information gathering using qualitative methods from patients, carers, laypersons, therapists, mental health practitioners and service managers concerning their experiences and views
Stage 2: Producing guidance on adapting the CBT manual based on the information
Stage 3: Translation and adaptation of therapy material/manual
Stage 4: Field testing the adapted CBT manual through an RCT and further refinement of guidelines

adapted therapy is field tested and further refinements are made based on the findings (Naeem et al., 2016).

Foci of adaptation

Adaptation in our work has focused on three fundamental areas of delivery, which we refer to as the ‘Triple-A’ principle:

- I. Awareness of relevant cultural issues and preparation for therapy. This can be further subdivided into: (a) culture and culture-related issues including religion and spirituality, family and community, and language and communication; (b) system and environmental aspects including individual capacity and circumstances, systems of support, services, and helpseeking pathways into care; and (c) cognitive biases and unhelpful beliefs that are directly related to the problem and its treatment.
- II. Assessment and engagement.
- III. Adjustments in therapy.

These areas and sub-areas are further discussed below in the framework of adaptation.

Framework for cultural adaptation of CBT

The process of adaptation starts with preparation for the therapy, i.e. pre-engagement phase as access to therapy, its delivery and most importantly availability for a given culture is as important as the actual modification of the intervention. The adaptation framework then progresses to engagement and assessment, followed by the therapy using relevant cultural adaptations to improve outcomes.

Rathod and colleagues (2015) have developed a framework that encompasses the Triple-A principles for cultural adaptation of CBT focusing on the following aspects:

- (i) Philosophical orientation;
- (ii) Practical considerations of societal and health system-related factors;
- (iii) Technical adjustments of methods and skills; and
- (iv) Theoretical adaptations of concepts.

We have developed this adaptation model based on our work, using descriptions initially described by Tseng and colleagues (2005) and modifying them. The key principle is that every individual has a unique culture that is influenced by their wider culture, sub-culture and further developed through unique life experiences. There must be flexibility in applying the




Figure 2. Oscillation of an individual's cultural values: Shanaya Rathod and David Kingdon (2009). CBT across cultures. *Psychiatry* 8, 370–371. Copyright ©2009 Elsevier Ltd; printed with permission.

culturally adapted therapy, and clinicians should be aware of their own biases, and propensity to stereotype when working with people from minority groups. Below we briefly describe the framework that is explained in detail with case examples in the manual (Rathod *et al.*, 2015).

Philosophical orientation or differences in world view

The fundamental view of life can be different between individuals and different cultures to a degree, and this can affect perceptions of health and illness, and relationship with services and professionals, including treatment and goals of therapy. Rathod and colleagues (2015) describe philosophical orientation along the following constructs:

- a. *Acculturation* – acculturation is the process of adjustment that involves integration of the original and new cultural values and attitudes. This can occur during the period of prolonged contact with the host culture after immigration or internal movement in countries. The blending of cultural values is often a gradual process (Garcia and Zea, 1997), and can occur over several generations. The process of acculturation can result in integration, assimilation, separation, and/or marginalization – describing different trajectories of individuals influencing behaviours when interacting with others (Berry, 2005).

An individual can present with a mixture of different attitudes in different phases of life, through the influences of positive and negative life experiences and other factors. Figure 2 describes this. Understanding of an individual's level of acculturation helps to

understand how to engage with them and develop explanations in therapy based on their value systems.

Case example

Anish, a 30-year-old IT specialist, had been in the UK for three years when he developed symptoms of depression. The key stressors for him were job pressures, immigration and the need to adjust with loss of family support. He had not had enough time to integrate after migration. When he discussed his situation with his parents in Bangladesh, they called him home for a break. He, therefore, did not present to services, but on the other hand, family support mitigated against social stressors in the UK.

Discussion: This example demonstrates how the level of acculturation can influence behaviour in relation to access to services. In a different scenario, Anish may have been a second generation immigrant and could have been well integrated (identifies with both cultures) or assimilated (identify with the new culture) with the UK culture and used systems of care early.

- b. *Beliefs and attributions to illness*: culture influences beliefs around health and attributions to illness. In many cultures, the mind and body are viewed as one. Therefore depression and other mental illnesses can present as somatic symptoms. As an example, the distress and symptoms of depression can be attributed to neurasthenia (meaning weakening of nerves) by some in the Chinese culture (Kleinman, 1982). McCabe and Priebe (2004) discussed that biological explanations were much more frequently cited by Caucasians than African Caribbean and West Africans when describing symptoms of mental illness. Our research found key attributions around ‘previous wrongdoing’, supernatural beliefs, and social and biological factors in the sample of participants from South Asian Muslim, Black African and African Caribbean populations. Other attributions like the use of drugs and being arrested have also been quoted (Rathod *et al.*, 2010). Typical religious attributions can include ‘will of God or *obeah* (casting of an evil eye)’ by another.

Stigma plays an important role in cultural attributions to illness. Shame is often associated with stigma. Women can be more affected than men, especially within the South Asian Muslim communities. This gender difference can be attributed to gender bias including role expectations, rights and responsibilities.

In relation to cognitive biases, having a good cultural understanding of the individual’s background may be helpful when considering cultural adaptations in relation to their assumptions, beliefs, norms and illness attributions. An understanding of the attributions helps clinicians in engaging people and addressing their distress.

- c. *Help-seeking behaviours defining pathways into care*: an individual’s attributions to health, illness and beliefs about health and illness often determine help-seeking pathways into care. Mistrust or trust for local services also dictates the speed of access by different communities. Some individuals may choose to seek help from their cultural or traditional

healers first or in addition to conventional services. Understandably, delays in presenting to appropriate services may occur and prolong the duration of untreated symptoms.

Case example

Sadiq is a 21-year-old gentleman who developed early symptoms of psychosis. He was irritable with his family and spent a lot of time in his room. His family noticed changes in his behaviour and decided to send him to Pakistan and get him married. They thought that if he was given responsibility, he might 'pull himself together'. On return to the UK with his wife, his mental state deteriorated further. He developed frank hallucinations and remained distracted and fearful. His parents took him to the local Mosque, and the Imam gave him a Taweez (arm locket). Six months later, he had been very unwell and finally his GP referred him to the local crisis team.

One of the key predictors of prognosis in psychosis is duration of untreated psychosis. In recognition of this key prognostic factor, the access and waiting time standard has been introduced in the UK (NHS England, 2015) and integrated care pathways have been developed that ensure early access to services and evidence-based interventions for first episode psychosis patients (Rathod *et al.*, 2016). In Sadiq's case, the duration of untreated psychosis was 6 months before he presented to services. This example demonstrates how

Table 2. Cultural barriers to accessing therapy (from Rathod *et al.*, 2015); reproduced with permission from John Wiley and Sons)

Cultural barriers to therapy

Mistrust of services/practitioners
 Worries about confidentiality/breach
 Poor information on psychological therapies/accessibility
 Language and terminology leading to being misunderstood
 Fear of being stigmatized
 Previous experience
 Stereotyping by therapists
 Doubt regarding CBT being empowering enough
 Lack of understanding of cultural norms, values by the therapist
 Perception of cultural incompetence
 Clinician's beliefs in the power of drugs
 Faith/spirituality and religion
 Individualism vs collectivism
 Gender issues
 Racism/colonial history
 Interpretation problems
 Financial burden
 Practical matters, e.g. environment of therapy, transport issues

help-seeking pathways into care can influence access to services and awareness of how culture influences these pathways can help clinicians in understanding differing orientation to care and health.

- d. *Cultural orientation towards psychotherapy*: cultural beliefs regarding interventions can define whether people from certain cultural groups will engage with therapies like CBT.

Practical considerations

There is a complex relationship between religion, politics and health services in any country. In some LMICs, social factors, like poverty, urbanization, internal migration and lifestyle changes, are moderators of the burden of mental illness and service provision (Rathod *et al.*, 2017). It helps if clinicians and therapists understand to what extent policies promote integration with the other sectors such as justice, social care and development of services to ensure a more comprehensive (prevention, promotion and treatment) and holistic approach to the delivery of mental health services and experiences of individuals in the system.

Racism and effects

Racial discrimination is universal across cultures, even within diverse ethnic groups. This can apply to discrimination internally within LMIC or for minority groups in HMIC. When one is feeling vulnerable, it is difficult to talk about these experiences. Similarly, clinicians may avoid discussions about racism or discrimination. As a result, individuals can feel dissatisfied as they have not been able to discuss their main concerns in therapy. Sometimes therapists report that discussions about racism can be anxiety provoking; some admit to avoiding it due to fear of being politically incorrect or because they might say something that could easily be misinterpreted. Some therapists report that they do not feel well trained to address issues pertaining to racism or discrimination in therapy.

On the other hand, some individuals prefer talking about these problems to someone of the same colour or background as they feel they would be understood better. Moreover, they argue that a therapist from their background and colour would relate better to their experiences. A culturally competent therapist will need to recognize the potential significance of the experience of everyday racism and its impact on an individual's mental health. One has to acknowledge the disadvantages to meeting a therapist from the same background as well. This can include fear of loss of confidentiality in the community or fear of being judged.

Technical adjustments

This involves adjustment of methods, including the mode and manner of therapy and various clinical issues within the therapy for individuals of different backgrounds. Technical adjustment involves an understanding of:

- a. *Setting and environment of therapy*: therapists use clinic setting or individual's homes and in some circumstances meet at an external venue. Meeting people in their homes gives the opportunity to see their home environments, neighbourhoods, social supports and meet family members. In either situation, the environment should be culturally welcoming;

information should be available in appropriate languages and if needed interpreters should be used (Rathod and Naeem, 2012).

In many LMIC distance and transport can be an issue. Individual adjustments including length and number of sessions need to be made. Technology can be an enabler, although connectivity can be a problem.

- b. *Therapeutic relationship*: the therapeutic relationship or alliance is accepted to be one of the critical factors, if not the crucial factor, in determining the outcome of psychotherapy (Warwar and Greenberg, 2000). Therapists need to strive to build the therapist alliance at every stage of therapy including:
1. Engagement and assessment;
 2. Individualized collaborative case formulation;
 3. Effective sequential interventions based on the case formulation;
 4. Relapse prevention and termination.

First impressions are important (Anez et al., 2005). The personal style of the therapist can play a role in the development of therapeutic alliance. Some suggest (Taylor et al., 2006) a more personal and friendly style – which conveys warmth and trust – in the first session as helpful in fostering a therapeutic alliance.

Limited and measured personal disclosure in a safe manner is often a way of gaining trust. One may ask ‘how would you know how I feel?’. Therapists can answer using personal experience: ‘I do not as your experience is unique to you. However, I do know that we all have times of distress and how we feel during these times’.

Case example

Simi presented to a therapist in a very distressed state. She had lost her business and felt that she could never go on after this loss. Engaging her in treatment was very difficult as she felt extremely hopeless. After struggling for many sessions, the therapist chose to share a personal experience of surviving cancer with Simi and how they recovered and built their resilience after the illness. Simi engaged after this session and started rationalizing – ‘If you can survive this, I can also survive the loss of my business’.

Often limited self-disclosure is about building trust and respect more than the disclosing personal information.

- c. *Therapeutic style*: Western therapists use the rational, collaborative and cognitive approaches to understanding the nature and origin of problems and coping strategies to deal with them. Eastern therapies stress the importance of experience and use meditation and mindfulness techniques. Therefore, there may be a mismatch of cultural values in therapy. However, an experienced therapist can use useful techniques that suit the individual by suitable adaptations.
- d. *Family structures and goals*: in many cultures, families function as a unit and family members are closely involved in the therapeutic process. Families can be used as a strength as they can help engage individuals, help with homework and also provide the

social support that aids recovery. The adverse impact of complex relationships also needs acknowledgment.

- e. *The role of religion*: spiritual development is a vital part of all cultures (Laungani, 2004). Sometimes ignorance or interpretation of spiritual experiences as manifestations of psychopathology and lack of confidence can occur, which results in the key issue not being addressed. Clinicians in one study reported feeling disempowered when faced with the subject of religion and spirituality. The influence of religion and spirituality remains strong in Eastern cultures despite Westernization and acculturation (Williams *et al.*, 2006). Some services flourish through close collaboration with religious institutions.

Theoretical modifications

Some changes in theoretical concepts need to be made for the best fit for an individual and their cultural strength. This can involve the use of different ways of explaining concepts. Some examples are cited below:

- (a) *Body and mind*: in many cultures, the body and mind are considered one entity. The discussions used in therapy around the relationship between physical symptoms and mental illness reflect the holistic view of health and appreciation of this concept varies between cultures. Understanding what the individual values can be helpful in deciding explanations used in therapy.
- (b) *Self and ego boundaries*: in some individuals from Eastern cultures, boundaries between self, family and society are narrow and influence interactions. For instance, the interdependent relationships with family and communities often determine how an individual seeks help.
- (c) *Individuality and collectiveness*: when a therapist is insensitive to cultural norms and cultural explanations of idioms of distress, there could be obstacles in the cognitive and behaviour change processes, especially if the explanations used for change do not line with cultural values. In collectivist cultures, there is an interdependence with group members and the self is defined by group identity. Individuals prioritize communal needs over their own. In many Western cultures, the self is autonomous and independent of groups. Priority is given to personal goals, and the need often involves exchange relationships, i.e. ‘what’s in this relationship for me?’.

Collaborative empiricism underpinning psychological interventions can be a challenge when working with people from LMIC. Some literature suggests that individuals take on a passive role as they probably would with their traditional healer or ‘guru’. However, if a therapist takes an overtly directive style they may be viewed by individuals as controlling (an agent of a dominant culture). Some individuals may respond better to a structured and prescriptive approach to therapy (Iwamasa *et al.*, 2006). Similarly, in some, exploration of opinions about the problem and collaboration may lead to doubts about the therapist’s competencies, particularly when the locus of control is believed to be external and the individual has a desire for a quick fix. A paternalistic attitude can sometimes be more therapeutic.

Conclusion

This paper highlights the process, methods, foci and framework of cultural adaptation of CBT. As far as we are aware this is the first framework for adapting CBT for individuals from non-Western cultures that is evidence based. Adaptation of CBT can be accomplished through a series of steps that involve co-production with the key stakeholders. The process of adaptation should focus on both theoretical and philosophical considerations as well as on practical issues in improving access to therapy and in adjusting therapy techniques for the need of individuals and local population.

Main points

- (1) Culturally adapted CBT shows better results in individuals from different cultural groups.
- (2) Adaptation of therapy requires a robust methodology.
- (3) Every individual has a unique culture that is influenced by their wider culture, sub-culture and further developed through unique life experiences.
- (4) There must be flexibility in applying the culturally adapted therapy, and clinicians should be aware of their biases, and propensity to stereotype when working with people from minority groups.

Conflicts of interest

None of the authors has received funding to prepare this manuscript. The authors are experts in this field and have received research grants, lectured on the topic and published widely in this area.

Ethical statement

All authors have abided by the Ethical Principles of Psychologists and Code of Conduct as set out by the APA. Ethical approval was not needed for this paper as it does not present a trial.

Suggestion for follow-up reading

Rathod S, Kingdon D, Pinninti N, Turkington D, Phiri P (2015). *Cultural Adaptation of CBT for Serious Mental Illness: A Guide for Training and Practice*. Wiley-Blackwell.

References

- Anez LM, Paris M Jr, Bedregal LE, Davidson L, Grilo CM** (2005). Application of cultural constructs in the care of first generation Latino clients in a community mental health setting. *Journal of Psychiatric Practice* **11**, 221–230. doi: [00131746-200507000-00002](https://doi.org/10.1017/S1754470X18000247)
- Beck A** (1964). Thinking and depression. II. Theory and therapy. *Archives of General Psychiatry* **10**, 561–571.
- Bernal G, Bonilla J, Bellido C** (1995). Ecological validity and cultural sensitivity for outcome research: issues for the cultural adaptation and development of psychosocial treatments with Hispanics. *Journal of Abnormal Child Psychology* **23**, 67–82.

- Berry J** (2005). Acculturation: living successfully in two cultures. *International Journal of Intercultural Relations* **29**, 697–712.
- Barrera M, Castro FG** (2006). A heuristic framework for the cultural adaptation of interventions. *Clinical Psychology: Science and Practice* **13**, 311–316.
- Carter MM, Sbrocco T, Gore KL, Marin NW, Lewis EL** (2003). Cognitive-behavioral group therapy versus a wait-list control in the treatment of African American women with panic disorder. *Cognitive Therapy and Research* **27**, 505–518.
- Clark DA, Beck AT, Alford BA** (1999). *Scientific Foundations of Cognitive Theory and Therapy of Depression*. New York, NY: John Wiley and Sons, Inc.
- Domenech Rodríguez MM, Weiling E** (2004). Developing culturally appropriate, evidence-based treatments for interventions with ethnic minority populations. In M Rastogin and E Weiling (eds), *Voices of Color: First Person Accounts of Ethnic Minority Therapists* (pp. 313–333). Thousand Oaks, CA: Sage.
- El Rhermoul F-Z, Naeem F, Kingdon D, Hansen L, Toufiq J** (2018) A qualitative study to explore views of patients, carers and mental health professionals' views on depression in Moroccan women. *International Journal of Culture and Mental Health* **11**, 178–193. doi: [10.1080/17542863.2017.1355397](https://doi.org/10.1080/17542863.2017.1355397)
- Garcia G, Zea C** (eds) (1997). *Psychological Interventions and Research with Latino Populations*. Boston, MA: Allyn and Bacon.
- Hays PA, Iwamasa GY** (2006). *Culturally Responsive Cognitive-Behavioral Therapy*. Washington, DC: American Psychological Association.
- Hinton DE, Pham T, Tran M, Safren SA, Otto MW, Pollack MH** (2004). CBT for Vietnamese refugees with treatment-resistant PTSD and panic attacks: a pilot study. *Journal of Traumatic Stress* **17**, 429–433.
- Hwang W, Wood JJ, Lin K, Cheung F** (2006). Cognitive behavioural therapy with Chinese Americans. Research, theory and clinical practice. *Cognitive and Behavioural Practice* **13**, 293–303.
- Iwamasa GY, Hsia C, Hinton D** (2006). Cognitive-behavioral therapy with Asian Americans. In PA Hays and GY Iwamasa (eds), *Culturally Responsive Cognitive Behavioral Therapy: Assessment, Practice, and Supervision* (pp. 117–140). Washington, DC: American Psychological Association.
- Kleinman A** (1982). Neurasthenia and depression: a study of somatization and culture in China. *Culture, Medicine and Psychiatry* **6**, 117–190.
- Kubany ES, Hill EE, Owens JA** (2003). Cognitive trauma therapy for battered women with PTSD: preliminary findings. *Journal of Traumatic Stress* **16**, 81–91.
- Lau AS** (2006). Making the case for selective and directed cultural adaptations of evidence-based treatments: examples from parent training. *Clinical Psychology: Science and Practice* **13**, 295–310.
- Laungani P** (2004). *Asian Perspectives in Counselling and Psychotherapy*. Taylor and Francis. ISBN 0203697081, 9780203697085
- Leong T, Lee H** (2006). A cultural accommodation model for cross cultural psychotherapy: illustrated with the case of Asian American. *Psychotherapy* **43**, 410–423.
- Li W, Zhang Li, Luo X, Liu B, Liu Z, Lin F, Liu Z, Xie Y, Hudson M, Rathod S, Kingdon D, Husain N, Liu X, Ayub M, Naeem F** (2017). A qualitative study to explore views of patients', carers' and mental health professionals' to inform cultural adaptation of CBT for psychosis (CBTp) in China. *BMC Psychiatry* **17**, 131. doi: [10.1186/s12888-017-1290-6](https://doi.org/10.1186/s12888-017-1290-6)
- McCabe R, Priebe S** (2004). Explanatory models of illness in schizophrenia: Comparison of four ethnic groups. *British Journal of Psychiatry* **185**, 25–30.
- Naeem F, Gul M, Irfan M, Munshi T, Asif A, Rashid S, Khan MN, Ghani S, Malik A, Aslam M, Farooq S, Husain N, Ayub M** (2015a). Brief culturally adapted CBT (CaCBT) for depression: a randomized controlled trial from Pakistan. *Journal of Affective Disorders* **177**, 101–107.

- Naeem F, Phiri P, Nasar A, Gerada A, Munshi T, Ayub M, Rathod S** (2016). An evidence-based framework for cultural adaptation of cognitive behaviour therapy: process, methodology and foci of adaptation. *World Cultural Psychiatry Research Review* **11**, 61–70.
- Naeem F, Saeed S, Irfan M, Kiran T, Mehmood N, Gul M, Munshi T, Ahmad M, Kazmi A, Husain N, Farooq S, Ayub M, Kingdon D** (2015b). Brief culturally adapted CBT for psychosis (CaCBTp): a randomized controlled trial from a low income country. *Schizophrenia Research* **164**, 143–148.
- Naeem F, Waheed W, Gobbi M, Ayub M, Kingdon D** (2010). Preliminary evaluation of culturally sensitive CBT for depression in Pakistan: findings from developing culturally sensitive CBT Project (DCCP). *Behavioural and Cognitive Psychotherapy* **39**, 165–173.
- Nagel T, Robinson G, Condon J, Trauer T** (2009). Approach to treatment of mental illness and substance dependence in remote indigenous communities: results of a mixed methods study. *Australian Journal of Rural Health* **17**, 174–182.
- NHS England** (2015). Guidance to support the introduction of access and waiting time standards for mental health services.
- Okasha A, Okasha T** (2000). Notes on mental disorders in pharaonic Egypt. *History of Psychiatry* **11**, 413–424.
- Patel V, Araya R, Chatterjee S, Chisholm D, Cohen A, De Silva M et al.** (2007). Treatment and prevention of mental disorders in low-income and middle-income countries. *The Lancet* **370**, 991–1005.
- Rahman A, Malik A, Sikander S, Roberts C, Creed F** (2008). Cognitive behaviour therapy based intervention by community health workers for mothers with depression and their infants in rural Pakistan: a cluster-randomised controlled trial. *The Lancet* **372**, 902–909.
- Rathod S, Kingdon D, Smith P, Turkington D** (2005). Insight into schizophrenia: the effects of cognitive behavioral therapy on the components of insight and association with sociodemographics – data on a previously published randomised controlled trial. *Schizophrenia Research* **74**, 211–219.
- Rathod S, Kingdon D, Pinninti N, Turkington D, Phiri P** (2015). *Cultural Adaptation of CBT for Serious Mental Illness: A Guide for Training and Practice*. Wiley-Blackwell.
- Rathod S, Garner C, Griffiths A, Dimitrov B, Newman-Taylor K, Woodfine C, Hansen L, Tabraham P, Ward K, Asher C, Phiri P, Naeem F, North P, Munshi T, Kingdon D** (2016). Protocol for a multicentre study to assess feasibility, acceptability, effectiveness and direct costs of TRIumPH (Treatment and Recovery In Psychosis): Integrated Care Pathway for Psychosis. *British Medical Journal Open* **6**, e012751. doi: [10.1136/bmjopen-2016-012751](https://doi.org/10.1136/bmjopen-2016-012751)
- Rathod S, Giga L, Degnan A et al.** (2017a) The current status of culturally adapted mental health interventions: a practice-focused review of meta-analysis. *Neuropsychiatric Disease and Treatment* **13**, 1–14
- Rathod S, Naeem F** (2012). Can you do meaningful cognitive-behavioural therapy with an interpreter? In Bhui K, *Elements of Culture and Mental Health: Critical Questions for Clinicians*. Royal College of Psychiatrists.
- Rathod S, Phiri P, Harris S, Underwood C, Thagadur M, Padmanabi U, Kingdon D** (2013). Cognitive behaviour therapy for psychosis can be adapted for minority ethnic groups: a randomised controlled trial. *Schizophrenia Research* **143**, 319–326.
- Rathod S, Phiri P, Kingdon D, Gobbi M** (2010). Developing culturally sensitive cognitive behaviour therapy for psychosis for ethnic minority patients by exploration and incorporation of service users' and health professionals' views and opinions. *Behavioural and Cognitive Psychotherapy* **38**, 511–533.
- Rathod S, Pinninti N, Irfan M, Gorczynski P, Rathod P, Gega L, Naeem F** (2017b). Mental health service provision in low and middle income countries. *Health Services Insights* **10**. doi: [10.1177/1178632917694350](https://doi.org/10.1177/1178632917694350)
- Resnicow K, Soler R, Braithwait L, Ahluwalia S, Butler J** (2000). Cultural sensitivity in substance abuse prevention. *Journal of Community Psychology* **28**, 271–290.

- Rojas G, Fritsch R, Solis J, Jadresic E, Castillo C, González M et al.** (2007). Treatment of postnatal depression in low-income mothers in primary-care clinics in Santiago, Chile: a randomised controlled trial. *The Lancet* **370**, 1629–1637.
- Scorzelli J, Scorzelli MR** (1994). Cultural sensitivity and cognitive therapy in India. *The Counselling Psychologist* **22**, 603–610.
- Schwarzbaum SE** (2004). Low-income Latinos and dropout: strategies to prevent dropout. *Journal of Multicultural Counselling and Development* **32**, 296–306.
- Taylor BA, Beatriz-Gambourg M, Rivera M, Laureano D** (2006). Constructing cultural competence perspectives of family therapists working with Latino families. *American Journal of Family Therapy* **34**, 429–445.
- Tseng WS** (1999). Culture and psychotherapy: review and practical guidelines. *Transcultural Psychiatry* **36**, 131–179.
- Tseng W-S, Chang SC, Nishizono M** (2005). *Asian Culture and Psychotherapy: Implications for East and West*. Honolulu, HI: University of Hawaii Press.
- Warwar S, Greenberg L** (2000). *Handbook of Counseling Psychology* (3rd edn). New York, NY: John Wiley and Sons, Inc.
- Williams W, Hean K, Beverly H** (2006). Cultural considerations in using cognitive behaviour therapy with Chinese people: a case study of an elderly Chinese woman with generalized anxiety disorder. *New Zealand Journal of Psychology* **35**, 153–162.

Learning objectives

- (1) To understand why the cultural adaptation of cognitive behaviour therapy (CBT) is important.
- (2) To understand the process of development of an evidence-based framework of adaptation.
- (3) To learn adaptation techniques relevant to practising personalized CBT with individuals from different cultures.