



Weeds

VOLUME 8
JULY 1960
NUMBER 3

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

C. J. WILLARD, *Editor*, Department of Agronomy, H. & F. Bldg.,
1827 Neil Ave., The Ohio State University, Columbus 10, Ohio.

W. C. JACOB, *Business Manager*, Department of Agronomy, Univer-
sity of Illinois, Urbana, Illinois.

EDITORIAL COMMITTEE

O. C. LEE, Department of Botany and Plant Pathology, Purdue
University, Lafayette, Indiana.

R. A. PETERS, Department of Plant Science, University of Connecti-
cut, Storrs, Connecticut.

W. C. ROBOCKER, Crops Research Division, ARS, USDA, State
College of Washington, Pullman, Washington.

E. G. RODGERS, Department of Agronomy, University of Florida,
Gainesville, Florida.

WEEDS is a quarterly journal published by the Weed Society of America. Editorial office is located at Department of Agronomy, H. and F. Bldg., The Ohio State University, 1827 Neil Ave., Columbus 10, Ohio. Printing is by the W. F. Humphrey Press Inc., Geneva, New York. Subscription price is \$6.00 yearly for four issues; single copies \$1.50. Address all communications regarding subscriptions, advertising, and reprints to W. C. Jacob, Department of Agronomy, University of Illinois, Urbana, Illinois. Inquiries concerning information on manuscripts and other material for publication should be addressed to the Editorial Office. All checks, money orders, and other remittances should be made payable to the Weed Society of America.

Entered as second-class matter at the post office at Urbana,
Illinois with additional entry at Geneva, New York.

Table of Contents


	<i>Page</i>
Effect of 2,4-Dichlorophenoxyacetic Acid on the Nucleic Acid and Protein Content of Seedling Tissue. S. H. West, J. B. Hanson and J. L. Key.	333
The Degradation of Isolated Microsomes from Cucumber and Corn Seedlings. S. H. West and J. B. Hanson	341
Control of Weeds in Legume Seedings with 4-(2,4-DB), Dalapon, and TCA. Elroy J. Peters and Frank S. Davis	349
Chemical Weed Control in Young Grapes. C. C. Doll	368
Herbicidal Value of Arsenic Trioxide in Eastern United States. Albert L. Leaf and Robert E. Smith, Jr.	374
Comments on Tests of Herbicide Mixtures. Donald P. Gowing	379
Effect of Histidine on the Inhibitory Action of 3-Amino-1,2,4-triazole. J. L. Hilton	392
Field Bindweed Control by Combinations of Cropping, Cultivation, and 2,4-D. O. G. Russ and L. E. Anderson	397
Characteristics of Johnsongrass Rhizomes. L. E. Anderson, A. P. Appleby, and J. W. Weseloh	402
Chemistry of Some Amino Acid Derivatives of 2,2-Dichloropropionic and 2,3,6-Trichlorobenzoic Acids. Charles F. Krewson and Edward J. Saggese	407
Effect of Sucrose on the Toxicity of Several Phenylurea Herbicides to Barley. W. A. Gentner and J. L. Hilton	413
Field Bindweed (<i>Convolvulus arvensis</i>) Control with Soil Applications of Phenoxy Herbicides. A. F. Wiese and H. E. Rea	418
The Control of Weeds Around Young Apple Trees. J. S. Leefe and R. P. Longley	422
Mesquite Root Inhibition Tests to Study Inhibitory Activity, Absorption and Translocation of 2,4-D and 2,4,5-T. Richard Behrens and Howard L. Morton	427
The Fate of Long Chain Esters of 2,4-D in Plants. D. James Morr� and B. J. Rogers	436
The Effect of Certain Triazines on C ¹⁴ O ₂ Fixation in Red Kidney Beans. Floyd M. Ashton, Gunter Zweig, and George W. Mason	448
Biochemical Analyses, Germination, and Production of Black and Brown Seed of <i>Halogeton glomeratus</i> . M. Coburn Williams	452
Brief Papers	
Iso-octyl Ester of 2,4,5-T in Hardwood Control. Carter B. Gibbs	462
An Experimental Plot Sprayer. K. F. Best and A. Wenhardt	464
Notes on the Distribution of the Russian Thistle (<i>Salsola kali</i> L. var. <i>tenuifolia</i>) in Mexico. Paulino Rojas-M. and M. Rojas-Garciduenas	466
Minutes of the Business Meeting and Committee Reports, Weed Society of America Meeting, February 22 to 25, 1960	469
Constitution of the Weed Society of America	482
Report of the Terminology Committee, Weed Society of America	487
News and Notes	521
Sustaining Members	522
Bibliography of Weed Investigations, October to December, 1959	523

Advertisers Index

U. S. Borax & Chemical Corp.	ii
Stauffer Chemical Co.	iii
Chipman Chemical Co.	iv
Spraying Systems Co.	v
duPont de Nemours & Co.	vi



**Four
easy
ways to
Destroy
Weeds**



Nonselective Herbicides

for Dependable Action

1. UREABOR®

A nonselective, granular complex of sodium borate and substituted urea. Low application rates are a feature. Apply with the special new PCB Spreader for best results.

2. BENZABOR®

A blend of trichlorobenzoic acid and sodium borates. Kills deep-rooted, noxious weeds. Low application rates give maximum control with the utmost economy; use the convenient PCB Spreader.

3. POLYBOR-CHLORATE®

Highly soluble; for spray or dry application. It gives a quick knock-down; destroys top growth and roots. A general nonselective herbicide.

4. Concentrated BORASCU®

A nonselective, granular material. Apply by hand or with a mechanical spreader. Long residual action.

UNITED STATES BORAX



& CHEMICAL CORPORATION

630 Shatto Place, Los Angeles 5, California

= NEWS ABOUT

VAPAM[®] =

Summary of Application Methods

After careful soil preparation, a wide range of application methods can be used for effective control of weeds, germinating weed seed, nematodes, and soil fungi with VAPAM fumigant. Most of the methods do not require special equipment.

For Seed Beds, Plant Beds, Lawns and Other Limited Areas—A sprinkling can, hose proportioner or rotary tiller are all effective. In some cases, it might be practical to use soil injection.

Field Application to Beds or Rows—Soil injection, bedding-over or soil covering are the quickest and easiest methods for most growers. Basin flooding is the recommended method for treating tree replant sites.

For Treating Entire Fields—Soil injection, sprinkler irrigation, and check or flood irrigation are recommended.

Full details on these application methods, plus the treatment of potting soils, are on the VAPAM label.

Copy of Label Available

Write to Stauffer for a copy of the label, or for further information on VAPAM. Address your inquiry to Stauffer Chemical Company, Agricultural Chemicals Division, 380 Madison Ave., New York 17, N. Y., or 636 California St., San Francisco 8, Calif.

©VAPAM is Stauffer Chemical Company's trade-mark (registered in principal countries) for sodium methyl dithiocarbamate, a soil fumigant.





ATLACIDE: Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water.

ATLACIDE-2,4-D: Offers dual killing action of sodium chlorate and 2,4-D.

CHLOREA: A combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray.

CHLOREA GRANULAR Similar to Chlorea, but a "pelletized" material. No mixing...easy to apply by hand or with mechanical spreader.

CHLORAX "40": A composition of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys vegetation and algae in ponds and lakes. Controls certain turf weeds. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

SODIUM ARSENITE: A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas "A". Applied dry or as a spray.

CHIP-CAL: New low-lime calcium arsenate formulation in granular and spray powder forms. Used as crabgrass preventatives for established turf. Applied any time before crabgrass seed starts to germinate.

2,4-D & 2,4,5-T: 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

METHOXONE: Contains MCP sodium salt. Used for control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

BUTOXONE: A liquid containing 4-(2,4-DB). For selective control of many broadleaf weeds in established alfalfa and certain other legumes.

Write for Weed Control Booklets

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

Bound Brook, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.


Bessemer, Ala.

Manufacturers of Weed Killers Since 1912




the precision nozzle for effective spraying


Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.



TEEJET SPRAY NOZZLE male pipe connection



TEEJET SPRAY NOZZLE female pipe connection



INTER-CHANGE-ABLE ORIFICE TIPS flat and cone spray types

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

SPRAYING SYSTEMS CO.
Engineers and Manufacturers

3296 RANDOLPH STREET

BELLWOOD, ILLINOIS

For Agriculture and Industry . . .

Du Pont WEED KILLERS

*offer new economies
and efficiency in killing
weeds, grass and brush*

KARMEX® diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KARMEX® DL for pre-emergence weed control in cotton.

TELVAR® monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KLOBEN® neburon for weed and grass control in nursery plantings and tomatoes.

DYBAR® fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

TRYSBEN® 200, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broadleaf weeds; and certain woody vines and brush.

AMMATE® X for long-term, low-cost brush control.



Better Things for Better Living . . . *through Chemistry*