

visits among seniors are frequently instigated by a fall at home. Some of these patients develop intracranial hemorrhage (ICH) because of falling. There has been little research on the frequency of ICH in elderly patients who fall, and on which clinical factors are associated with ICH in these patients. The aim of this study was to identify the incidence of ICH, and the clinical features which are associated with ICH, in seniors who present to the ED having fallen. **Methods:** This was a prospective cohort study conducted in three EDs. Patients were included if they were age >65 years, and presented to the ED within 48 hours of a fall on level ground, off a bed/chair/toilet or down one step. Patients were excluded if they fell from a height, were knocked over by a vehicle or were assaulted. ED physicians recorded predefined clinical findings (yes/no) before any head imaging was done. Head imaging was done at the ED physician's discretion. All patients were followed for 6 weeks (both by telephone call and chart review at 6 weeks) for evidence of ICH. Associations between baseline clinical findings and the presence of ICH were assessed with multivariable logistic regression. **Results:** In total, 1753 patients were enrolled. The prevalence of ICH was 5.0% (88 patients), of whom 74 patients had ICH on the ED CT scan and 14 had ICH diagnosed during follow-up. 61% were female and the median age was 82 (interquartile range 75-88). History included hypertension in 76%, diabetes in 29%, dementia in 27%, stroke/TIA in 19%, major bleeding in 11% and chronic kidney disease in 11%. 35% were on antiplatelet therapy and 25% were on an anticoagulant. Only 4 clinical variables were independently associated with ICH: bruise/laceration on the head (odds ratio (OR): 4.3; 95% CI 2.7-7.0), new abnormalities on neurological examination (OR: 4.4; 2.4-8.1), chronic kidney disease (OR: 2.4; 1.3-4.6) and reduced GCS from baseline (OR: 1.9; 1.0-3.4). Neither anticoagulation (OR: 0.9; 0.5-1.6) nor antiplatelet use (OR: 1.1; 0.6-1.8) appeared to be associated with ICH. **Conclusion:** This prospective study found a prevalence of ICH of 5.0% in seniors after a fall, and that bruising on the head, abnormal neurological examination, abnormal GCS and chronic kidney disease were predictive of ICH. **Keywords:** intracranial hemorrhage, predictors, seniors

#### PL04

##### Comparison of the cost and the quality of the care provided to low acuity patients in an emergency department and a walk-in clinic

S. Berthelot, MD, MSc, M. Mallet, MA, D. Simonyan, MSc, J. Guertin, PhD, L. Moore, PhD, C. Boilard, BSc, J. Boulet, BSc, C. Fortier, BSc, P. Olivier, BSc, B. Huard, BSc, K. Vachon, BSc, A. Lesage, BSc, É. Lévesque, BSc, A. Mokhtari, BSc, L. Baril, MD, O. Yip, BSc, M. Bouchard, BSc, M. Létourneau, BA, A. Pineault, BA, M. Lafrenière, MD, S. Blais, MBA, Université Laval, Québec, QC

**Introduction:** Low acuity patients have been controversially tagged as a source of emergency department (ED) misuse. Authorities for many Canadian health regions have set up policies so these patients preferably present to walk-in clinics (WIC). We compared the cost and quality of the care given to low acuity patients in an academic ED and a WIC of Québec City during fiscal year 2015-16. **Methods:** We conducted an ambidirectional (prospective and retrospective) cohort study using a time-driven activity-based costing method. This method uses duration of care processes (e.g., triage) to allocate to patient care all direct costs (e.g., personnel, consumables), overheads (e.g., building maintenance) and physician charges. We included consecutive adult patients, ambulatory at all time and discharged from the ED or WIC

with a diagnosis of upper respiratory tract infection (URTI), urinary tract infection (UTI) or low back pain. Mean cost [95%CI] per patient per condition was compared between settings after risk-adjustment for age, sex, vital signs, number of regular medications and co-morbidities using generalized log-gamma regression models. Proportions [95% CI] of antibiotic prescription and chest X-Ray use in URTI, compliance with provincial guidelines on use of antibiotics in UTI, and column X-Ray use in low back pain were compared between settings using a Pearson Chi-Square test. **Results:** A total of 409 patients were included. ED and WIC groups were similar in terms of age, sex and vital signs on presentation, but ED patients had a greater burden of comorbidities. Adjusted mean cost (2016 CAN\$) of care was significantly higher in the ED than in the WIC ( $p < 0.0001$ ) for URTI (78.42[64.85-94.82] vs. 59.43[50.43-70.06]), UTI (78.88 [69.53-89.48] vs. 53.29[43.68-65.03]), and low back pain (87.97 [68.30-113.32] vs. 61.71[47.90-79.51]). For URTI, antibiotics were more frequently prescribed in the WIC (44.1%[34.3-54.3] vs. 5.8% [1.2-16.0];  $p < 0.0001$ ) and chest X-Rays, more frequently used in the ED (26.9%[15.6-41.0] vs. 13.7%[7.7-22.0];  $p = 0.05$ ). No significant differences were observed in the compliance with guidelines on use of antibiotics in UTI and in the use of column X-Ray in low back pain. **Conclusion:** Total cost of care for low acuity patients is lower in walk-in clinics than in EDs. However, our results suggest that quality-of-care issues should be considered in determining the best alternate setting for treating ambulatory emergency patients.

**Keywords:** healthcare costs, low-acuity patients, quality of healthcare

## Oral Presentations

#### LO01

##### Development and validation of an adjustment score for ruling out MI using a single high-sensitivity cardiac troponin T assay in patients with chest pain and kidney dysfunction

A. McRae, MD, PhD, S. Vatanpour, PhD, J. Ma, MSc, E. Lang, MD, J. Andruchow, MD, MSc, G. Innes, MD, MSc, M. James, MD, PhD, A. Worster, MD, MSc, P. Kavsak, PhD, University of Calgary, Calgary, AB

**Introduction:** Very low concentrations of high-sensitivity cardiac troponin can rule-out myocardial infarction (MI) at ED arrival in patients with chest pain. However, this single troponin rule-out strategy works poorly in patients with renal impairment and elevated baseline troponin levels. The objective of this study was to develop and validate a troponin adjustment strategy to accurately rule-out MI with a single hs-cTnT measurement in patients with kidney dysfunction. **Methods:** We used data from three cohorts of ED chest pain patients to develop an adjustment score for a high-sensitivity troponin T (hs-cTnT) assay in patients with kidney dysfunction. The derivation cohort ( $n = 8846$ ) used administrative and registry data. Two validation cohorts ( $n = 1187$  and  $1092$ ) were prospectively-collected. The score assigned points for increasing hs-cTnT levels and subtracted points for lower estimated glomerular filtration rate (eGFR). In the derivation cohort, hs-cTnT concentrations achieving 98.5% sensitivity in of patients with eGFR  $\geq 60$ , 45-59, 30-44, 15-29 and  $< 15$  were assigned ascending positive integer values. Negative integer values were assigned to eGFR values 45-59, 30-44, 15-29 and  $< 15$ . The scores for troponin and eGFR were summed for each patient, with scores ranging from  $-4$  to  $+5$ . The proportion of patients with 7-day MI ruled out by a score  $\leq 0$ , sensitivity, NPV, negative likelihood ratio (LR-) and area under the curve (AUC) were quantified