The Journal of Laryngology and Otology

EDITED BY
G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

	A family with high-tonal objective tinnitus .	2 .	J. D. GLANVILLE,	
			R. R. A. Coles and Brenda M. Sullivan	I
	The role of perilymph in human labyrinth	ine	and Brenda M. Sullivan	
	physiology		ROBERT G. HUGHES	II
1	Grommets and glue ears: a clinical trial .		S. H. RICHARDS, D. KILBY, J. D. SHAW and H. CAMPBELL	17-
1	Clinical electronystagmography		R. E. HOOPER and D. WORGAN	23
	Methods for early detection of hearing loss in infa-	nts	M. M. ALTMAN and R. Shenhav	35
	Variations in vocal wave duration		YVAN LEBRUN and JANNIE HASQUIN	43
	Sarcoma of the ethmoid in children		J. L. W. WRIGHT, P. D. PHELPS and A. G. MACIVER	Em
	To also and its the studies of allines leaves		G. S. LUM	57 69
	In-vivo and in-vitro studies of rhinoscleroma .		G. S. LUM	09
	Clinical records—			
	Pemphigus of the larynx	- 6	P. N. PATHAK	81
	Localized amyloidosis of the larynx		P. Beasley	83
	Primary lingual tuberculoma	-	P. K. KAKAR and V. P. Sood	89
	A pedunculated tonsillar lymphangioma .		P. G. VISVANATHAN	93
	General notes			97

Headley Brothers Ltd

Ashford, Kent

Annual Subscription £8 net, U.S.A. \$23

Monthly, LI net, post free

The Journal of Laryngology and Otology

(Founded in 1887 by Morell Mackenzie and Norris Wolfenden)

G. H. BATEMAN

ASSISTANT EDITOR
LIONEL TAYLOR

- r. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.
- 2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.


The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951), *Journal of Laryngology and Otology*, 65, 33. The titles of all Journals should be given without abbreviation.

- It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.
- 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.
- 4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to Headley Brothers Ltd., The Invicta Press, Ashford, Kent. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.
- 5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.
- 6. Editorial communications may be addressed to The EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent.
- 7. The annual subscription is £8 (eight pounds sterling) post free; U.S.A. \$23 post free, and payable in advance.
- 8. Single copies of current or back numbers (when available) will be on sale at $f_{\rm I}$ (including postage) each.
- All subscriptions, advertising and business communications should be sent to the publishers, or subscription agents.

HEADLEY BROTHERS LTD.

The Invicta Press, ASHFORD, Kent.

O Journal of Laryngology and Otology Ltd., 1971



- * Permit accurate pure tone audiometry in conditions of ambient noise too high for unshielded earphones
- ★ Easily fitted to any audiometer earphone using standard TDH 39 inserts with MX-41 AR Cushions
- * Complete freedom from threshold shift—no change in audiometer calibration at time of fitting
- * Subsequent periodic recalibration can be effected on standard ISO 6 cc. coupler without removal of inserts from Audiocups
- * Invaluable where a silent cabin or sound-proof room is not available
- * Give extra attenuation when a booth is available.

Write for detailed leaflet and price list to:-

AMPLIVOX HEARING CONSERVATION LTD.

Beresford Avenue, Wembley, Middlesex. England. Tel: 01-902 8991.

Oto-, Rhino-, Laryngology

Important, up-to-date, and scientifically accurate abstracts of reports from the world's biomedical literature dealing with Oto-, Rhino-, Laryngology are brought to your desk in this monthly publication.

First published in 1948,
Oto-, Rhino-, Laryngology currently contains some 4,750 abstracts each year.

MAIN CHAPTER HEADINGS

- 1. SKULL
- 2. FACE AND LIPS
- 3. MOUTH
- 4. TONGUE
- 5. SALIVARY GLANDS
- 6. NECK
- 7. PHARYNX
- 8. ESOPHAGUS
- 9. LARYNX
- 10. TRACHEA
- 11. BRONCHI, LUNGS, MEDIASTINUM
- 12. OLFACTORY SYSTEM
- 13. EXTERNAL NOSE
- 14. NASAL CAVITY
- 15. NASAL SEPTUM

- 16. NASOPHARYNX
- 17. NASAL SINUSES
- 18. EXTERNAL EAR
- 19. MIDDLE EAR
- 20. INTERNAL EAR
- 21. OTOSCLEROSIS
- 22. MENIERE DISEASE
- 23. FACIAL NERVE
- 24. VESTIBULAR SYSTEM
- 25. CEREBELLOPONTINE ANGLE TUMORS
- 26. HEARING
- 27. SPEECH
- 28. PHONIATRICS
- 29. ANESTHESIA

Specimen copies are available on request.

Annual Subscription Rate: U.S. \$50.00/£20. 19. 0d. Sterling/Dfl. 180,00

EXCERPTA MEDICA

Herengracht 119-123, Amsterdam, The Netherlands

Floxapen simplifies antibiotic treatment in hospital...

and after

Floxapen, like Orbenin* (cloxacillin B.P.), is active against virtually all staphylococci, streptococci and pneumococci, irrespective of their sensitivity to other antibiotics.

However, it is simpler to use than Orbenin, the established first-line treatment for Gram-positive infections in hospitals, because:

The standard dose, orally or by injection, is only 250 mg. g.i.d.

Equivalent serum levels are obtained by either route.

Transfer from injectable to oral therapy may be justified at an earlier stage of treatment.

Floxapen also offers the penicillin advantages of low toxicity and bactericidal action.

For outpatient treatment, Floxapen is ideal; its simple oral dosage and minimal side-effects encourage patients to follow the correct dosage and to complete the course of therapy.

Indications: Infections due to sensitive Gram-positive organisms.

Dosage: Oral and Intramuscular:

Adults: 250 mg. q.i.d.
Children: 2-10 years-half adult dose;
Under 2 years—quarter adult dose.
Oral doses should be taken one hour before meals. In severe infections dosages may be increased

if required.

Availability: Capsules (black/caramel) and Vials, each containing

250 mg. flucloxacillin.

Contra-indications: Penicillin hypersensitivity. Subconjunctival injection.

Side-effects: Rare and mild,

as with other penicillins.

Further information is available on request.





Floxapen* (flucloxacillin) is a product of British research at Beecham Research Laboratories, Brentford, England, originators of the new penicillins



THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$24.00 per year Canada \$25.00 per year Foreign \$25.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D. FDITOR

222 PINE LAKE ROAD, COLLINSVILLE, ILLINOIS 62234



P. C. WERTH LTD. Contractors to H.M. Government and Local Authorities

Specialist Supply, Maintenance, Calibration Services for Audiological Test and Teaching Equipment

17 STRATFORD PLACE, OXFORD ST., LONDON W1N ODH

Telephone: 01-629 5559

Clinical, Diagnostic, Screening and Bekesy Audiometers. Acoustic Impedance Apparatus. Stereo Auditory Trainers. Acoustic Cabins. Artificial Mastoid, etc.

Kamplex DA11 Twin-Channel Clinical Audiometer

Our latest model with many new features:


Probably the most keenly priced twin-channel audiometer currently available.

Fowler, Sisi, Luscher and Rainville test facilities inhuilt


Complete speech facilities.

Narrow band and white noise masking to 100 db.

Pure tone attenuation to 120 db.



New for the infected ear OTOSEPTIL eardrops



Containing a broad spectrum antibiotic and possessing antifungal properties

Candida and aspergillus are found in about 20% of cases of otitis externa.

Otoseptil ear drops destroy both pathogenic bacteria and fungi commonly found in ear infections.

Otoseptil ear drops contain:
Neomycin Undecylenate – antibacterial and antifungal.
Tyrothricin – active against gram-positive cocci.

Tyrothricin — active against gram-positive cocci.

(Containing gramicidin and tyrocin).

Hydrocortisone alcohol — anti-inflammatory.

Ethylene oxide polyoxypropylene glycol condensate—
to ensure rapid tissue contact.

Otoseptil ear drops control infection, relieve pain, erythema and itching.


Neomycin Undecylenate was researched by H R Napp Ltd.

Presentation: 8 ml bottle with separate dropper.

Samples and full information available to the medical profession upon request.

H R Napp Ltd Watford wd2 7RA England

A new product by Napp - Developed by International Research Group. Dedicated to Physician and Patient for over 40 years.






Demonstrations can be arranged, full details from Degenhardt and Co Ltd Carl Zeiss House 31/36 Foley Street London W1P 8AP 01-636 8050

Cable: Zeissag London

Telex: 24300



the drop that soothes and clears infected ears

soothes by reducing inflammation with hydrocortisone clears by rapidly destroying virtually all bacteria commonly found in ear infections


OTOSPORIN drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration, Supplied in dropper bottles of 5 ml. Full information is available on request.

*Trade Mark

** OTOSPORIN



Burroughs Wellcome & Co.
(The Wellcome Foundation Ltd.)Dartford, Kent



PETERS CLINIC AUDIOMETERS AND BÉKÉSY

ARE THE IDEAL TOOLS FOR MORE ADVANCED AUDIOMETRY

The Audiometers offer:

- Two continuously variable attenuators accurate to I db.
- SISI Test (built-in) and automatic alternate loudness balance switching.
- Maximum attenuation of 120 db for loudness discomfort level tests.
- Narrow band SYNCHRONOUS masking (a new semi-automatic masking technique).
- Full speech test facilities with white noise masking.
- Plug-in circuit boards and operator output checks.
- Conversion to a true BÉKÉSY AUDIOMETER at will.
- Extra features not listed are offered by the AP/6 compared to the AP/5 AUDIOMETER.

write for full details to:

ALFRED PETERS AND SONS LTD GELL STREET SHEFFIELD 3