

SEROXAT

Now at a New Lower Price



What else could we improve but the price?

Seroxat is at a new
low price of 71p per day*.

Breacadh an lae. Re orga ag breacadh.

*MIMS Ireland, December 1994. 20 mg (30's). Full prescribing information is available on request from: Smith Kline & French, Corrig Avenue, Dun Laoghaire, Co. Dublin.
©1994 SmithKline Beecham Pharmaceuticals. 'Seroxat' is a trade mark. 9/94/SX:AD0234E.

PRESCRIBING INFORMATION

Presentation: 'Seroxat' Tablets, PA 48/50/1-2, each containing either 20 mg or 30 mg paroxetine as the hydrochloride. 20 mg : 30(OP); 30 mg :

30(OP).

Indications: Treatment of symptoms of depressive illness of all types including depression accompanied by anxiety. Prevention of relapse and also

recurrence of further depressive episodes.

Dosage: Adults: 20 mg a day. Review response within two to three weeks and if necessary increase dose by 10 mg increments to a maximum of 50 mg according to response. Give once a day in the morning with food. The tablets should not be chewed. Continue treatment for a sufficient period, which may be several months. Stop treatment gradually. Elderly: 20 mg a day increasing by increments of 10 mg up to 40 mg a day according to response. Children: Not recommended.

Severe renal impairment (creatinine clearance <30 ml/min) or severe hepatic impairment: 20 mg a day. Restrict incremental dosage if required to

achieve a plasma level of 1017/S0790966700002172 Published online by Cambridge University Press

Contra-indications: Hypersensitivity to paroxetine and related drugs; use with MAO inhibitors; unstable epilepsy or convulsive disorders; severe renal failure.

Precautions: History of mania. Cardiac conditions: caution. Caution in patients with controlled epilepsy (monitor carefully); stop treatment if

seizures develop. Caution patients about driving and operating machinery.

Drug interactions: Do not use with or within two weeks after MAO inhibitors; leave a two-week gap before starting MAO inhibitor treatment.

Possibility of interaction with tryptophan. Great caution with warfarin and other oral anticoagulants. Use lower doses if given with drug metabolising enzyme inhibitors; adjust dosage if necessary with drug metabolising enzyme inducers. Combination with other highly bound protein drugs may alter plasma levels of either. Alcohol is not advised. Care with other CNS active drugs. Keep dosage of concomitant benzodiazepines low. Use lithium with caution and monitor lithium levels. Increased adverse effects with phenytoin; similar possibility with other anticonvulsants. Pregnancy and lactation: Use in pregnancy only if essential and avoid during lactation.

Adverse reactions: Most commonly nausea, somnolence, sweating, tremor, asthenia, dry mouth, insomnia, sexual dysfunction.

Overdosage: Symptoms include nausea, vomiting, tremor, dilated pupils, dry mouth, irritability. No specific antidote. General treatment as for

overdosage with any antidepressant. Early use of activated charcoal suggested.

Product authorisation holder: SmithKline Beecham Pharmaceuticals Ltd, Corrig Avenue, Dun Laoghaire, Co. Dublin.