



## Erratum for ‘Geometric Height Inequalities and the Kodaira–Spencer Map’<sup>★</sup>

MINHYONG KIM

*Department of Mathematics, University of Arizona, Tucson, AZ 85721, U.S.A.*  
*e-mail: kim@math.arizona.edu*

(Received: 8 July 1998; in final form: 17 December 1998)

(1) Reference [2] on p. 44, line 2, should be [3]; reference [4] on p. 45, beginning of last paragraph, should be [5]; reference [5] on p. 46, line 7, should be [6]; reference [6] in the beginning of section 3 should be [7].

(2) Two points in Section 4 of the above paper need clarification. Firstly, the base change to  $B$  that occurs at the beginning needs to be a separable one if the assumption about the Kodaira–Spencer map is to be preserved. Secondly, the descent argument near the bottom of p. 52 only works when the map  $Y \rightarrow X_B$  is also separable. Therefore, the proof from the beginning of Section 4 up to the italicized statement on p. 53 is incomplete. We need to correct as follows:

If the map  $Y \rightarrow X$  is separable, so is the base change  $B \rightarrow S$  and the map  $Y \rightarrow X'$  and the argument is correct as it stands. If  $Y \rightarrow X$  is inseparable, we can factor it (after blowing up, if necessary) to  $Y \rightarrow Y' \rightarrow X$  where  $\rho: Y \rightarrow Y'$  is purely inseparable and  $g: Y' \rightarrow X$  is separable. The map  $P: T \rightarrow X$  lifts to  $t_P: T \rightarrow Y$ ; let  $t': T \rightarrow Y'$  denote the composition  $\rho \circ t_P$ . Now, since  $\rho: Y \rightarrow Y'$  is purely inseparable, there is a non-trivial kernel for the map  $\Omega_{Y'} \rightarrow \rho_*(\Omega_Y)$ . Let  $L' \subset \Omega_{Y'}$  be a saturated line subsheaf contained in this kernel. Then the map  $t'^*\Omega_{Y'} \rightarrow \Omega_T$  factors through  $t'^*\rho_*(\Omega_Y)$ :

$$t'^*\Omega_{Y'} = t_P^*\rho^*\Omega_{Y'} \rightarrow t_P^*\rho^*\rho_*(\Omega_Y) \rightarrow t_P^*\Omega_Y \rightarrow \Omega_T.$$

In particular,  $t'^*L' \hookrightarrow t'^*\Omega_{Y'} \rightarrow \Omega_T$  is zero. But since  $Y' \rightarrow X$  is separable, so that  $\Omega_{Y'}$  and  $g^*\Omega_X$  are generically equal, there is also a line subsheaf  $Q \subset g^*(\Omega_X)$  which also vanishes when pulled-back to  $T$  and mapped down to  $\Omega_T$ . Now we can apply the descent of the proof in Section 4.

### Acknowledgement

I am grateful to the referee for kindly suggesting improvements in the first version of this erratum.

<sup>★</sup> *Composito Math.* **105** (1997), 43–54.