V.P.Arkhipova, T.A. Lozinskaya, E.I. Moskalenko, and T.G. Sitnik Sternberg Astronomical Institute Moscow, USSR

ABSTRACT. Spectral and monochromatic observations of two thin-filamentary nebulae S22 and YM29 have been carried out in the lines [O III] 5007A, [N II] 6584A and [S II] 6717+6732A. Radial stratification of the emission typical for photoionization excitation has been found: displacement between [O III] - bright regions and [S II] - and [N II] - bright regions is equal $\Delta R \sim 0.1R$ (about 0.01 to 0.2 pc), the first ones being closer to the nuclei. Both nebulae are characterized by more diffuse morphology in the [O III] line and by thin-filamentary one in the [N II] and [S II] lines.

Both objects seem to be old planetary nebulae of PNI type probably having stellar wind from their nuclei.

Optical 4000-7000A and UV 1150-3150A spectra of YM29 nucleus have been analyzed in looking for stellar wind indicators. The spectra seem to be similar to sd0 spectra with black-body temperature of 58000 ± 6000 K. Overall spectral structure may probably show some hints of PCyg-type lines, but one needs to obtain spectra with better resolution and statistics. The nebula YM29 and its nucleus have been also observed in X-ray range 0.2-3.5 KeV, but clear indications on strong wind have not been found. There are neither compact nor diffuse sources with the 3σ upper limit for 0.2-3.5 KeV flux: $F_{\text{nucl}} \le 1.1 \cdot 10^{-13}$ ergs/cm²s and $F_{\text{YM29}} \le 8 \cdot 10^{-14}$ ergs/cm²s under "standard" spectrum $J(E) \sim E^{-1.5}$ photons/cm²s and $N_{\text{H}} = 6 \cdot 10^{19}$ cm⁻².

The detailed results are given in Arkhipova, V.P., Lozinskaya, T.A., Moskalenko, E.I., 1986, Pisma Astron. Zh. (Sov. Astron. J. Lett.), v. 12, p. 890; Lozinskaya, T.A., Sitnik, T.G., Toropova, M.S., 1984, Pisma Astron. Zh. (Sov. Astron. J. Lett.,), v. 10, p. 122; Lozinskaya, T.A., Sitnik, T.G., Toropova, M.S., 1986, Astron. Zh. (Sov. Astron. J.), v. 63, p. 255.