Article: EPA-1675

Topic: P04 - Child and Adolescent Psychiatry

THE DIFFERENTIAL MODEL OF PSYCHOLOGICAL CORRECTION INCLUDED THE NATURAL AND PREMATURE FACTORS OF MENTAL AND BEHAVIORAL DISORDERS IN CHILDREN AND ADOLESCENTS

I. Vashkite<sup>1</sup>, B. Mykhaylov<sup>1</sup>, M. Vodka<sup>2</sup>, T. Aliieva<sup>1</sup>

<sup>1</sup>Psychotherapy, Kharkov Medical Academy of Postgraduate Education Kharkov, Kharkov, Ukraine; <sup>2</sup>Director, International Charity Foundation «Alexander Feldman Fund», Kharkov, Ukraine

**Educational Objectives**. Nonpsychotic mental disorders in children, bio-psycho-social assistance to children with mental and behavioral disorders, Center for comprehensive psychosocial rehabilitation of children, medical and psychological support, multi-modal system of providing medical and psychological assistance, psychotherapy is combined with animal-assisted therapy, psychocorrectional work in the «in / out door» system.

Purpose. Develop a medical and psychological treatment system, support and prophylaxis disadaptive psychological disorders in children.

Methods. Diagnostic and advisory work, correctional work, rehabilitation work, social and psychological work, methodical work.

**Results.** After the medical and psychological adjustment in four study groups (children with F90.0 – 1st group, F94 – 2nd group, F81 – 3nd group and F82 – 4th group) the following results were obtained:

-Positive '+' marker of social adaptation (the stabilization of family relationships, positive academic performance and normalization of interpersonal relationships with peers) in 1st group was 74,5%, in 2nd group – 86,0%, in 3nd group – 50,0% and in 4th group – 48,1%;

## The conclusions.

- 1. The system of medical and psychological support for children with mental and behavioral disorders.
- 2. The basic criteria for the effectiveness of medical and psychological treatment: the formation of a constructive type of family relationships and social adaptation.
- 3. Testing results on the primary stage of the work of the Center has demonstrated its effectiveness in 74.5% of children diagnosed with F 90, in 86.0% with F 94, in 50.0% with F 81 and in 48.1% with F 82, who underwent medical and psychological correction.