

## Special Issue Article

# The association between social emotional development and symptom presentation in autism spectrum disorder

Kyle B. Reid<sup>1</sup>, Lori-Ann R. Sacrey<sup>1</sup>, Lonnie Zwaigenbaum<sup>1</sup>, Sarah Raza<sup>1</sup>, Jessica Brian<sup>2</sup>, Isabel M. Smith<sup>3</sup>, Susan Bryson<sup>3</sup>, Vickie Armstrong<sup>3</sup>, Wendy Roberts<sup>4</sup>, Peter Szatmari<sup>5</sup>, Tracy Vaillancourt<sup>6</sup> and Caroline Roncadin<sup>7</sup>

<sup>1</sup>Department of Pediatrics, University of Alberta, Edmonton, Alberta, Canada; <sup>2</sup>Bloorview Research Institute and Department of Pediatrics, University of Toronto, Toronto, Ontario, Canada; <sup>3</sup>IWK Health Centre and Departments of Pediatrics and Psychology and Neuroscience, Dalhousie University, Halifax, Nova Scotia, Canada; <sup>4</sup>Integrated Services for Autism and Neurodevelopmental Disorders, Toronto, Ontario, Canada; <sup>5</sup>Department of Psychiatry, University of Toronto, Centre for Addiction and Mental Health, Hospital for Sick Children, Toronto, Ontario, Canada; <sup>6</sup>Counselling Psychology, Faculty of Education, University of Ottawa, Ottawa, Ontario, Canada and <sup>7</sup>Ron Joyce Children's Health Centre, Hamilton Health Sciences, Hamilton, Ontario, Canada

### Abstract

Understanding differences in social-emotional behavior can help identify atypical development. This study examined the differences in social-emotional development in children at increased risk of an autism spectrum disorder (ASD) diagnosis (infant siblings of children diagnosed with the disorder). Parents completed the Brief Infant-Toddler Social-Emotional Assessment (BITSEA) to determine its ability to flag children with later-diagnosed ASD in a high-risk (HR) sibling population. Parents of HR ( $n = 311$ ) and low-risk (LR; no family history of ASD;  $n = 127$ ) children completed the BITSEA when their children were 18 months old and all children underwent a diagnostic assessment for ASD at age 3 years. All six subscales of the BITSEA (Problems, Competence, ASD Problems, ASD Competence, Total ASD Score, and Red Flags) distinguished between those in the HR group who were diagnosed with ASD ( $n = 84$ ) compared to non-ASD-diagnosed children (both HR-N and LR). One subscale (BITSEA Competence) differentiated between the HR children not diagnosed with ASD and the LR group. The results suggest that tracking early social-emotional development may have implications for all HR children, as they are at increased risk of ASD but also other developmental or mental health conditions.

**Keywords:** autism spectrum disorder, BITSEA, high-risk cohort, infant sibling, ITSEA

(Received 4 November 2019; revised 11 March 2020; accepted 21 April 2020)

Social-emotional development includes the ability to express and manage positive and negative emotions, develop interpersonal relationships, as well as explore the environment to learn about one's surroundings (Pontoppidan, Niss, Pejtersen, Julian, & Vaever, 2017). Previous research has suggested that atypicalities in social-emotional development appear in community samples of children by 2 years of age and predict later mental health (Alink et al., 2006; Briggs-Gowan & Carter, 2008; Briggs-Gowan, Carter, Bosson-Heenan, Guyer, & Horwitz, 2006; van Zeijl et al., 2006; Wakschlag & Danis, 2009). For example, social-emotional atypicalities in 12- to 36-month-olds, as indexed by the Brief Infant-Toddler Social-Emotional Assessment (BITSEA), predicted children who later met diagnostic criteria for mental health disorders at elementary school entry (Briggs-Gowan & Carter, 2008).

Social-emotional development can be difficult to measure in children under the age of two. Infancy and toddlerhood are marked

by rapid changes in all areas of development, including social-emotional aspects, but it is difficult to separate typical (e.g., tantrums, separation anxiety) from atypical social-emotional development, as these behaviors may fall along a continuum, from typical to atypical, depending on contextual factors (Shields, Cicchetti, & Ryan, 1994; Lasch, Wolff, & Elison, 2019). Nevertheless, there is an increasing awareness of the need for monitoring social-emotional development in young children, as early problems have been associated with decreased social competence in preschool-aged children (Schmidt, Demulder, & Denham, 2010), poor academic performance in school-aged children (Campbell, Spieker, Burchinal, & Poe, 2006), and later mental health challenges (Campbell, Shaw, & Gilliom, 2000; Campbell et al., 2006; Fox, 2004; Shaw, Keenan, & Vondra, 1994). Early interventions for preschool-aged children for social-emotional problems show efficacy in improving later socialization and academic performance (Gross et al., 2003; Kelleher, Campo, & Gardner, 2006; Reid, Webster-Stratton, & Baydar, 2004; Thomas & Zimmer-Gembeck, 2007).

The BITSEA (Briggs-Gowan et al., 2006) is a 42-item parent-completed questionnaire for children between the ages of 11 and 48 months used to rate social and emotional problem behavior and social competence in the general population. Previous studies using the BITSEA have shown its utility in

**Author for correspondence:** Lonnie Zwaigenbaum, Department of Pediatrics, University of Alberta; Autism Research Centre, Glenrose Rehabilitation Hospital, Edmonton, Alberta, T5G 0B7. E-mail: [lonnie.zwaigenbaum@ahs.ca](mailto:lonnie.zwaigenbaum@ahs.ca)

**Cite this article:** Reid KB et al (2020). The association between social emotional development and symptom presentation in autism spectrum disorder. *Development and Psychopathology* 32, 1206–1216. <https://doi.org/10.1017/S0954579420000711>

predicting mental health challenges and parent and teacher ratings of psychopathology in children screened before age 4 years (Briggs-Gowan & Carter, 2008; Briggs-Gowan et al., 2013; Giserman Kiss et al., 2017). The BITSEA developers have added scales specific to autism spectrum disorder (ASD) in an attempt to support its use as an early detection tool for ASD (Briggs-Gowan et al., 2006; Gardner et al., 2013; Giserman Kiss & Carter, 2017). They reported acceptable levels of sensitivity (>.70) for distinguishing children with ASD from children without ASD children under 4 years old (Giserman Kiss, Feldman, Sheldrick, & Carter, 2017; Kruizinga et al., 2014). Although the hallmarks of ASD are impairments in social communication and restricted, repetitive behavior and interests (American Psychiatric Association, 2013), social-emotional difficulties have been recalled by parents of children with ASD (De Giacomo & Fombonne, 1998) and prospectively noted by parents of infant siblings of children who are later diagnosed with ASD (Sacrey et al., 2015) by their first birthday.

Understanding the associations between early social-emotional development and ASD symptom presentation is important, as social-emotional atypicalities often co-occur with ASD. This literature cited above noted accuracy for identifying ASD in children using a social-emotional development questionnaire. This literature, however, only included children who are already diagnosed with ASD and compared them to a cohort of typically developing children. The utility of the BITSEA to identify social-emotional atypicalities in infant siblings of children diagnosed with ASD has yet to be explored. These younger siblings are at increasing risk of also being diagnosed with ASD as the number of older siblings with ASD increases (16% for simplex families to 36% for multiplex families; McDonald et al., 2019). Furthermore, examining social-emotional difficulties in siblings of children with ASD is important because they are also at heightened risk for attention deficit hyperactivity disorder, conduct problems, anxiety disorders, and other neurodevelopmental and mental health conditions (Jokiranta-Olkoniemi et al., 2016). In this study, we assessed whether parent endorsement of social-emotional behavioral problems and competencies, as captured by the BITSEA, differentiated between high-risk (HR) toddlers who later received an ASD diagnosis (HR-ASD), HR toddlers who did not receive an ASD diagnosis (HR-N), and low-risk (LR) controls. We also examined the association between social-emotional development and clinical presentation at 36 months. We predicted that parents of children diagnosed with ASD at age 3 years would report more social-emotional atypicalities at 18 months compared to parents of children who would not be diagnosed with ASD.

## Methods

### Participants

Infant siblings of children with ASD were recruited from families attending one of five multidisciplinary ASD diagnostic and treatment centers in Canada: the Glenrose Rehabilitation Hospital in Edmonton, the Hospital for Sick Children and Holland Bloorview Kids Rehabilitation Hospital in Toronto, the IWK Health Centre in Halifax, and the McMaster Children's Hospital in Hamilton. This study was approved by the research ethics boards at each institution, and all families gave written informed consent before enrollment into the study. This study was part of a larger multisite prospective longitudinal study exploring early development in infant siblings of children with

ASD, including social-emotional development, language and cognitive ability, and symptom presentation in infant siblings who themselves also receive a diagnosis not receive a diagnosis of ASD (see Brian et al., 2008 for more details).

For the HR group, diagnosis of ASD in the older sibling (i.e., proband) was confirmed through a review of diagnostic records or clinical assessment using *Diagnostic and statistical manual of mental disorders* (fourth edition, text revision) (DSM-IV-TR) criteria. The HR infant siblings and probands had no identifiable neurological conditions, genetic conditions, or severe sensory or motor impairments. LR controls were recruited from local communities on the basis that they had no first- or second-degree relatives with an ASD diagnosis. All infant participants were born at 36–42 weeks' gestation and had a birth weight greater than 2500 g.

Children from the larger HR cohort were included in this study if (a) they had undergone a 3-year diagnostic assessment, and (b) their parents had completed an Infant-Toddler Social Emotional Assessment (ITSEA; from which the BITSEA items were drawn) at 18 months. Of the 465 HR children and 178 LR children with 3-year follow-up, 154 and 50, respectively, did not have a completed ITSEA and were excluded. The children without ITSEAs had completed their 18-month assessment prior to inclusion of the ITSEA in the study protocol. We adopted Kruizinga et al.'s (2014) exclusion criteria recommending that children who were missing more than three subscale items be excluded from analyses ( $n = 7$ ; 2 HR-ASD, 5 HR-N). Overall, data from 311 HR infant siblings and 128 LR controls were included in the current analyses. Participant characteristics are displayed in Table 1.

### Measures

BITSEA (Briggs-Gowan, Carter, Irwin, Wachtel, & Cicchetti, 2002) is a 42-item parent questionnaire designed to screen for social-emotional and behavioral problems, in addition to delays in social-emotional competence (Briggs-Gowan et al., 2006). The BITSEA is the brief version of the ITSEA (Briggs-Gowan & Carter, 2000). Our sample of parents completed the ITSEA when their children were 18 months of age, and items that comprise the BITSEA were extracted (as per Kruizinga et al., 2014). Each item is rated on a three-point Likert scale: 0 = *not true/rarely*, 1 = *somewhat true/sometimes*, and 2 = *very true/often*. As a Level 1 screen (Briggs-Gowan et al., 2002), the BITSEA has two subscales, Social Competency (hereafter, Competency) and Problem Behaviors (hereafter, Problems). Subscales were calculated if fewer than three items were missing (Giserman Kiss & Carter, 2017). For the scales with two or fewer missing items, 3-year outcome group mean values for the missing item were placed in the empty cell, resulting in replacement of less than 1% of values for each group (0.09% of cells in the LR group and 0.37% of cells in the HR group).

To evaluate potential as a Level 2 screen (i.e., targeted to children at-risk on the basis of specific symptoms or positive family history; Zwaigenbaum et al., 2015), we used the criteria of Gardner et al. (2013) to calculate the ASD-specific subscales, ASD Competency (items 1, 10, 13, 15, 22, 25, 29, & 31), ASD Problems (items 9, 14, 21, 35, 36, 37, 38, 39, & 40), Total ASD Screening Score (ASD Problems—ASD Competency), and Red Flags (items 2, 14, 18, 24, 32, 34, 35, 36, 37, 38, 39, 40, 41, 42; Gardner et al., 2013). The items included in each subscale are displayed in Table 2. As can be seen, the items comprising ASD Social Competence ( $n = 8$ ) and BITSEA Social Competency ( $n = 11$ ) are very similar, whereas the items comprising BITSEA Problem Behaviors ( $n = 31$ ) are more inclusive of a range of

**Table 1.** Participant characteristics

Characteristics	LR (a)		HR-N (b)		HR-ASD (c)		Statistics		
Sex	<b>boy:girl</b>		<b>boy:girl</b>		<b>boy:girl</b>		$\chi^2$	<i>p</i>	<b>Post hoc</b>
M:F	67:61		115:112		61:23		12.62	<0.01	c > a,b boys
Age at Assessment	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>F</b>	<b>p</b>	<b>Post hoc</b>
18 months	18.47	0.70	18.40	0.59	18.36	0.51	0.96	0.38	-
36 months	39.60	4.26	39.37	3.92	39.12	4.12	0.51	0.60	-
ADI-R									
Total score	4.38	3.90	6.71	5.71	23.44	10.73	229.04	<0.001	c > a,b; b > a
ADOS severity score									
SA	2.14	1.40	2.58	1.68	6.63	1.87	225.14	<0.001	c > a,b;
RRB	3.52	2.41	4.83	2.49	7.48	2.10	69.41	<0.001	c > a,b; b > a
Total	1.78	1.27	2.45	1.64	6.75	1.84	284.58	<0.001	c > a,b; b > a
MSEL standard scores									
ELC	119.59	15.44	108.84	17.37	90.16	22.50	65.80	<0.001	c < a,b; b < a
Visual Reception	119.89	15.14	115.08	17.21	97.57	28.75	35.66	<0.001	c < a,b
Fine motor	112.97	18.95	101.45	21.39	84.95	20.08	47.66	<0.001	c < a,b; b < a
Receptive Language	113.26	14.29	104.57	14.35	89.17	18.89	61.93	<0.001	c < a,b; b < a
Expressive Language	112.88	13.38	105.49	14.80	90.71	17.93	54.78	<0.001	c < a,b; b < a
Vineland standard scores									
ABC	100.59	14.09	94.50	14.33	85.04	15.69	17.03	<0.001	c < a,b; b < a
Communication	110.26	10.86	103.24	14.00	88.27	14.71	41.22	<0.001	c < a,b; b < a
Daily living Skills	93.01	10.07	90.54	12.50	76.50	10.89	34.11	<0.001	c < a,b
Socialization	93.38	9.48	92.18	12.57	77.71	10.69	33.95	<0.001	c < a,b
Motor skills	100.59	14.09	94.50	14.33	95.04	15.69	17.02	<0.001	c < a,b

LR = low-risk toddlers (a); HR-N = high-risk toddlers who did not meet criteria for autism spectrum disorder (b); HR-ASD = high-risk toddlers diagnosed with autism spectrum disorder (c); SD = Standard Deviation; ABC = Adaptive Behavior Composite; ADI-R = Autism Diagnostic Interview—Revised; ADOS = Autism Diagnostic Observation Schedule; MSEL = Mullen Scales of Early Learning, Vineland = Vineland Adaptive Behavior Scales.

potentially problematic behaviors than the ASD Problem Behaviors ( $n = 9$ ), which are more characteristic of ASD.

#### *The Autism Diagnostic Observation Schedule*

(ADOS; Lord et al., 1989) is a semi-structured assessment with excellent interrater reliability; it uses standardized activities and ‘presses’ to elicit communication, social interaction, imaginative use of play materials, and repetitive behavior (Lord et al., 1989; Lord et al., 2000). We used the updated ADOS algorithms of Gotham, Risi, Pickles, and Lord (2007), organized into two domains: Social Affect (including communication and social items) and Restricted Repetitive Behaviors. The original ADOS consists of four modules, each of which is appropriate for individuals of differing language levels. The current study used Modules 1 (minimal or no language), 2 (regular use of non-echoed 3-word phrases) and 3 (fluent language). Comparability across modules and language levels was optimized using the 36-month ADOS severity metric (Gotham, Pickles, & Lord, 2009). The ADOS was administered at age 36 months by research-reliable raters.

#### *The Autism Diagnostic Interview-Revised*

(ADI-R; Lord, Rutter, & Le Couteur, 1994) is an investigator-directed interview that elicits information regarding social

development, verbal and non-verbal communication skills, and the presence of repetitive, stereotyped interests and behavior required to make an ICD-10 or DSM-IV-TR diagnosis of ASD. The questions are designed to distinguish qualitative impairments from developmental delays. The ADI-R discriminates well between ASD and other forms of developmental disability, and inter-rater reliability is excellent (Lord et al., 1994). The ADI-R was administered to parents of 36-month-olds by research-reliable interviewers.

#### *The Mullen Scales of Early Learning*

(MSEL; Mullen, 1995) consists of five scales, four of which (Visual Reception, Receptive Language, Expressive Language, and Fine Motor) assess nonverbal, cognitive, and language ability, while the fifth scale measures gross motor development (from 0 to 29 months only). An Early Learning Composite is calculated based on scores from the first four scales for children aged 0–69 months. Inter-rater and test-retest reliability are excellent (Mullen, 1995). The MSEL was administered at 36 months of age.

#### *The Vineland Adaptive Behavior Scales*

(Vineland; Sparrow, Balla, Cicchetti, Harrison, & Doll, 1984) is a semi-structured parent interview designed to assess adaptive behavior across four subdomains outlined by typical

**Table 2.** Items included in each of the BITSEA and ASD subscales

ASD Social Competence	ASD Problem Behaviors
Shows pleasure when they succeed	Has less fun than other children
Looks for you when upset	Does not react when hurt
Looks at you when you say their name	Has trouble adjusting to changes “Spaces out”—unaware of surroundings
Is affectionate with loved ones	Avoids physical contact
Points to show you something far away	Does not make eye contact
Tries to help when someone is hurt (give a toy)	Puts things in a special order, over and over
Imitates playful sounds when you ask them to	Repeats a particular movement, over and over
Hugs or feeds doll or stuffy	Repeats the same action, over and over
BITSEA Social Competence	BITSEA Problem Behaviors
All ASD Social Competence items	All ASD Problem Behaviors items
Follows rules	Takes a while to feel comfortable in new place
Can pay attention for a long time (not including TV)	Gets hurt so often you can't take your eyes off them
Plays well with other children	Acts aggressive when frustrated
Red Flags	Wakes up at night and needs help to fall asleep again
Gets hurt so often you can't take your eyes off them	Cries or tantrums until exhausted
Does not react when hurt	Afraid of certain animals, things, or places
Runs away in public places	Cries or hangs onto you when you try to leave
Gags or chokes on food	Worries a lot or is very serious
When upset, gets still, doesn't move	Won't touch certain things because of how they feel
Seems very unhappy, sad, or depressed	Runs away in public places
Hurts self on purpose	Often gets very upset
“Spaces out”—unaware of surroundings	Gags or chokes on food
Avoids physical contact	Has trouble falling asleep or staying asleep
Does not make eye contact	When upset, gets still, doesn't move
Puts things in a special order, over and over	Refuses to eat
Repeats a particular movement, over and over	Is destructive and breaks things
Repeats the same action, over and over	Hits, shoves, or kicks other children
Eats things that are not edible	Hits, bites, and kicks you
	Seems very unhappy
	Purposefully tries to hurt you
	Hurts themselves on purpose
	Eats things that are not edible

developmental milestones that are anchored to specific ages - Communication, Daily Living, Socialization, and Motor skills (with the last domain limited to children younger than 30 months). The scale has excellent reliability and concurrent validity and is sensitive to impairments experienced by children with ASD (Carter et al., 1998; Volkmar, Carter, Sparrow, & Cicchetti, 1993). The Vineland or Vineland II was administered at 36 months of age.

### Diagnostic Assessment

Each participant underwent an independent gold-standard diagnostic evaluation at age 3 years conducted by a clinician blind to previous study visits. Diagnoses were assigned using DSM-IV-TR criteria, based on the best judgement of the clinician (developmental pediatrician, child psychiatrist, or clinical psychologist, all with at least 10 years' diagnostic experience),

considering information from the ADOS, ADI-R, and concurrent developmental assessments using the MSEL & Vineland.

### Statistical Analyses

A series of one-way ANOVAs was conducted to compare the BITSEA subscales (BITSEA Problems, BITSEA Competence, ASD Problems, ASD Competence, Total ASD Score, and Red flags) by 3-year diagnostic outcome group (HR-ASD, HR-N, LR). Post hoc comparisons were explored using Bonferroni corrections. We included comparisons between the LR groups and the two HR groups for two reasons. First, previous reports of social-emotional differences in children with ASD included a group of already diagnosed children and typically developing peers. In this respect, our HR-ASD group serves as the diagnosed group and the LR controls serves as an analogue of typically developing peers. Second, for the HR siblings who were not diagnosed with ASD, we wanted to clarify their place on a continuum of social-emotional behavior. That is, if they differ from HR siblings who were diagnosed with ASD (*rated as showing more social competence and less problem behavior*), are they also different from LR controls (*rated as showing less social competence and more problem behavior*) due to the heightened risk for other developmental / mental health conditions in siblings of children with ASD compared to typically developing siblings (Jokiranta-Olkonienmi et al., 2016).

To examine the association between social-emotional development and clinical outcomes in the HR sample, Pearson correlations and receiver operator characteristics (ROC) were calculated. All six BITSEA subscales were correlated with ADOS severity scores, ADI-R Total algorithm scores, MSEL subscales and Early Learning Composite, and Vineland subscales and Adaptive Behavior Composite. To control for multiple comparisons, we used Benjamini and Hochberg (1995) corrections. In this method, the  $p$ -values are ordered smallest to largest. The alpha level for each test is then set at  $(k \cdot \alpha) / m$ , with  $k$  corresponding to the  $p$ -value's rank (lowest  $p = 1$ ) and  $m$  corresponding to the number of comparisons, which in this case was 13 (each subscale was run separately). This method decreases the chance of false positives; comparisons stop once one of the  $t$ -tests is rejected (this method uses ' $q$ ' rather than ' $p$ ' to denote the critical alpha level). In addition, area under the curve (AUC) was calculated to assess 18-month BITSEA's sensitivity and specificity with respect to ASD diagnosis. To examine specifically the BITSEA's potential predictive properties within an HR context, ROC analyses were limited to the HR cohort (ASD + versus ASD -). The AUC was calculated to represent an accuracy index, with higher AUC values representing better sensitivity and specificity (AUC of 0.50 = chance relationship; AUC 0.70–0.90 = moderate,  $\geq 0.90$  high accuracy; Akobeng, 2007). Youden indexes (defined as the maximum vertical distance between the ROC curve and the chance/diagonal line [Youden's Index ( $J$ ) = sensitivity + specificity - 1]) were calculated to determine optimal cut-scores for each subscale (Akobeng, 2007). In this method, the highest  $J$  value represents the recommended cut score (Akobeng, 2007). Estimates of screening accuracy were calculated, including: (1) sensitivity, the proportion of children with ASD who were correctly classified by the BITSEA; (2) specificity, the proportion of children not diagnosed with ASD correctly classified by the BITSEA; (3) positive predictive value (PPV), the proportion of children exceeding the BITSEA cut-point who were diagnosed with ASD; and (4) negative predictive

value (NPV), the proportion of children who did not exceed the BITSEA cut-point who were not diagnosed with ASD (Fischer, Bachmann, & Jaeschke, 2003). Statistical analyses were completed using SPSS version 25.

## Results

### Participant Characteristics

Three groups were identified for comparison based on the 36-month diagnostic assessments: (1) HR infant siblings who received a diagnosis of ASD ('HR-ASD';  $n = 84$ ; 61 boys and 23 girls); (2) HR infant siblings who did not receive a diagnosis of ASD ('HR-N';  $n = 227$ ; 115 boys and 112 girls); and (3) LR controls who did not receive a diagnosis of ASD ('LR';  $n = 128$ ; 67 boys and 61 girls). A significant sex difference ( $\chi^2 = 12.62$ ,  $p = 0.02$ ) showed a higher boy-to-girl ratio in the HR-ASD group than in the LR and HR-N groups, who did not differ ( $ps < 0.01$  and  $= .42$ , respectively). No differences were found relating to the child's age at completion of the BITSEA ( $F(2,438) = 0.96$ ,  $p > 0.05$ ).

As expected, there were both overall and pair-wise group differences between the HR-ASD, HR-N, and LR controls on the ADOS Social Affect ( $ps < .001$ ), Restricted Interests and Repetitive Behavior ( $ps < .001$ ), and Overall Severity scores ( $ps < 0.001$ ), ADI-R Total score ( $ps < 0.002$ ), MSEL standard scores ( $ps < 0.001$ ), and Vineland standard scores ( $ps < 0.001$ ), with the exception of the Daily Living Skills standard score, which did not differentiate between the HR-N and LR groups ( $p = .14$ ). Overall, HR-ASD toddlers had higher ADOS and ADI-R scores and lower scores on the Vineland and MSEL relative to the HR-N and LR groups. Descriptive data on these 36-month measures are summarized in Table 2.

### Completers versus Non-Completers

Independent-sample  $t$ -tests were conducted to determine whether clinical characteristics differed between the BITSEA completers and non-completers at 36 months. There was no significant difference between the two groups for ADOS severity scores at 36 months [SA ( $t(637) = -0.57$ ,  $p = .57$ ); RRB ( $t(364.19) = -1.93$ ,  $p = .054$ ); Total Severity ( $t(637) = -0.67$ ,  $p = .50$ ), nor the ADI-R Total score at 36 months ( $t(601) = -0.90$ ,  $p = 0.37$ ). Similarly, completers versus non-completers did not differ on MSEL Early Learning Composite ( $t(616) = -0.90$ ,  $p = .37$ ) nor the Vineland Adaptive Behavior Composite ( $t(632) = 0.82$ ,  $p = .41$ ).

### BITSEA Subscales

An overall group difference was observed for each BITSEA subscale, including BITSEA Problems ( $F(2,436) = 17.62$ ,  $p < .001$ ), BITSEA Competence ( $F(2,436) = 43.60$ ,  $p < .001$ ), and Red Flags ( $F(2,436) = 25.53$ ,  $p < .001$ ). In addition, there was a group difference on the ASD-specific subscales, including ASD Problems ( $F(2,436) = 22.92$ ,  $p < .001$ ), ASD Competence ( $F(2,436) = 38.51$ ,  $p < .001$ ), and ASD Total Score ( $F(2,436) = 43.77$ ,  $p < .001$ ). Overall, the HR-ASD group had lower scores on two subscales (the Competencies) and higher scores on the remaining four subscales compared to the HR-N and LR groups ( $ps < .003$ ), who did not differ, except on BITSEA Competence ( $p < .001$ ). These results are summarized in Table 3.

### Associations between Social-Emotional Development and Clinical Presentation

To examine associations between children's social-emotional problems and competencies at 18 months of age and their clinical characteristics at 36 months of age, correlations were computed between BITSEA subscales and ADOS severity scores (SA, RRB, and overall severity scores), ADI-R algorithm scores, Vineland subscale scores and Adaptive Behavior Composite, and the MSEL subscale scores and Early Learning Composite. Because the ADOS was not designed to detect meaningful clinical variation in typically developing children, we have only included the HR group in the correlational analyses. Pearson's correlations were run for each of the HR groups separately (HR-N and HR-ASD) and multiple comparisons were corrected using Benjamini and Hochberg (1995) corrections. The results are presented in Table 4.

**HR-N Group.** There were no significant correlations between any BITSEA subscale and the ADOS severity scores or MSEL subscales or Early Learning Composite ( $ps > .01$ ).

There were significant correlations between the BITSEA and other parent-report measures. All BITSEA subscales were associated with the ADI-R Total score ( $ps < .01$ ). For the Vineland, (2) the Communication Scale was related to BITSEA Problems ( $ps = .002$ ), (3) the Daily Living Scale was related to BITSEA Competence ( $p = .001$ ), ASD Competence ( $p = .001$ ), and ASD Total Score ( $p = .002$ ), (3) the Motor Skills Scale was related to BITSEA Competence ( $p < .001$ ), ASD Competence ( $p < .001$ ), and ASD Total Score ( $p < .001$ ), and (4) the Vineland Adaptive Behavior Composite (ABC) was related to BITSEA Problems ( $p = .003$ ) and Red Flags ( $p = .005$ ). The Socialization Scale was not related to any BITSEA subscale ( $ps > .01$ ).

**HR-ASD Group.** There were no significant correlations between any BITSEA subscale and the ADOS severity scores ( $ps > .05$ ). In contrast, there was a significant correlation between the ADI-R Total and the BITSEA Competence, Red Flags, ASD Competence, and ASD Total Score subscales ( $ps < .002$ ).

For the MSEL, there was a significant correlation between the Fine Motor and the Receptive Language subscales with the BITSEA Problems subscale ( $p = .007$  and  $p = .003$ , respectively). There were no significant relationships between the Expressive Language Scale, the Visual Reception Scale, or the Mullen Early Learning Composite and BITSEA subscales ( $p < .05$ ).

Significant correlations were also found for the Vineland subscales: (1) the Daily Living Skills scale was related to BITSEA Competence ( $p = .004$ ) and ASD Competence ( $p = .005$ ), and (2) the Motor Skills scale was related to BITSEA Competence ( $p = .004$ ) and ASD Competence ( $p = .007$ ). There were no significant relationships between the Communication Scale, the Socialization Scale, or the Vineland ABC and BITSEA subscales ( $p < .05$ ).

### Receiver Operating Characteristic (ROC) Curve Analyses

Analyses were completed on the HR group (HR-ASD and HR-N) to determine the ability of social-emotional development at 18 months to distinguish between HR children who would and would not be diagnosed with ASD at age 3. The AUC analyses for each subscale were significant (i.e., differed from 0.5) and

**Table 3.** Descriptive statistics for the Brief Infant-Toddler Social Emotional Assessment (BITSEA) subscales

BITSEA subscales	LR (a)		HR-N (b)		HR-ASD (c)		F	P	Post hoc
	Mean	SD	Mean	SD	Mean	SD			
BITSEA Competence	17.21	2.49	15.85	3.22	12.98	4.19	43.60	<0.001	c > a,b; b > a
BITSEA Problem Behavior	6.52	4.62	7.41	5.09	10.99	7.77	17.62	<0.001	c > a,b
ASD Competence	13.25	1.93	12.51	2.45	10.23	3.32	38.51	<0.001	c > a,b
ASD Problem Behavior	0.85	1.20	0.65	1.27	2.02	2.62	22.92	<0.001	c > a,b
Total ASD Screening score	-12.40	2.45	-11.86	3.14	-8.21	5.06	43.78	<0.001	c > a,b
Red Flags	1.70	1.69	1.70	1.94	3.69	3.62	25.53	<0.001	c > a,b

Note. LR = low-risk toddlers (a); HR-N = high-risk toddlers who did not meet criteria for autism spectrum disorder (b); HR-ASD = high-risk toddlers diagnosed with autism spectrum disorder (c)

ranged from 0.64 (BITSEA Problem) to 0.73 (Total ASD Screening score).

Youden Index analysis [(sensitivity + specificity) - 1] was used to identify the optimal BITSEA subscale cut-scores at 18 months of age. The ASD-specific scales yielded relatively poor sensitivity (0.40–0.52) but high specificity (0.84–0.90) values. The non-ASD Competency and Problems scales had similar results; that is, relatively poor sensitivity (0.38–0.46) and high specificity (0.84–0.87). In contrast, the Red Flags scale had higher sensitivity than ASD and non-ASD scales (0.71) but relatively poor specificity (0.60). When comparing the optimal parameters, discrimination between HR children who would and would not be diagnosed with ASD was best achieved by the Total ASD screening score, as determined by the highest AUC value, Youden Index, and sensitivity value among the relevant BITSEA subscales. Table 5 displays the results of ROC analyses for each subscale.

### True Positives versus False Negatives

We were interested in determining clinical differences between the children with ASD who were correctly (true positive) versus incorrectly (false negative) classified by the Total ASD screening cut-off of 9 on the BITSEA. We used Mann-Whitney U tests to compare scores on the ADOS, ADI-R, MSEL, and Vineland for correctly versus incorrectly classified children. Overall, correctly classified children with ASD had higher scores on the Vineland Adaptive Behavior Composite ( $M \pm SD = 89.39 \pm 15.70$  versus  $76.40 \pm 13.85$ ;  $Z = -2.41$ ,  $p = .016$ ) and lower scores on the ADI-R Total algorithm score ( $M \pm SD = 20.42 \pm 10.69$  versus  $27.14 \pm 9.70$ ;  $Z = -2.87$ ,  $p = .004$ ).

There were no statistically significant differences between correctly and incorrectly classified children with ASD on the ADOS SA ( $Z = -1.3$ ,  $p = .17$ ), ADOS RRB ( $Z = -0.023$ ,  $p = .98$ ), ADOS Total Severity score ( $Z = -1.11$ ,  $p = .26$ ), nor the MSEL Early Learning Composite ( $Z = -0.068$ ,  $p = .95$ ). Similarly, no differences were observed between the groups on any of the MSEL ( $ps > .30$ ) or Vineland subscales ( $ps > .06$ ).

### Discussion

The aim of this study was to examine, in infants at high risk, the associations between early social-emotional behavior and competencies and later clinical presentation with respect to ASD in a HR population. Primary caregivers completed the BITSEA at 18 months of age and all children underwent an ASD diagnostic

assessment at age three. There were three main results: (1) social-emotional development, as indexed by BITSEA subscales, differentiated children with ASD from children without ASD at 18 months of age, with only one subscale (BITSEA Competence) differentiating between the HR-N and LR groups; (2) social-emotional development was associated with clinical presentation at age three, as indexed by significant associations between BITSEA subscales and ADI-R algorithm scores in both HR groups; and (3) screening cut-off thresholds for social-emotional development, as indexed by the BITSEA, did not meet recommended criteria for screening of ASD (Zwaigenbaum et al., 2015). The results demonstrate that the BITSEA provides clinically meaningful information regarding social and emotional problems in HR siblings but is not an adequate stand-alone screen for ASD based on individual classification metrics.

Parents of toddlers later diagnosed with ASD endorsed more social-emotional problems compared to parents of toddlers without ASD at 18 months of age. Five of the six BITSEA subscales show good discrimination of ASD from non-ASD, with only one subscale, BITSEA Competence, differentiating between the non-diagnosed children (HR-N and LR). These results are similar to previous reports using the BITSEA as an index of social-emotional development in children at risk for ASD and children with confirmed diagnoses of ASD. Gardner et al. (2013) had parents complete the BITSEA when their children were 12 and 24 months of age and compared children based on elevated scores on the Modified Checklist for Autism in Toddlers (M-CHAT). Parents of children in the elevated M-CHAT scores group endorsed significantly more social-emotional development atypicalities (as indexed by the same BITSEA subscales used in this study) at both time points. Higher endorsements of social-emotional atypicalities were also reported for children with ASD compared to typically developing controls (Boone, Brown, & Keim, 2018; Giserman Kiss et al., 2017; Karabekiroglu, Briggs-Gowan, Carter, Rodopman-Arman, & Akbas, 2010; Kruizinga et al., 2014; Raza et al., 2019). Combined, these results suggest that the BITSEA can help parents identify social-emotional difficulties their children may be experiencing, which can inform clinical referrals and potentially, diagnoses.

Exploration of the associations between early social-emotional development and later clinical outcomes indicated a strong relationship between social-emotional problems and competencies and ADI-R algorithm scores in both HR groups. Similarly, Daily Living Skills and Motor subscales on the Vineland were associated with both the BITSEA and ASD Competence scales in both HR groups. These results corroborate the valuable

**Table 4.** Relationships between Brief Infant-Toddler Social Emotional Assessment (BITSEA) subscales and clinical characteristics in the high-risk sample

Clinical characteristic	BITSEA Competence	BITSEA Problems	Red Flags	ASD Competence	ASD Problems	ASD total score
<b>high-risk non-asd</b>						
<b>Range N</b>	125–222	125–222	125–222	125–222	125–222	125–222
<b>Significance at</b>	q<.01	q<.01	q<.007	q<.01	q<.003	q<.01
<b>ADOS SA</b>	–.06	–.04	.003	–.04	–.04	.02
<b>ADOS RRB</b>	–.10	.04	.09	–.10	.04	.10
<b>ADOS total severity</b>	–.04	–.07	–.02	–.04	–.04	.02
<b>ADI-R total score</b>	<b>–.41*</b>	<b>.34*</b>	<b>.40*</b>	<b>–.40*</b>	<b>.31*</b>	<b>.44*</b>
<b>MSEL fine motor</b>	.14	–.06	–.06	.11	–.09	–.12
<b>MSEL visual reception</b>	.03	–.06	–.06	.04	–.05	–.06
<b>MSEL receptive language</b>	.12	–.02	–.03	.10	–.05	–.10
<b>MSEL expressive language</b>	.005	.01	–.03	.05	.03	–.03
<b>MSEL ELC</b>	.08	–.14	–.10	.06	–.06	–.07
<b>Vineland communication</b>	.11	<b>–.28*</b>	–.21	.10	–.12	–.13
<b>vineland daily living skills</b>	<b>.29*</b>	–.18	–.18	<b>.28*</b>	–.12	<b>–.27*</b>
<b>Vineland socialization</b>	.22	–.07	–.10	<b>.23*</b>	–.04	–.20
<b>Vineland motor skills</b>	<b>.37*</b>	–.16	–.17	<b>.33*</b>	–.14	<b>–.32*</b>
<b>Vineland ABC</b>	–.04	<b>.26*</b>	<b>.25*</b>	–.006	–.24	.10
<b>High-Risk with ASD</b>						
<b>Range N</b>	48–84	48–84	48–84	48–84	48–84	48–84
<b>Significance at</b>	q<.008	q<.008	q<.004	q<.01	q<.001	q<.003
<b>ADOS SA</b>	–.02	–.14	–.08	–.07	–.02	.04
<b>ADOS RRB</b>	.02	–.04	–.04	–.02	.04	.04
<b>ADOS Total Severity</b>	.04	–.21	–.13	–.03	.04	.004
<b>ADI-R Total Score</b>	<b>–.40*</b>	.33	<b>.35*</b>	<b>–.43*</b>	.27	<b>.42*</b>
<b>MSEL fine motor</b>	.09	<b>.29*</b>	.21	.13	.15	–.003
<b>MSEL visual reception</b>	.10	–.08	–.08	.08	–.05	–.08
<b>MSEL receptive language</b>	.09	<b>.32*</b>	.27	.11	.17	.02
<b>MSEL expressive language</b>	.09	–.05	–.07	.05	–.05	–.06
<b>MSEL ELC</b>	.09	–.15	–.19	.03	–.20	–.12
<b>Vineland Communication</b>	.20	–.31	–.29	.16	–.29	–.25
<b>Vineland daily living skills</b>	<b>.41*</b>	–.10	–.08	<b>.40*</b>	–.14	–.32
<b>Vineland Socialization</b>	.28	–.04	.009	.25	.005	–.16
<b>vineland motor skills</b>	<b>.41*</b>	–.11	–.10	<b>.38*</b>	–.09	–.29
<b>Vineland ABC</b>	.35	–.20	–.16	.27	–.17	–.26

**Note:** HR-N = high-risk toddlers who did not meet criteria for autism spectrum disorder; HR-ASD = high-risk toddlers diagnosed with autism spectrum disorder; ADI-R = Autism Diagnostic Interview—Revised; ADOS = Autism Diagnostic Observation Schedule; MSEL = Mullen Scales of Early Learning; ELC = Early Learning Composite; Vineland = Vineland Adaptive Behavior Scales; ABC = Adaptive Behavior Composite; p-values corrected for familywise error rate using Benjamini and Hochberg (1995) correction (\* = *q* significant) and Range N is the range of lowest to highest number of participants in each comparison.

contributions of parents with respect to their ability to identify early differences in children who later are classified as having behavioral problems, developmental delays, or ASD (Robins *et al.*, 2001; Glascoe, 2003; Sacrey *et al.*, 2015; Sacrey *et al.*, 2018). In contrast, no ADOS score was related to social-emotional development as measures by the BITSEA in either HR group. The lack of association between the BITSEA and ADOS may result from their different purposes (detecting social-emotional

development versus ASD symptoms), the narrow focus of items included in the ADOS severity algorithms, and the nature of the assessment context (i.e., a one-time observation over roughly a one-hour period) compared to the broader array of behavior queried on the BITSEA and the larger sampling frame from which parents can pull. This may also have contributed to the small number of associations between the BITSEA and the MSEL. It has been reported that parents and clinicians do not

**Table 5.** ROC characteristics of BITSEA subscales for the HR-ASD and HR-N groups only

Subscale	AUC (CI)	Cutoff	Sens	Spec	LHR+	LHR-	OR	J	PPV	NPV
ASD Problems	0.68 (0.61–0.75)	-	-	-	-	-	-	-	-	-
		1 <sup>a</sup>	0.42	0.86	3.05	0.33	4.52	0.28	0.53	0.80
		2	0.26	0.74	3.96	0.25	5.02	0.20	0.59	0.77
ASD Competence	0.71 (0.64–0.77)	8	0.27	0.93	3.88	0.26	4.97	0.20	0.59	0.78
		9 <sup>a</sup>	0.40	0.90	4.18	0.24	6.34	0.31	0.44	0.80
		10	0.49	0.81	2.58	0.39	4.08	0.30	0.49	0.81
ASD Total Score	0.73 (0.67–0.80)	-10	0.60	0.75	2.41	0.41	4.49	0.35	0.47	0.83
		-9 <sup>a</sup>	0.52	0.84	3.21	0.31	5.65	0.36	0.54	0.83
		-8	0.43	0.87	3.24	0.31	4.92	0.30	0.27	0.73
BITSEA Problems	0.64 (0.57–0.71)	10	0.40	0.79	1.91	0.52	2.53	0.19	0.41	0.78
		11 <sup>a</sup>	0.38	0.84	2.34	0.43	3.16	0.22	0.46	0.79
		12	0.32	0.86	2.28	0.44	2.89	0.18	0.46	0.77
BITSEA Competence	0.70 (0.63–0.77)	11	0.33	0.94	4.25	0.24	5.40	0.20	0.58	0.79
		12 <sup>a</sup>	0.46	0.87	3.51	0.28	5.69	0.33	0.56	0.81
		13	0.54	0.79	2.53	0.39	4.30	0.32	0.48	0.82
Red Flags	0.69 (0.62–0.76)	-	-	-	-	-	-	-	-	-
		1	0.71	0.60	1.80	0.56	3.81	0.32	0.40	0.85
		2	0.48	0.74	1.83	0.55	2.59	0.22	0.40	0.79

Note. AUC = area under the curve; CI = confidence interval (95%); LHR+ = positive likelihood ratio; Sens = sensitivity; Spec = specificity; LHR- = negative likelihood ratio; OR = odds ratio; J = Youden's index; PPV = positive predictive value; NPV = negative predictive value. <sup>a</sup>Shading indicates the optimal cutoff (via Youden's J). No statistics were generated for ASD Problems subscale with cut-off of 0 due to BITSEA questions being on 3-point Likert scale.

often agree with respect to signs of ASD (Stadnick et al., 2017), even when the two assessments (parent questionnaire and clinician observation) query the *same* ASD-related behavior (Sacrey et al., 2018). The BITSEA, ADI-R, and Vineland rely on parent report and thus share method variance, which may account for the stronger relationship compared to the ADOS and MSEL.

The similar pattern of associations between the BITSEA subscales and the clinical assessments for both HR groups suggests that the questionnaire may not be suited to differentiating ASD from non-ASD in an HR sample. We would temper this statement, however, by acknowledging that HR siblings of children with ASD are at increased risk of being diagnosed with other neurodevelopmental conditions compared to siblings of typically developing children, such as anxiety disorders (Shivers, Deisenroth, & Taylor, 2013; 2019) and ADHD (Ghirardi et al., 2018; Miller et al., 2019). As such, the similar pattern of relationships between the BITSEA and the clinical assessments may partially be due to parents recognizing social-emotional differences in their children who may be diagnosed with conditions such as attention deficit hyperactivity disorder, rather than ASD. With this in mind, the BITSEA may be a useful method of collecting and tracking social-emotional behaviors and differences in siblings of children with ASD.

Our ROC analyses produced sensitivity and specificity scores that are below recommended levels for screening (Cicchetti, Volkmar, Klin, & Showalter, 1995; Council on Children with Disabilities, 2006; Dumont-Mathieu & Fein, 2005; Volkmar et al., 1988). This is in contrast to previous reports of screening accuracy of the BITSEA, with sensitivity estimates ranging from .70 to .93 and specificity estimates ranging from .68 to .89

(Boone et al., 2018; Giserman Kiss et al., 2017; Karabekiroglu et al., 2010; Kruijzinga et al., 2014). Within the current group of children who were diagnosed later with ASD, we compared the clinical characteristics of those who were identified versus not identified by the Total ASD Score cutoff (subscale with highest AUC, sensitivity, and specificity). Surprisingly, the screen-positive children did not differ from the screen-negative children on the three ADOS severity scores, the Vineland and MSEL subscales, nor on the MSEL Early Learning Composite. The screen-positive children only differed on the Vineland Adaptive Behavior Composite (higher scores than screen-negative children) and ADI-R Total algorithm scores (lower scores than screen-negative children), in line with the correlation findings. Although the BITSEA lacked discriminatory ability to classify ASD, the BITSEA Competence scale differentiated between the HR-ASD and HR-N groups, as well as the HR-N and LR groups, suggesting that this scale may be able to index behaviors related to inherited risk of ASD. A similar relationship was found by Lasch et al. (2019), who observed a strong association between the ITSEA competence subscale and reciprocal social behavior measured with the Video-Referenced Rating of Reciprocal Social Behavior (vrRSB; Marrus et al., 2015), an index of heritable traits for ASD (Marrus et al., 2020).

These results highlight important considerations regarding social-emotional development in HR siblings. Although more social-emotional challenges are reported in HR siblings with ASD compared to non-diagnosed siblings, these differences are not specific to ASD. Furthermore, both the lack of differences on the ADOS and MSEL, as well as the counter-intuitive directions of differences on the ADI-R and Vineland (screen-positive



children had *better performance* on the measures than screen-negative children) provide further evidence for the lack of discriminatory ability of the BITSEA. We would be remiss not to point out that the previous studies compared children diagnosed with ASD to typically developing peers. These screening accuracy estimates are likely inflated due to exclusion of children with ambiguous presentations, such as those of HR non-ASD children who often present with social-emotional and other developmental differences (for a recent review of clinical characteristics of non-diagnosed siblings, see Pisula & Ziegart-Sadowska, 2015). When compared to the findings of Gardner et al. (2013), who employed an arguably more comparable sample (completed BITSEAs at age 1 and 2 years and screening accuracy computed using ASD risk classifications on the M-CHAT), our results are similar. Both samples (ours and those of Gardner et al.) had AUC within the .60 to .70 range and estimates of screening accuracy were below recommended values (Cicchetti et al., 1995; Council on Children with Disabilities, 2006; Dumont-Mathieu & Fein, 2005; Volkmar et al., 1988). Again, it is important to acknowledge that the siblings of children with ASD are at a heightened risk for many conditions (Ghirardi et al., 2018; Jokiranta-Olkonemi et al., 2016; Miller et al., 2019; Shivers et al., 2013; 2019) and the early manifestation of these may contribute to the BITSEA's inability to differentiate HR siblings with ASD from those who do not have ASD (but who may have other conditions not yet diagnosed).

There is evidence to support the use of the BITSEA with a high-risk population in a primary care setting even though our results do not support its use as a stand-alone screener for ASD. Asking caregivers to fill out a questionnaire about social-emotional development may reduce discomfort that can accompany questionnaires that query ASD specifically, especially for families who have not raised any concerns about ASD-related behavior in their younger children (Giserman Kiss & Carter, 2017; Zwaigenbaum et al., 2015). Furthermore, querying social-emotional differences in HR siblings provides clinically meaningful information by age one; up to 30% of caregivers have retrospectively recalled social-emotional developmental challenges (De Giacomo & Fombonne, 1998) and parents of HR siblings have prospectively endorsed more such challenges in children who are later diagnosed with ASD (Sacrey et al., 2015) during this period.

This study provides an in-depth analysis of the association between social-emotional problems and competencies and clinical outcomes with respect to ASD in a sample of HR siblings. This study benefits overall from a prospective longitudinal design, large HR toddler cohort with and without ASD, and comprehensive dataset encapsulating many clinical measures over time. This study is not without limitations, however. First, families of HR toddlers in this study have at least one child already diagnosed with ASD. As such, a better awareness of early developmental and behavioral signs associated with ASD may have influenced parent reporting on the BITSEA. Second, ongoing feedback parents receive regarding their child's development may influence parent report on the BITSEA and other measures. Third, additional variability may have been introduced due to BITSEA items having been pulled from the longer ITSEA (that is, responses to items that are not included as part of the BITSEA may have influenced how parents responded to items that were included in the BITSEA). Finally, prospectively studying our sample only to 36 months of age may have limited our ability to appreciate differences in social-emotional development trajectories among HR siblings. Despite these limitations, parents of

HR toddlers who were diagnosed with ASD at 36 months endorsed more social-emotional differences relative to parents whose children were not diagnosed with ASD. Because social-emotional challenges are associated in the long-term with poor academic performance (Campbell et al., 2006) and neurodevelopmental and mental health disorders (Campbell et al., 2000; Campbell et al., 2006; Fox, 2004; Shaw et al., 1994), collecting information on early social-emotional development is valuable for children at risk of any psychopathology, as early interventions have been shown to improve outcomes in preschool-aged children, with associated long-term consequences (Gross et al., 2003; Kelleher et al., 2006; Reid et al., 2004; Thomas & Zimmer-Gembeck, 2007).

Our results highlight the potential benefit of collecting information on multiple areas of development for HR siblings. Although the ROC analyses do not support use of the BITSEA as a stand-alone screener for ASD, it can be used as an adjunct measure to capture clinically meaningful information that may not be derived from an in-person clinical assessment. It is important to consider social-emotional development in the overall assessment and potentially intervention planning for children at risk of or diagnosed with ASD.

#### Funding Source

This study was supported by Women & Children's Health Research Institute (WCHRI), Brain Canada, Kids Brain Health Network (formerly NeuroDevNet), Canadian Institutes of Health Research (CIHR), Azrieli Foundation, Autism Speaks Canada, and Stollery Children's Hospital Foundation Chair in Autism.

#### References

- Akobeng, A. K. (2007). Understanding diagnostic tests 3: Receiver operating characteristic curves. *Acta Paediatrica*, 96, 644–647.
- Alink, L. R., Mesman, J., Van Zeijl, J., Stolk, M. N., Juffer, F., Koot, H. M., ... Van IJzendoorn, M. H. (2006). The early childhood aggression curve: Development of physical aggression in 10 to 50 month old children. *Child Development*, 77, 954–66.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5\*)*. Arlington, VA: American Psychiatric Pub.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B (Methodological)*, 57, 289–300.
- Boone, K. M., Brown, A. K., & Keim, S. A. (2018). Screening accuracy of the Brief Infant Toddler Social-Emotional Assessment to identify autism spectrum disorder in toddlers born at less than 30 weeks gestation. *Child Psychiatry and Human Development*, 49, 493–504. doi:10.1007/s10578-017-0768-2
- Brian, J., Bryson, S. E., Garon, N., Roberts, W., Smith, I. M., Szatmari, P., & Zwaigenbaum, L. (2008). Clinical assessment of autism in high-risk 18-month-olds. *Autism*, 12, 433–456. doi:10.1177/1362361308094500
- Briggs-Gowan, M. J., & Carter, A. S. (2000). *The Infant Toddler Social and Emotional Assessment (ITSEA)*. Pre-Publication Version. New Haven, CT: Yale University.
- Briggs-Gowan, M. J., & Carter, A. S. (2008). Social-emotional screening status in early childhood predicts elementary school outcomes. *Pediatrics*, 121, 957–962.
- Briggs-Gowan, M. J., Carter, A. S., Bosson-Heenan, J., Guyer, A. E., & Horwitz, S. M. (2006). Are infant-toddler social-emotional and behavioural problems transient? *Journal of the American Academy of Child and Adolescent Psychiatry*, 45, 849–858.
- Briggs-Gowan, M. J., Carter, A. S., Irwin, J. R., Wachtel, K., & Cicchetti, D. V. (2002). *Brief Infant-Toddler Social and Emotional Assessment (BITSEA) manual, version 2.0*. New Haven, CT: Yale University.

- Briggs-Gowan, M. J., Carter, A. S., McCarthy, K., Augustyn, M., Caronna, E., & Clark, R. (2013). Clinical validity of a brief measure of early childhood social-emotional/behavioural problems. *Journal of Pediatric Psychology*, 38, 577–587.
- Campbell, S. B., Shaw, D. S., & Gilliom, M. (2000). Early externalizing behavior problems: Toddlers and preschoolers at risk for later maladjustment. *Developmental Psychopathology*, 12, 467–488.
- Campbell, S. B., Spieker, S., Burchinal, M., & Poe, M. D. (2006). Trajectories of aggression from toddlerhood to age 9 predict academic and social functioning through age 12. *Journal of Child Psychology and Psychiatry*, 47, 791–800.
- Carter, A. S., Volkmar, F. R., Sparrow, S. S., Wang, J., Lord, C., Dawson, G., ... Schopler, E. (1998). The Vineland adaptive behaviour scales: Supplementary norms for individuals with autism. *Journal of Autism and Developmental Disorders*, 28, 287–302.
- Cicchetti, D. V., Volkmar, F., Klin, A., & Showalter, D. (1995). Diagnosing autism using ICD-10 criteria: A comparison of neural networks and standard multivariate procedures. *Child Neuropsychology*, 1, 26–37.
- Council on Children with Disabilities, Section on Developmental Behavioural Pediatrics, Bright Futures Steering Committee and Medical Home Initiatives for Children with Special Needs Project Advisory Committee. (2006). Identifying infants and young children with developmental disorders in the medical home: An algorithm for developmental surveillance and screening. *Pediatrics*, 118, 405–420. doi:10.1542/peds.2006-1231
- De Giacomo, A., & Fombonne, E. (1998). Parental recognition of developmental abnormalities in autism. *European Child & Adolescent Psychiatry*, 7(3), 131–136. doi:10.1007/s007870050058
- Dumont-Mathieu, T., & Fein, D. (2005). Screening for autism in young children: The modified checklist for autism in toddlers (M-CHAT) and other measures. *Mental Retardation and Developmental Disabilities Research Reviews*, 11, 253–262.
- Fischer, J. E., Bachmann, L. M., & Jaeschke, R. (2003). A readers' guide to the interpretation of diagnostic test properties: Clinical example of sepsis. *Intensive Care Medicine*, 29, 1043–1051.
- Fox, N. A. (2004). Temperament and early experience form social behaviour. *Annals of the New York Academy of Science*, 1038, 171–178.
- Gardner, L. M., Murphy, L., Campbell, J. M., Tylavsky, F., Palmer, F. B., & Graff, J. C. (2013). Screening accuracy for risk of autism spectrum disorder using the brief infant-toddler social and emotional assessment (BITSEA). *Research in Autism Spectrum Disorders*, 7, 591–600.
- Ghirardi, L., Brikell, I., Kuja-Halkola, R., Freitag, C. M., Franke, B., Asherson, P., ... Larsson, H. (2018). The familial co-aggregation of ASD and ADHD: a register-based cohort study. *Molecular Psychiatry*, 23, 257–262. doi:10.1038/mp.2017.17
- Giserman Kiss, I., & Carter, A. S. (2017). Brief infant-toddler social and emotional assessment (BITSEA). In *Encyclopedia of autism spectrum disorders* (pp. 1–5). New York, NY: Springer.
- Giserman Kiss, I., Feldman, M. S., Sheldrick, R. C., & Carter, A. S. (2017). Developing autism screening criteria for the brief infant toddler social emotional assessment. *Journal of Autism and Developmental Disorders*, 47, 1269–1277. doi:10.1007/s10803-017-3044-1.
- Glascoc, F. P. (2003). Parents' evaluation of developmental status: How well do parents' concerns identify children with behavioral and emotional problems? *Clinical Pediatrics*, 42(2), 133–138. doi:10.1177/000992280304200206
- Gotham, K., Pickles, A., & Lord, C. (2009). Standardizing ADOS scores for a measure of severity in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 39, 693–705.
- Gotham, K., Risi, S., Pickles, A., & Lord, C. (2007). The autism diagnostic observation schedule: Revised algorithms for improved diagnostic validity. *Journal of Autism and Developmental Disorders*, 37, 613.
- Gross, D., Fogg, L., Webster-Stratton, C., Garvey, C., Julion, W., & Grady, J. (2003). Parent training of toddlers in day care in low-income urban communities. *Journal of Consulting and Clinical Psychology*, 71, 261–278.
- Jokiranta-Olkoniemi, E., Cheslack-Postava, K., Sucksdorff, D., Suominen, A., Gyllenberg, D., Chudal, R., ... Sourander, A. (2016). Risk of psychiatric and neurodevelopmental disorders among siblings of probands with autism spectrum disorder. *JAMA Psychiatry*, 73, 622–629. doi:10.1001/jamapsychiatry.2016.0495
- Karabekiroglu, K., Briggs-Gowan, M. J., Carter, A. S., Rodopman-Arman, A., & Akbas, S. (2010). The clinical validity and reliability of the brief infant-toddler social and emotional assessment (BITSEA). *Infant Behavior and Development*, 33, 503–509. doi:10.1016/j.infbeh.2010.07.001
- Kelleher, K. J., Campo, J. V., & Gardner, W. P. (2006). Management of pediatric mental disorders in primary care: Where are we now and where are we going? *Current Opinions in Pediatrics*, 18, 649–653.
- Kruizinga, I., Visser, J. C., van Batenburg-Eddes, T., Carter, A. S., Jansen, W., & Raat, H. (2014). Screening for autism spectrum disorders with the brief infant-toddler social and emotional assessment. *PLoS One*, 9, e97630.
- Lasch, C., Wolff, J. J., & Elison, J. T. (2019). Examining criterion-oriented validity of the Repetitive Behavior Scales for Early Childhood (RBS-EC) and the Video-Referenced Rating of Reciprocal Social Behavior (vrRSB). *Development and Psychopathology*, 28, 1–11. doi:10.1017/S0954579419001159.
- Lord, C., Risi, S., Lambrecht, L., Cook, E. H., Leventhal, B. L., DiLavore, P. C., ... Rutter, M. (2000). The autism diagnostic observation Schedule—Generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30, 205–223.
- Lord, C., Rutter, M. L., Goode, S., Heemsbergen, J., Jordan, H., Mawhood, L., & Schopler, E. (1989). Autism diagnostic observation schedule: A standardized observation of communicative and social behaviour. *Journal of Autism and Developmental Disorders*, 19, 185–212.
- Lord, C., Rutter, M., & Le Couteur, A. (1994). Autism diagnostic interview-revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 24, 659–685.
- Marrus, N., Glowinski, A. L., Jacob, T., Klin, A., Jones, W., Drain, C. E., ... Sant, S. M. (2015). Rapid video-references ratings of reciprocal social behavior in toddlers: A twin study. *The Journal of Child Psychology and Psychiatry*, 56, 1338–1346. doi:10.1111/jcpp.12391
- Marrus, N., Grant, J. D., Harris-Olenak, B., Albright, J., Bolster, D., Haber, J. R., et al. (2020). Genetic architecture of reciprocal social behavior in toddlers: Implications for heterogeneity in the early origins of autism spectrum disorder. *Development and Psychopathology*, in press.
- McDonald, N. M., Senturk, D., Scheffler, A., Brian, J. A., Carver, L. J., Charman, T., ... Klin, A. (2019). Developmental trajectories of infants with multiplex family risk for autism: A baby siblings research consortium study. *JAMA Neurology*, 77(1), 73–81. doi:10.1001/jamaneuro.2019.3341.
- Miller, M., Musser, E. D., Young, G. S., Olson, B., Steiner, R. D., & Nigg, J. T. (2019). Sibling recurrence risk and cross-aggregation of attention-deficit/hyperactivity disorder and autism spectrum disorder. *JAMA Pediatrics*, 173, 147–152. doi:10.1001/jamapediatrics.2018.4076
- Mullen, E. M. (1995). *Mullen scales of early learning*. MN: AGS Circle Pines.
- Pisula, E., & Ziegart-Sadowska, K. (2015). Broader autism phenotype in siblings of children with ASD: A review. *Molecular Autism*, 16, 13217–13258. doi:10.3399/ijms10613217
- Pontoppidan, M., Niss, N. K., Pejtersen, J. H., Julian, M. M., & Væver, M. S. (2017). Parent report measures of infant and toddler social-emotional development: A systematic review. *Family Practice*, 34(2), 127–137. doi:10.1093/fampra/cmx003
- Raza, S., Sacrey, L. R., Zwaigenbaum, L., Bryson, S., Brian, J., Smith, I. M., ... Garon, N. (2019). Relationship between early social-emotional behavior and autism spectrum disorder: A high-risk sibling study. *Journal of Autism and Developmental Disorders*, in press. doi:10.1007/s10803-019-03977-3
- Reid, M. J., Webster-Stratton, C., & Baydar, N. (2004). Halting the development of conduct problems in Head Start children: the effects of parent training. *Journal of Clinical Child and Adolescent Psychology*, 33, 279–291.
- Robins, D. L., Fein, D., Barton, M. L., & Green, J. A. (2001). The Modified checklist for autism in toddlers: An initial study investigating the early detection of autism and pervasive developmental disorders. *J Autism Dev Disord*, 31, 131–144. doi:10.1023/A:1010738829569
- Sacrey, L. R., Bryson, S., Zwaigenbaum, L., Brian, J., Smith, I. M., Roberts, W., ... Garon, N. (2018). The autism parent screen for infants: Predicting risk of autism spectrum disorder based on parent-reported behaviour observed at 6–24 months of age. *Autism*, 22, 322–334.
- Sacrey, L. R., Zwaigenbaum, L. R., Bryson, S. E., Brian, J., Smith, I. M., Roberts, W., ... Armstrong, V. (2015). Can parents' concerns predict autism spectrum disorder? A prospective study of high-risk siblings from 6 to 36 months of age.

- Journal of the American Academy of Child, 54*(6), 470–478. doi:10.1016/j.jaac.2015.03.014
- Schmidt, M., Demulder, E., & Denham, S. (2010). Kindergarten social-emotional competence: Developmental predictors and psychosocial implications. *Early Child Development and Care, 172*, 451–462.
- Shaw, D. S., Keenan, K., & Vondra, J. I. (1994). Developmental precursors of externalizing behaviour: Ages 1 to 3. *Developmental Psychology, 30*, 355–364.
- Shields, AM, Cicchetti, D, & Ryan, RM. (1994). The development of emotional and behavioral self-regulation and social competence among maltreated school-age children. *Development and Psychopathology, 6*(1), 57–75. doi:10.1017/S0954579400005885
- Shivers, C. M., Deisenroth, L. K., & Taylor, J. L. (2013). Patterns and predictors of anxiety among siblings of children with autism spectrum disorders. *Journal of Autism and Developmental Disorders, 43*, 1336–1346. doi:10.1007/s10803-012-1685-7
- Shivers, C. M., Jackson, J. B., & McGregor, C. M. (2019). Functioning among typically developing siblings of individuals with autism spectrum disorder: A meta-analysis. *Clinical and Child Family Psychology Review, 22*, 172–196. doi:10.1007/s10567-018-0269-2
- Sparrow, S. S., Balla, D. A., Cicchetti, D. V., Harrison, P. L., & Doll, E. A. (1984). *Vineland Adaptive Behaviour Scales*. Minnesota: Circle Pines.
- Stadnick, N., Chlebowski, C., Baker-Ericzen, M., Dyson, M., Garland, A., & Brookman-Frazer, L. (2017). Psychiatric comorbidity in autism spectrum disorder: Correspondence between mental health clinician report and structured parent interview. *Autism, 21*, 841–851.
- Thomas, R., & Zimmer-Gembeck, M. J. (2007). Behavioural outcomes of parent-child interaction therapy and Triple P-positive parenting program: A review and meta-analysis. *Journal of Abnormal Child Psychology, 35*, 475–495.
- van Zeijl, J., Mesman, J., Stolk, M. N., Alink, L. R., Van IJzendoorn, M. H., Bakermans-Kranenburg, M. J., ... Koot, H. M. (2006). Terrible ones? Assessment of externalizing behaviours in infancy with the child behaviour checklist. *Journal of Child Psychology Psychiatry and Allied Disciplines, 47*, 801–10.
- Volkmar, F. R., Carter, A., Sparrow, S. S., & Cicchetti, D. V. (1993). Quantifying social development in autism. *Journal of the American Academy of Child & Adolescent Psychiatry, 32*, 627–632.
- Volkmar, F. R., Cicchetti, D. V., Dykens, E., Sparrow, S. S., Lechman, J. F., & Cohen, D. J. (1988). An evaluation of the autism behaviour checklist. *Journal of Autism and Developmental Disorders, 18*, 81–97.
- Wakschlag, L. S., & Danis, B. (2009). Characterizing early childhood disruptive behaviour. In C. H. Zeanah (Ed.), *Handbook of infant mental health* (3rd ed). New York, NY: Guilford Press.
- Zwaigenbaum, L., Bauman, M. L., Fein, D., Pierce, K., Buie, T., Davis, P. A., ... Choueiri, R. (2015). Early screening of autism spectrum disorder: Recommendations for practice and research. *Pediatrics, 136*, S59.