Paleobiology, 49(2), 2023, pp. 374–375 DOI: 10.1017/pab.2023.15



CORRIGENDUM

Linking host plants to damage types in the fossil record of insect herbivory – CORRIGENDUM

Sandra R. Schachat, Jonathan L. Payne, and C. Kevin Boyce

https://doi.org/10.1017/pab.2022.35, published online by Cambridge University Press, 9 January 2023.

This article published with an error in the legend of Figure 6 (Schachat et al. 2023). In

the bottom of the three panels, the labels "1,000" and "750" are swapped from their proper positions. The correct figure appears below.

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Paleontological Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (http://creativecommons.org/licenses/by-nc-nd/4.0), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article. 0094-8373/23



Corrigendum 375




FIGURE 6. Mean values and 95% confidence intervals for beta-diversity metrics generated by resampling and subsampling data for the two most abundant host plants from (A) Willershausen, (B) Colwell Creek Pond, and (C) Williamson Drive. At the highest sample sizes, represented in dark blue, the data were resampled rather than subsampled. *Macronrytrs. scheuchzeri = Macroneuropteris scheuchzeri; Sgllrphyllm. = Sigillariophyllum.*

The authors apologize for this error.

Literature Cited

Schachat, S. R., J. L. Payne, and C. K. Boyce. 2023. Linking host plants to damage types in the fossil record of insect herbivory. Paleobiology 49:232–258.