


CORRIGENDUM

Discovery of the skull of Stephanorhinus kirchbergensis (Jäger, 1839) above the Arctic Circle – CORRIGENDUM

Irina V. Kirillova, Olga F. Chernova, Jan van der Made, Vladimir V. Kukarskikh, Beth Shapiro, Johannes van der Plicht, Fedor K. Shidlovskiy, Peter D. Heintzman, Thijs van Kolfschoten, Oksana G. Zanina

doi: <https://doi.org/10.1017/qua.2017.53>, Published by Cambridge University Press, 30 August 2017.

In Kirillova *et al.* (2017), Figure 4 appeared with incorrect values along the x- and y-axes. Below is the correct figure:



REFERENCES

- Kirillova, I.V., Chernova, O.F., van der Made, J., Kukarskikh, V.V., Shapiro, B., van der Plicht, J., Shidlovskiy, F.K., Heintzman, P.D., van Kolfschoten, T., Zanina, O.G., 2017. Discovery of the skull of Stephanorhinus kirchbergensis (Jäger, 1839) above the Arctic Circle. *Quaternary Research* <https://doi.org/10.1017/qua.2017.53>.

Figure 4. (color online) The widths of the anterior (DTa) and posterior (DTp) lobes of the second upper molar (M^2) of the Chondon rhinoceros compared with those of other species of *Stephanorhinus*: *S. kirchbergensis* from type locality Kirchberg, Mosbach, Neumark Nord, Ehringsdorf, Bilzingsleben, Rhinegravels; *S. hemitoechus* from Murr, Neumark Nord, Rhinegravels; *S. hundsheimensis* from Voigtsdorf, Süssenborn, Mosbach, Soleihac, Mauer; and *S. etruscus* and *S. aff. etruscus* from the Upper Valdarno and Atapuerca TD4. All teeth from tooth rows (so their identity as M^2 is clear), save for one of the type specimens of *S. kirchbergensis* from Kirchberg (marked as “K”). For comparisons of further cheek teeth and provenance of the data, see the Supplementary Materials.