

# YSOs in Taurus-Auriga-Perseus and Orion

Sarolta Zahorecz<sup>1</sup>, L. Viktor Tóth<sup>1</sup>, Gábor Marton<sup>2</sup>,  
Toshikazu Onishi<sup>3</sup>, Lajos G. Balázs<sup>2</sup>, Orsolya Fehér<sup>1</sup>,  
Akiko Kawamura<sup>4</sup>, Yoshimi Kitamura<sup>5</sup>, Mónika Lisztes<sup>1</sup>,  
Atsushi Nishimura<sup>3</sup>, László Pásztor<sup>6</sup>, Sándor Pintér<sup>1</sup>, István Rácz<sup>1</sup>,  
Motohide Tamura<sup>4</sup>, Rogel M.D. Sese<sup>7</sup> and Munetaka Ueno<sup>5</sup>

<sup>1</sup>Loránd Eötvös University, Department of Astronomy, Pázmány Péter sétány 1, 1117  
Budapest, Hungary, email: S.Zahorecz@astro.elte.hu

<sup>2</sup>Konkoly Observatory of the Hungarian Ac. of Sci., PO Box 67, 1525 Budapest, Hungary

<sup>3</sup>Department of Physical Science, Osaka Prefecture University, Gakuen 1-1, Sakai, Osaka  
599-8531, Japan

<sup>4</sup>National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

<sup>5</sup>JAXA, 3-1-1 Yoshinodai, Sagamihara, Kanagawa, 229-8510, Japan

<sup>6</sup>Research Institute for Soil Science and Agricultural Chemistry of the Hungarian Academy of  
Sciences, GIS Lab, Budapest, Hungary

<sup>7</sup>College of Arts and Sciences, UPLB College, Laguna 4031 Philippines

**Abstract.** Physical parameters were derived for 100 young stellar objects in the TAPO region.

**Keywords.** stars: formation — stars: pre-main-sequence — infrared: stars

---

We made an analysis of 4441 AKARI FIS (Kawada *et al.* 2007; Yamamura *et al.* 2010) point sources in the Tau-Aur-Per and Orion region. Spectral energy distribution (SED) was drawn for about 550 point sources based on AKARI FIS, AKARI IRC (Ishihara *et al.* 2010), Spitzer Space Telescope (SST, Werner *et al.* 2004) Infrared Array Camera (IRAC; Fazio *et al.* 2004), Mid-Infrared Photometer for Spitzer (MIPS; Rieke *et al.* 2004), 2 Micron All Sky Survey (Skrutskie *et al.* 2006) and Wide-Field Infrared Survey Explorer (WISE, Wright *et al.* 2010) and various other photometric data. We determined the physical parameters (e.g. stellar mass, temperature and radius, disk size and mass) of 100 YSOs with the SED Fitting Tool of Robitaille *et al.* 2007 and for other 450 sources we determined an evolutionary stage based on the slope of the SED in mid-IR and FIR.

**Acknowledgement.** The European Union and the European Social Fund have provided financial support to the project under the grant agreement no. TAMOP-4.2.1/B-09/1/KMR-2010-0003. This research was partly supported by the Hungarian Research Fund (OTKA) and the HAS-JSPS mobility program.

## References

- Ishihara, D., Onaka, T., Kataza, H. *et al.* 2010, *A&A*, 514, 1  
Fazio, G. G., Hora, J. L., Allen, L. E. *et al.* 2004, *ApJS*, 154, 10  
Rieke, G. H., Young, E. T., Engelbracht, C. W. *et al.* 2004, *ApJS*, 154, 25  
Kawada, M., Baba, H., Barthel, P. D., Clements, D. *et al.* 2007, *PASJ*, 59, 389  
Robitaille, T. P., Whitney, B. A., Indebetouw, R., & Wood, K. 2007, *ApJS*, 169, 328  
Skrutskie, M. F., Cutri, R. M., Stiening, R. *et al.* 2006, *AJ*, 131, 1163  
Yamamura, I. *et al.* 2010, *AKARI/FIS Bright Source Catalogue Version 1.0 Release Note*  
Werner, M. W., Roellig, T. L., Low, F. J. *et al.* 2004, *ApJS*, 154, 1  
Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K. *et al.* 2010, *AJ*, 140, 1868