

# CLUSTERING OF RED GALAXIES NEAR A RADIO-LOUD QUASAR AT $z = 1.086$

I. TANAKA AND T. YAMADA

*Astronomical Institute, Tohoku University,  
Aoba-ku, Sendai 980-77, Japan*

A. ARAGÓN-SALAMANCA AND T. KODAMA

*Institute of Astronomy,  
Madingley Road, Cambridge, CB3 0HA, UK*

K. OHTA

*Department of Astronomy, Kyoto University,  
Kyoto, 606-01, Japan*

AND

N. ARIMOTO

*Institute of Astronomy, University of Tokyo,  
Mitaka, Tokyo 181, Japan*

We investigated the environment of the radio-loud quasar 1335.8+2834 at  $z = 1.086$  where an excess surface number density of galaxies was reported by Huthings et al. (1993). By obtaining near-infrared and new deep optical images of the field, we found a clustering of objects with very red optical-NIR color,  $4 < R - K < 6$  and  $3 < I - K < 5$  near the quasar. The colors and magnitude of the reddest objects are consistent with those predicted for old (2–4 Gyr) passively evolving elliptical galaxies at  $z = 1.1$ .

There is a large fraction ( $> 60\%$  after field correction) of objects with moderately red colors,  $4 < R - K < 4.5$ , which have a similar sky distribution as the reddest objects. They are interpreted as cluster galaxies with on-going star formation. For more details, see Yamada et al. (1997) and Tanaka et al. (1997, in preparation).

## References

- Hutchings, J. B., Crampton, D., & Persram, D. 1993, *AJ*, **106**, 1324  
Yamada, T., Tanaka I., N., Aragón-Salamanca, A., Kodama, T., Ohta, K., & Arimoto, N. 1992, *ApJ*, **487**, L125