

IAU Symposium

376

April 17th–21st 2023

Budapest, Hungary

Proceedings of the International Astronomical Union

At the crossroads of astrophysics and cosmology: Period–luminosity relations in the 2020s

Edited by


Richard de Grijs
Patricia A. Whitelock
Márcio Catelan

ISSN 1743-9213

International Astronomical Union



CAMBRIDGE
UNIVERSITY PRESS



AT THE CROSSROADS OF ASTROPHYSICS AND COSMOLOGY:
PERIOD–LUMINOSITY RELATIONS IN THE 2020S

IAU SYMPOSIUM 376

Uncrowded Cepheids observed with the *James Webb Space Telescope (JWST)* in the near-infrared. High-resolution *JWST* observations can test confusion-limited *Hubble Space telescope (HST)* observations that are used to measure the Hubble constant. *JWST* provides superior source separation to negate crowding noise resulting in a factor of > 2.5 reduction in the dispersion of the Cepheid period–luminosity relation in galaxies that host Type Ia supernovae (image shown for NGC 5584) and the geometric distance reference, NGC 4258, with the period–luminosity relation shown here. The agreement between Cepheid distance measures between *HST* and *JWST* is excellent and provides the strongest evidence to date that systematic errors in *HST* Cepheid measurements do not play a significant role in the present ‘Hubble tension,’ now at 5σ confidence. The source of the Hubble tension remains an outstanding mystery of potentially cosmological origin and warrants further study. (Credit: Adam Riess, Johns Hopkins University, USA)

IAU SYMPOSIUM PROCEEDINGS SERIES

Chief Editor

JOSÉ MIGUEL RODRIGUEZ ESPINOSA, General Secretariat

Instituto de Astrofísica de Andalucía

Glorieta de la Astronomía s/n

18008 Granada

Spain

IAU-general.secretary@iap.fr

Editor

DIANA WORRALL, Assistant General Secretary

HH Wills Physics Laboratory

University of Bristol

Tyndall Avenue

Bristol


BS8 1TL

UK

IAU-assistant.general.secretary@iap.fr

INTERNATIONAL ASTRONOMICAL UNION
UNION ASTRONOMIQUE INTERNATIONALE

International Astronomical Union



**AT THE CROSSROADS OF
ASTROPHYSICS AND
COSMOLOGY:
PERIOD–LUMINOSITY
RELATIONS IN THE 2020S**

**PROCEEDINGS OF THE 376th SYMPOSIUM OF
THE INTERNATIONAL ASTRONOMICAL UNION
BUDAPEST, HUNGARY**

17–21 April, 2023

Edited by

RICHARD DE GRIJS

Macquarie University, Sydney, Australia

PATRICIA A. WHITELOCK

South African Astronomical Observatory and University of Cape Town, South Africa

and

MÁRCIO CATELAN

Pontificia Universidad Católica de Chile, Santiago, Chile



**CAMBRIDGE
UNIVERSITY PRESS**

CAMBRIDGE UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom
1 Liberty Plaza, Floor 20, New York, NY 10006, USA
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© International Astronomical Union 2024

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of the International Astronomical Union.

First published 2024

Printed in Great Britain by Henry Ling Limited, The Dorset Press, Dorchester, DT1 1HQ

Typeset in System L^AT_EX 2 ϵ

*A catalogue record for this book is available from the British Library of Congress
Cataloguing in Publication data*

This journal issue has been printed on FSCTM-certified paper and cover board. FSC is an
independent, non-governmental, not-for-profit organization established to promote the
responsible management of the world's forests. Please see www.fsc.org for information.

ISBN 9781009353045 hardback
ISSN 1743-9213

Table of Contents

Preface	viii
Editors	xii
List of Participants	xiii
The Cepheid Extragalactic Distance Scale: Past, Present and Future <i>Wendy L. Freedman and Barry F. Madore</i>	1
The Local Value of H_0 <i>Adam G. Riess and Louise Breuval</i>	15
The Tip of the Red Giant Branch as a Cosmological Tool <i>Myung Gyoon Lee</i>	30
Period–Luminosity Relations in the Local Group of Galaxies <i>Igor Soszyński</i>	48
Theoretical Stellar Pulsation Physics <i>M. Marconi, G. De Somma, R. Molinaro and I. Musella</i>	68
Anomalous Cepheids: Updated Theoretical Period–Luminosity–Color and Period–Wesenheit Relations <i>Giulia De Somma, Marcella Marconi, Santi Cassisi and Roberto Molinaro</i>	84
Pulsation of chemically peculiar stars <i>Ernst Paunzen</i>	91
A study of the stellar photosphere–hydrogen ionization front interaction in δ Scuti stars <i>Mami Deka, Shashi M. Kanbur, Sukanta Deb and Susmita Das</i>	98
A multi-wavelength analysis of BL Her stars: Models versus Observations <i>S. Das, L. Molnár, S. M. Kanbur, M. Joyce, A. Bhardwaj, H. P. Singh, M. Marconi, V. Ripepi and R. Smolec</i>	105
Impact of the <i>Gaia</i> ESA mission on the primary Period–Luminosity Calibrators in the Milky Way: Cepheids and RR Lyrae <i>Gisella Clementini</i>	115
Primary Period–Luminosity-Relation Calibrators in the Milky Way: Cepheids and RR Lyrae Physical basis, Calibration, and Applications <i>M. A. T. Groenewegen</i>	128
Cepheids with giant companions: A new, abundant source of Cepheid astrophysics <i>Bogumił Pilecki</i>	150

Implication of the period-magnitude relation for massive AGB stars and its astronomical applications	159
<i>Akiharu Nakagawa, Tomoharu Kurayama, Hiroshi Sudou and Gabor Orosz</i>	
The structure of the Milky Way from period-luminosity relations	180
<i>Dorota M. Skowron</i>	
Granulation in Red Supergiants: The Scaling Relations	195
<i>Yi Ren, Zehao Zhang, Biwei Jiang, Igor Soszyński and Tharindu Jayasinghe</i>	
The distance scale of Type II Cepheids from near-infrared observations in the Magellanic Clouds	205
<i>T. Sicignano, V. Ripepi, R. Molinaro, A. Bhardwaj, M. Marconi and M.-R. L. Cioni</i>	
Sub-percent binary star masses and distances from interferometric observations	214
<i>Alexandre Gallenne</i>	
Globular cluster metallicities and distances from disentangling their RR Lyrae light curves	222
<i>A. Arellano Ferro</i>	
The PL diagram for δ Sct stars: back in business as distance estimators	239
<i>Antonio García Hernández, Javier Pascual-Granado, Mariel Lares-Martiz, Giovanni M. Mirouh, Juan Carlos Suárez, Sebastián Barceló Forteza and Andrés Moya</i>	
Period–Luminosity–Metallicity relations for Classical Pulsators at Near-infrared Wavelengths	250
<i>Anupam Bhardwaj</i>	
A multiphase study of classical Cepheids in the Magellanic Clouds- Models and Observations	267
<i>Kerdaris Kurbah, Shashi M. Kanbur, Sukanta Deb, Susmita Das, Mami Deka, Anupam Bhardwaj, Hugh Riley Randal and Selim Kalici</i>	
Double-mode RR Lyrae star — robust distance and metallicity indicators	275
<i>Shu Wang, Xiaodian Chen, Jianxing Zhang and Licai Deng</i>	
Empirical constraints for the instability strip from the analysis of LMC Cepheids	281
<i>F. Espinoza-Arancibia and B. Pilecki</i>	
The period-luminosity relations of red supergiants	292
<i>Biwei Jiang, Yi Ren and Ming Yang</i>	
Long-Period Variables as distance and age indicators in the era of <i>Gaia</i> and LSST	306
<i>Michele Trabucchi</i>	
China Space Station Telescope and Variable Star Studies	319
<i>Xiaodian Chen, Shu Wang, Licai Deng, Richard de Grijs, Xiaoyue Zhou, Xiaohan Chen and Jianxing Zhang</i>	

OH/IR stars and the Period-Luminosity Relation of Mira variables	328
<i>D. Engels, S. Etoka, F. Jiménez-Esteban, W. Herrmann, B. López-Martí</i>	
Author Index	335

Preface

This volume includes contributions from IAU Symposium 376, *At the crossroads of astrophysics and cosmology: Period–luminosity relations in the 2020s*.

In the early 1900s, Henrietta Leavitt, in 1908—and later Adams & Joy and Shapley & Walton, both in 1927—showed that bright Cepheid variables were characterised by a narrow range in spectral type or, equivalently, temperature at a given period. This, in turn, led to the establishment of a tight period–luminosity relation (PLR): more luminous stars are expected to have longer pulsation periods. The spectral-type/temperature restriction of Cepheids naturally led to the realisation that pulsating variables only occur in a narrow “instability strip” in the Hertzsprung–Russell diagram.

This classical instability strip is host to a range of periodic variable stars, including classical Cepheids (δ Cephei stars), Mira variables at the top of the asymptotic giant branch (AGB), Type II Cepheids, and RR Lyrae stars. SX Phoenicis and δ Scuti variables (jointly known as dwarf Cepheids or ultrashort-period variables) as well as anomalous Cepheids, and ZZ Ceti, V777 Herculis and GW Virginis pulsating white dwarfs are also found in specific instability strips. Many of these stellar types obey specific period–mean density relations. Curiously, not only pulsating stars exhibit PLRs: well-defined and long-established relationships between orbital periods and luminosities also exist for contact binaries of W Ursae Majoris (EW) type. Crucial remaining open issues in this broad field include the metallicity dependence of the zero point of the Cepheid PLR, the possible presence of a break at a pulsation period around 10 days, and the effects of binarity and circumstellar envelopes. Intrinsic PLR widths may offer unique insights into the physical processes shaping these relations and the underlying physical properties of the contributing stars (stellar structure, atmospheric parameters, and pulsation properties).

A promising approach to reducing the current systematic uncertainties associated with the present-day expansion rate of the Universe—the well-known tension in the Hubble parameter—is by trying to achieve improved local calibrations of primary distance indicators and their derivatives, including calibration of the photometric zero point of the Cepheid PLR, at both optical and—potentially with much reduced scatter—infrared wavelengths (e.g., starting from existing *Spitzer Space Telescope* data), for instance through trigonometric-parallax measurements of carefully selected Cepheid samples by the European Space Agency’s *Gaia* satellite. Improvements in the Cepheid and Mira distance scales will be achievable to a level of 3–4% or better, based on forthcoming mid-infrared observations with the *James Webb Space Telescope (JWST)*. Perhaps the most powerful use of the variety of PLRs present for variable stars is the ability to cross-check results from any individual technique. The diverse stellar types exhibiting PLRs are drawn from distinct stellar populations, with their own age and metallicity distributions. Working together, these techniques can explore population-based systematics in distance determination to provide insights into potential calibration biases in the current Type Ia supernovae calibration.

After much discussion, we decided to frame IAU Symposium 376 by initially focusing on the discrepancy in the Hubble constant based on different approaches, including the “standard” Cepheid PLR. This choice led to many productive and engaging discussions, triggered by numerous inspiring talks that highlighted the latest developments. In essence, this theme followed on from the Spring school on the distance ladder that had been organised at Konkoly Observatory during the week prior to the conference week.

From the outset, it became clear that the *JWST* will soon become a game changer given the amazing data already generated by this new observatory. Thanks to its flawless launch and smooth initial operations, the community is now looking at an observatory

that may last much longer than its nominal five-year operational lifetime. In turn, this offers an enormously expanded scope for studies of the distance ladder. Even at this early time, we are seeing significant improvements in spatial resolution with respect to that offered by the *Hubble Space Telescope* at similar (near-)infrared wavelengths. In fact, preliminary results imply that the scatter in the Cepheid PLR at 1.5 μm may be reduced by a factor of 2–3. Multiple participants presented very exciting results based on early *JWST* data.

Another eye-catching aspect brought up in many inspirational contributions is the phenomenal distance accuracy currently achievable from Cepheid PLRs. *Gaia* cluster Cepheids now show Large Magellanic Cloud-like scatter in their period–Wesenheit relations, while the distance to Messier 33 has now been established to better than 1.3%. This finally allows us to understand secondary effects, such as the impact of period fluctuations (e.g., in Messier 51) on distance estimates.

Similarly, tip of the red giant branch (TRGB) distance determination has come a long way over the past three or more decades and promises to deliver a powerful alternative approach to the standard PLR technique for Cepheids, pushing to ever greater distances. In addition, newly developed stellar population methods now allow us to apply the “J-AGB” method, exploiting the characteristics of carbon-rich stars.

Meanwhile, comparisons of distance estimates using different, independent methods are finally coming together. By pursuing minimisation of systematics and secondary variability effects, over the past decade the distance ladder has narrowed from a consideration of many complementary, independent techniques to a streamlined ladder with well-defined rungs, from Geometry to Cepheids, from Cepheids to SNe Ia, and from SNe Ia to redshifts. However, the field is now returning to the broader implications resulting from this much-improved ladder, with a greater focus on cross-checks with other methods. *Gaia*, *JWST*, and soon the Vera C. Rubin Observatory’s Legacy Survey of Space and Time will be game changers out to nearby galaxies at distances as far as tens of megaparsecs.

Despite an early focus on the challenges posed by the extragalactic distance scale, a significant fraction of the conference was dedicated to the underlying physics governing stellar pulsation and variability, for different types of tracers. Among the classical Cepheids, renewed interest focuses on multimode variability and period fluctuations, from short-period overtones to ultralong-period variability. Current developments promise exciting opportunities to much better determine the stellar mass–luminosity relation, often based on painstaking, detailed work where every object counts. Perhaps even more importantly, we are now reaching observational regimes where we can finally get a good handle on metallicity effects in the context of Cepheid variability.

AGB stars are coming back into fashion, particularly because of improved calibration methods and new theoretical developments. Parallaxes from masers obtained with very long baseline interferometry (VLBI), at nearby distances anchored by *Gaia* parallaxes, are providing very useful constraints. These are important developments in the era of the new extremely large telescopes which are more infrared-sensitive than their smaller counterparts. Moreover, the Square Kilometre Array, the next-generation Very Large Array (ngVLA), the Event Horizon Telescope, and other major radio initiatives open up new areas of research involving AGB and semi-regular variables as useful tracers of the physical processes taking place in and distances to numerous nearby galaxies.

RR Lyrae stars remain the workhorse PLR tracers for older stellar populations given their pre-eminent importance for near-field cosmology and for understanding the formation history of the Galaxy, particularly when combined with dynamical data. *Gaia* has also triggered a revival in this field, complemented by many cutting-edge ground-based projects. A major focus at the present time is on the metallicity dependence of the RR Lyrae PLRs, so that their empirical basis now supports theoretical arguments.

The wealth of data available at the present time and their high quality now allow us to consider secondary effects that may play a role in this field, such as the light-curve shape, multi-mode pulsation properties, period changes, binarity, circumstellar envelopes, etc.

This conference has clearly shown that stellar pulsation physics has reached a high level of maturity, so that our attention is increasingly focused on more diverse pulsator types other than classical Cepheids and RR Lyrae, including anomalous and Type II Cepheids, δ Scuti stars, magnetically and chemically peculiar stars, scaling relations and PLRs pertaining to supergiant pulsators, contact binary systems, and many others.

It has become clear that the importance of *Gaia* for studies of variable stars cannot be overstated. It has already collected more than a trillion CCD measurements down to $G \sim 21$ mag with uncertainties as small as 1 mmag. At the same time, the mission provides colours, astrometry, and spectroscopic time-series observations (for future release), thus making it the perfect tool for variability studies. At the present time, we already have access to *Gaia* observations of 271,000 RR Lyrae stars among some 10 million variables in Data Release 3 (DR3), with DR4 potentially reaching up to 100 million variable stars.

The key outstanding issue in this field, pertaining to PLRs in general, relates to the systematics affecting the current crop of *Gaia* parallaxes. We still need better ways to mitigate those effects, particularly as regards those objects that exhibit ‘astrophysical confusion’ in the sense of outflows, disks, etc. At present, it appears that the ‘counter-corrections’ required to derive accurate distances are often similar to the intrinsic, recommended corrections, so that more work is required to overcome those limitations. Many methods have been proposed (cluster membership, asteroseismology, binary systems with interferometric orbits, Miras with VLBI parallaxes, etc.), so it appears that the field is slowly moving to an acceptable solution in this area as well. The picture that is emerging is that there is not a single ‘best’ parallax correction that may or may not be magnitude-dependent, but we will have to resolve this issue using a sample-based approach.

Numerous contributions discussed the structure of the Milky Way, showing the significant synergies among different variability tracers, although one should keep in mind that different ages may trace different spatial components. Combined with dynamical measures, we now have a pretty good handle on the orbits of the Magellanic Clouds and their wake, on the warped and flaring young disk structure of the Milky Way, and even on the structure of our Galaxy on the other side of the Galactic Centre.

Theoretical advances are, of course, equally important as all those beautiful observational results. At this time, it seems that we are overwhelmed with high-quality data sets and theory is taking a bit of a back seat. Nevertheless, numerous teams are working on improving our theoretical understanding of the underlying pulsation physics, offering ever more detailed tests of models aided by the unmatched quality of observational data, including such models that focus on the boundaries of the classical instability strip. Major advances are seen in the context of understanding and modelling the effects of rotation and convection, where we are slowly moving from 1D to more realistic 3D models.

In view of all of these exciting developments, the future of this field looks extremely promising. Increasing data volumes are fundamentally changing our approaches, as we have already seen in the *Gaia* context, and this will only be accelerated when facilities such as the Rubin Observatory and its Legacy Survey of Space and Time come online. Likewise, the Roman Space Telescope/*WFIRST* will also contribute to more and larger data sets, while smaller, ground-based facilities will still be required to provide reference data, follow-up opportunities, and complementary temporal sampling.

Beyond photometry and light-curve analyses, spectroscopy represents the next frontier, and also here the future looks bright. Leading on from APOGEE and GALAH, the field is looking forward to exploiting such facilities as 4MOST, WEAVE, DESI,

SDSS-V, MOONS, and others, allowing us to start sampling the variable sky spectroscopically. In particular, radial velocity variability will be tackled with facilities like VELOCE (Cepheids) and, increasingly, *Gaia* (everything else?). It would be very helpful for the field as a whole to share analysis codes and improve accessibility to such tools so as to benefit a greater cross-section of the community.

Spectroscopic time-series observations clearly are the future of this field. Beyond the optical and near-infrared domains, X-ray variability, ultraviolet data, and radio approaches are increasingly hitting the forefront of the field, so we are warned to keep an eye out for new developments in those areas and an open mind as regards one's favourite wavelength range (even including gravitational waves?). Standardisation and cross-calibration remain a concern, but major efforts are undertaken to get this under control. We are eagerly looking forward to the era of the extremely large telescopes. And will artificial intelligence start to play an increasingly important role in our improved understanding? By the time that these developments have matured, a follow-up IAU Symposium will probably be warranted.

Richard de Grijs
Patricia A. Whitelock
Márcio Catelan

Editors

Richard de Grijs
Macquarie University, Sydney, Australia

Patricia A. Whitelock
South African Astronomical Observatory and University of Cape Town, South Africa

Márcio Catelan
Pontificia Universidad Católica de Chile, Santiago, Chile

Organising Committees

Scientific Organising Committee

Richard de Grijs (Chair), Macquarie University, Australia.
László Kiss (Co-chair), Konkoly Observatory, Hungary.
Rachael Beaton, Princeton University, USA.
Márcio Catelan, Pontificia Universidad Católica de Chile, Santiago, Chile.
Xiaodian Chen, National Astronomical Observatories, Chinese Academy of Sciences, Beijing, China.
Gisella Clementini, INAF, Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Italy.
Wolfgang Gieren, Universidad de Concepción, Chile.
Noriyuki Matsunaga, University of Tokyo, Japan.
Victoria Scowcroft, University of Bath, UK.
Róbert Szabó, Konkoly Observatory, Hungary.
Andrzej Udalski, University of Warsaw, Poland.
Patricia Whitelock, South African Astronomical Observatory and University of Cape Town, South Africa.

Local Organising Committee

Róbert Szabó (Chair), Konkoly Observatory, Hungary.
Adrienn Ádám (Co-chair), Konkoly Observatory, Hungary.
László Szabados (Co-chair), Konkoly Observatory, Hungary.
Attila Bódi, Konkoly Observatory, Hungary.
Susmita Das, Konkoly Observatory, Hungary.
Aliz Derekas, Eötvös University, Gothard Astrophysical Observatory, Hungary.
László Kiss, Konkoly Observatory, Hungary.
Adrienn Forró, Konkoly Observatory, Hungary.
Gábor Kovács, Konkoly Observatory, Hungary.
László Molnár, Konkoly Observatory, Hungary.
Emese Plachy, Konkoly Observatory, Hungary.
Dóra Tarczay-Nehéz, Konkoly Observatory, Hungary.
Zsófia Sódorné Bognár, Konkoly Observatory, Hungary.

List of Participants

Abdollahi, Mahdi
Ádám, Rozália
Afanasiev, Anton
Ahmad, Arief
Anderson, Richard
Andriantsaralaza, Miora
Čeki, Atila
Ardern, Steve
Arellano Ferro, Armando
Bódi, Attila
Baruch, John
Beaton, Rachael
Belokurov, Vasily
Benhida, Abdelmajid
Benkő, József
Bhardwaj, Anupam
Bienias, John
Bognár, Zsófia
Bora, Zsófia
Braga, Vittorio Francesco
Bras, Garance
Breuval, Louise
Catanzaro, Giovanni
Chen, Xiaodian
Chung, Chul
Clement, Christine
Clementini, Gisella
Colombo, Claudia
Cruz Reyes, Mauricio
Csoernyei, Geza
Dálya, Gergely
Das, Susmita
de Grijs, Richard
De Somma, Giulia
Deka, Mami
Derekas, Alíz
di Criscienzo, Marcella
Engels, Dieter
Espinoza-Arancibia, Felipe
Evans, Nancy
Eyer, Laurent
Fiorentino, Giuliana
Forró, Adrienn
Freedman, Wendy
Gallenne, Alexandre
Garca Hernández, Antonio
Gholami, Mahtab
Groenewegen, Martin
Hajdu, Gergely
Hintz, Eric
Hocdé, Vincent
Huang, Caroline D.
Iwanek, Patryk
Jeffery, Elizabeth
Jeon, Young-Beom
Jiang, Biwei
Jiménez-Esteban, Fran
Joyce, Meridith
Jurkovic, Monika
Kalup, Csilla
Kervella, Pierre
Khan, Saniya
Kiss, László
Kovács, Gábor
Kovács, Géza
Kurbah, Kerदारis
Latkovic, Olivera
Lee, Jae Woo
Lee, Myung Gyoon
Lengen, Bastian
Lewis, Megan
Liu, Niu
Marconi, Marcella
Martínez-Vázquez, Clara
Mateu, Cecilia
Medina Toledo, Gustavo
Menzies, John
Minniti, Javier
Molnár, László
Mondal, Prapti
Monelli, Matteo
Muraveva, Tatiana
Musella, Ilaria
Nakagawa, Akiharu
Narloch, Weronika
Netzel, Henryka
Olszewska, Justyna
Orosz, Gábor
Owens, Kayla
Paparó, Margit
Park, Seunghyun
Parto, Tahere
Paunzen, Ernst
Pawlak, Michał
Pilecki, Bogumił
Plachy, Emese
Ramezani, Tahereh
Ren, Yi
Ren, Fangzhou
Riess, Adam

Ripepi, Vincenzo
Sánchez-Benavente, Manuel
Sicignano, Teresa
Skowron, Dorota
Son, Junhyuk
Soszyński, Igor
Spetsieri, Zoi Tzogia
Storm, Jesper
Suárez, Juan Carlos
Surdha, Tashveena
Szabó, Róbert
Szabados, László
Tantalo, Maria

Taormina, Mónica
Tokarek, Jakub
Trabucchi, Michele
Trentin, Erasmo
Varga, Vázsony
Vázquez Ramió, Hector
Vinkó, József
Viviani, Giordano
Wang, Shu
Whitelock, Patricia A.
Wielgórski, Piotr
Zgirski, Bartłomiej
Zhang, Zehao

