

of high-functioning ASD such as social difficulties, dislike of change and repetitive/restrictive habits were prominent. Developmental history of the patient and the Autism-Spectrum-Quotient-50 also supported the clinical diagnosis of the ASD. Delusional disorder was excluded, and the therapy organized according to the structure characteristics of the high-functioning ASD which yielded to significant amelioration of depressive symptoms and increased perceived life quality of the patient.

Conclusions: Although coping mechanisms of the patients can be successful, identifying high-functioning ASD-structure even in an old-age can be quite helpful in diagnostic/therapeutic processes. An elaborate discussion of the subject through contemporary literature will be presented.

Disclosure: No significant relationships.

Keywords: differential diagnosis; old age psychiatry; Autism Spectrum Disorder

EPV0930

Evidences of neurodegenerative processes in patients with late-onset schizophrenia and cognitive impairment

V. Pochueva*, M. Savina, V. Sheshenin and N. Cherkasov

FSBSI "Mental Health Research Centre", Geriatric Psychiatry, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1667

Introduction: The large proportion of patients with late-onset schizophrenia (LoS) has cognitive impairment. We hypothesized that this group of patients could have more risk factors associated with neurodegeneration.

Objectives: The aimed to compare various clinical and risk factors in LoS patients with low and relatively preserved cognitive status.

Methods: 28 LoS patients (ICD-11) with duration of disease less than 10 years from a cohort of patients with late onset psychosis underwent clinical assessment (PANSS, HDRS-17), cognitive examination (MMSE, MoCA, FAB, verbal and symbolic memory, trail making test (part A, B)), structured interviewing on risk factors and CT. Hierarchical cluster analysis of cognitive test results was applied. Nonparametric statistic was used to compare control group (24 subjects with signs of psychosis or depression, age 58,1±10,8, 50% females) and patient`s groups.

Results: Patients were divided on two clusters: Cluster 1 with lower cognitive functions (n=20, age 62,2, 94% of females) and Cluster 2 with preserved cognitive functions (n=8, age 56,8, 100% of females). Patients of Cluster 1 were older, had more negative symptoms, higher atrophy scores, higher rate of leukoaraiosis on CT and more history of mild brain injury than patients of Cluster 2 and controls. There was no group differences in age of manifestation, other PANSS scores, rates of social phobia and number of habitual anxiety reactions between clinical groups.

Conclusions: LoS patients with cognitive deficiency had more factors associated with neurodegeneration, in particular history of mild brain injury.

Disclosure: No significant relationships.

Keywords: old age; late-onset schizophrenia; neuridegeneration; risk factors

EPV0932

Sexuality and the Quality of Life in Older People: A Correlational Study

N.E. Boyacıoğlu^{1*}, F. Oflaz², A. Yıldız Karaahmet³, B.K. Hodaei⁴ and Y. Afşin⁵

¹Istanbul Üniversitesi-Cerrahpaşa Health Science Faculty, Gerontology, Büyükcçekmece Yerleşkesi Alkent Mah. Yiğittürk Cad. No:1, Turkey; ²Koç University, School Of Nursing Faculty, Istanbul, Turkey; ³Halic University School of Health Sciences, Midwifery, Istanbul, Turkey; ⁴Istanbul University- Cerrahpasa, Midwifery, Istanbul, Turkey and ⁵Mimar Sinan Fine Arts University, Institute Of Science, Statistics Master Program, Istanbul, Turkey

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1668

Introduction: Sexuality, which is an essential part of human life, is an instinct with the potential to cause or be caused by health problems. Although qualitative and quantitative characteristics of sexual life evolves over time, it may continue until the age of eighties.

Objectives: This descriptive-correlational study aimed to analyze the relationship between general health status, quality of life and sexual life among senior people.

Methods: Study was conducted with the participation of 323 (169 female and 154 male) older people at the age of 65+. The participants were clients of the inpatient and outpatient services in a general hospital in Istanbul. General Health Questionnaire, Arizona Sexual Experiences Scale and Quality of Life Scale in Older People were used to collect data via online survey.

Results: The quality of life was better and sexual problems were lower for the participants who had a partner, higher education level, lower age, a regular job, sufficient income, no chronic disease, who defined their relationship as sufficient, frequently engaged in sexual activity and who considered themselves as attractive.

Conclusions: Sexuality in older people changes over time and continues to hold its importance. Researchers should consider the importance of the quality of life on sexual satisfaction in older people.

Disclosure: No significant relationships.

Keywords: Quality of Life; Older people; health status; sexuality

EPV0933

An observational study of the correlation of efi severity with depression

R. Rasheed, A. Patel* and Y. Shanthakumaran

Rigg Milner Medical Centre, Gp Education And Research, East Tilbury, United Kingdom

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1669

Introduction: Patients with high frailty indices experience poor mental health due to multiple co morbidity and social isolation.

Objectives: This was a retrospective observational analysis that studied the correlation of Electronic frailty indices and GAD scores with Depression scores in a rural population.

Methods: An annual frailty assessment is offered to elderly patients and we screen routinely for anxiety and depression using the PHQ-9 score and GAD score. This was an observational study