

## Relationship between eating behaviours and food and drink consumption in healthy postmenopausal women in a real-life context

Julie Goulet<sup>1</sup>, Véronique Provencher<sup>1</sup>, Marie-Ève Piché<sup>1</sup>, Annie Lapointe<sup>1</sup>, S. John Weisnagel<sup>2</sup>, André Nadeau<sup>2†</sup>, Jean Bergeron<sup>3</sup> and Simone Lemieux<sup>1\*</sup>

<sup>1</sup>*Institute of Nutraceuticals and Functional Foods, Laval University, 2440 Hochelaga Blvd, Québec G1K 7P4, Canada*

<sup>2</sup>*Endocrinology and Diabetes Research Unit, Québec, Canada*

<sup>3</sup>*Lipid Research Center, CHUQ, Québec, Canada*

(Received 14 May 2007 – Revised 28 November 2007 – Accepted 30 November 2007 – First published online 18 February 2008)

Associations between eating behaviours and dietary variables have not been thoroughly investigated in healthy postmenopausal women in a real-life uncontrolled context. To investigate how eating behaviours (cognitive dietary restraint, disinhibition and susceptibility to hunger) were associated with food and drink consumption, energy density and meal pattern in 112 healthy postmenopausal women (age 56.8 (SD 4.4) years) not on hormonal therapy. Women completed a 3 d weighed food record and filled out the Three-Factor Eating Questionnaire. The sample was divided according to the median of the distribution of cognitive dietary restraint and disinhibition (9 and 6 respectively). Both subgroups of women with high restraint level (presenting either high or low disinhibition) consumed a diet with a lower energy density than subgroups of women with lower restraint level. Women with high restraint–low disinhibition had a lower consumption of red meat and processed meat and a lower consumption of diet soft drinks than women with low restraint–high disinhibition. They were also characterised by a higher intake of whole grains than women with high restraint–high disinhibition and than women with lower restraint level (with either high or low disinhibition). Women with high restraint–high disinhibition levels showed differences in dietary variables when compared with subgroups of women with lower restraint level, namely for refined grains and diet soft drinks. We conclude that in healthy postmenopausal women, dietary consumption of specific food and drink may be related to particular eating behaviours. Women with high restraint and low disinhibition levels generally showed the most healthy dietary pattern.

**Postmenopausal women: Eating behaviours: Food habits: Meal pattern: Energy density: Three-Factor Eating Questionnaire**

The prevalence of obesity in Western countries has steadily increased in the last decade, and has become a major public health problem<sup>(1–3)</sup>. Obesity, and more specifically abdominal obesity, has been shown to increase the risk of CVD and type 2 diabetes<sup>(4,5)</sup> and in women, CVD remains the first cause of mortality after menopause<sup>(6–8)</sup>. While dietary treatment is the basic therapeutic approach for obesity, its long-term impact on weight management is disappointing<sup>(9)</sup>. In fact, approximately 20–30% of overweight or obese individuals succeed in maintaining their weight loss after 1–3 years following a hypoenergetic diet, and a lower percentage succeed after 3–5 years<sup>(9)</sup>. On the other hand, dieting in order to control weight is very prevalent in our modern society, especially among women. In fact, results from the National Health and Nutrition Examination Survey 2001–2 underline that restraining food intake is the most prevalent weight-control practice, either in women wanting to lose weight or trying not to gain weight<sup>(10)</sup>.

The Three-Factor Eating Questionnaire is one of the most widely used scales in behavioural research, developed in order to measure cognitive dietary restraint, disinhibition and susceptibility to hunger<sup>(11)</sup>. Even if each method to assess eating beha-

viours has its strengths and weaknesses, results from a study comparing four questionnaires of dietary restraint measurement has shown that the Three-Factor Eating Questionnaire cognitive dietary restraint scale was the most valid measure of the intent to diet and of actual energy restriction<sup>(12)</sup>.

The associations between eating behaviours and body-weight management have been studied extensively but some inconsistency remains. More precisely, some intervention studies underlined that dietary restraint was a predictor of weight loss<sup>(13)</sup>. On the other hand, other studies found no association between cognitive dietary restraint and BMI<sup>(14,15)</sup>. Higher scores of disinhibition, which is defined as a loss of control over eating in response to different stimuli that leads to an overconsumption of food<sup>(11)</sup>, have been associated more systematically to higher BMI<sup>(15–19)</sup>. The inconsistent associations between restraint and BMI could be partly explained by the fact that in some studies, increased restriction has been associated with increased disinhibition<sup>(20,21)</sup>.

Since change in food choices is the cornerstone of obesity management and because eating behaviours are related to obesity, better knowledge with regard to eating behaviours

\* **Corresponding author:** Dr Simone Lemieux, fax +1 418 656 5877, email Simone.Lemieux@al.n.ulaval.ca

† Deceased 25 September 2006.

and their relationship with food and drink consumption and meal pattern could be helpful to improve dietary treatment of obesity. Some studies evaluated the relationship between eating behaviours and food consumption<sup>(22–27)</sup>. Accordingly, cognitive dietary restraint has been previously associated with frequent use of reduced-energy and reduced-fat foods<sup>(26,28)</sup>. Furthermore, French *et al.* demonstrated that subjects who self-reported that they were dieting to lose weight (which is related to increased dietary restraint)<sup>(29)</sup> had lower self-reported intake of several high-fat/high-energy foods, including sweets, meat, soft drinks, French fries and dairy products<sup>(29)</sup>.

In most studies, associations between eating behaviours, BMI and food intakes have been examined in premenopausal women<sup>(23,25–27,30–32)</sup> and studies have generally not considered a wide set of dietary variables and rather focused on energy intake, macronutrient composition of the diet and only a few food groups. Fewer studies have been performed in healthy postmenopausal women not under treatment, under 'real-life settings'. Also, to our knowledge, no study has yet investigated whether the presence of disinhibition could modulate the association between restraint and the pattern of food consumption among postmenopausal women. Therefore, the present cross-sectional study aimed at examining the relationship between eating behaviours (cognitive dietary restraint, disinhibition and susceptibility to hunger) and habitual food and drink consumption as well as energy and nutrient intake, energy density and meal pattern. We also wanted to verify the hypothesis that the presence of disinhibition could modulate the association between restraint and dietary variables under study.

## Research design and methods

### Subjects

The present study was conducted in a sample of 112 postmenopausal women (aged between 46 and 68 years) in a real-life uncontrolled context recruited through the media in the Quebec City metropolitan area. Those reporting that they did not have menses for at least 1 year were considered as postmenopausal and were included in the study. A measure of the follicle-stimulating hormone was used to confirm the menopausal status (follicle-stimulating hormone value between 28 and 127 IU/l). All women included in the present study were free from metabolic disorders, were not using any type of hormonal therapy and were not under treatment for CHD, diabetes, dyslipidaemias or endocrine disorders (except stable thyroid disease). Five women included in the present study were smokers. None of the participants had received a diagnosis of type 2 diabetes before the study. All participants signed an informed consent document before entering the study, which was approved by the Laval University Medical Centre and the Laval University Research Ethics Committees.

### Dietary profile

Food intake was assessed by a 3 d weighed food record, which was completed during two weekdays and one weekend day. The food record was explained and reviewed

by the study registered dietitian. Copies of food record examples were also provided to each subject. In addition, participants were encouraged to consume the usual amount of typical foods and drinks. The food record included a section for recording information about recipes. Women were asked to weigh foods with a scale provided by the dietitian. Evaluation of nutrient intake derived from the 3 d food record was performed using the Nutrition Data System for Research software (version 4.03, developed by the Nutrition Coordination Center, University of Minnesota, Minneapolis, MN, Food and Nutrient Database 31, released in November 2000)<sup>(33)</sup>. Each 3 d food record was reviewed to determine daily portions for 121 categories of food items that were further regrouped in twenty-eight food groups. Food categories used were adapted from the food list derived from a FFQ developed by Goulet *et al.*<sup>(34)</sup> (see Appendix 1 for more details).

Each subject also had to identify in the 3 d food record the time of day and location for each meal consumed. Total energy intake for each day was divided by the total weight of the foods and beverages reported to determine daily energy density values<sup>(35)</sup>. Similarly, energy density was also calculated for each meal.

### Anthropometric measurements

Height (cm), body weight (kg) and BMI (kg/m<sup>2</sup>) were determined following the procedures recommended at the Airlie Conference<sup>(36)</sup>. Anthropometric measurements were performed after the completion of the weighed food record.

### Questionnaire about history of weight

A questionnaire about weight and diet history was administered by a registered dietitian. Participants had to indicate whether they had previously been on a diet to lose weight and when was their last dieting attempt. They also had to indicate their body weight at age 20 years.

### Three-Factor Eating Questionnaire

Postmenopausal women filled out a French version at home of the Three-Factor Eating Questionnaire. The Three-Factor Eating Questionnaire is a fifty-one-item questionnaire developed by Stunkard & Messick in 1985<sup>(11)</sup>. The purpose of this questionnaire is to assess three factors related to cognitions and behaviours associated with eating. These factors are cognitive dietary restraint, disinhibition, and susceptibility to hunger. More precisely, cognitive dietary restraint is a conscious control over food intake with concerns about shape and weight (twenty-one items, score ranging from 0 to 21). Disinhibition is an overconsumption of food in response to a variety of stimuli, such as emotional stress, associated with a loss of control on food intake (sixteen items, score ranging from 0 to 16). Finally, susceptibility to hunger refers to food intake in response to feelings and perceptions of hunger (fourteen items, score ranging from 0 to 14)<sup>(11)</sup>. This questionnaire has been validated, and all three of these scales have good test–retest reliability<sup>(11,37,38)</sup>.

### Statistical analyses

Pearson correlation analyses were performed to assess how eating behaviours were associated with BMI, energy intake and energy density. The whole sample of women was divided according to the median of the distribution of cognitive dietary restraint (score of 9) and of disinhibition (score of 6), resulting in the formation of four subgroups (low restraint–low disinhibition; low restraint–high disinhibition; high restraint–low disinhibition; high restraint–high disinhibition). Comparisons between groups were performed by ANOVA. In the presence of significant effects, Duncan's multiple comparison test was used to determine precisely the location of significant differences. The  $\chi^2$  frequency procedure was used to compare the frequency of dieters in the four groups described above. Moreover, participants were also divided in two subgroups using the median of the distribution of susceptibility to hunger score (4 units). Student's *t* tests were then computed to evaluate differences between these two subgroups.

Finally, ordinary least squares regression was performed to determine the independent contribution of cognitive dietary restraint, disinhibition and their interaction to the determination of dietary variables studied. All analyses were performed with the SAS statistical package version 8.02 (SAS Institute, Inc., Cary, NC, USA).

### Results

Postmenopausal women had a mean age of 56.8 (SD 4.4) years, a mean menopause duration of 8.3 (SD 6.9) years and a mean BMI of 28.5 (SD 5.9) kg/m<sup>2</sup>. BMI was significantly correlated with dietary restraint ( $r = -0.33$ ;  $P=0.0003$ ), disinhibition ( $r = 0.49$ ;  $P<0.0001$ ) and susceptibility to hunger ( $r = 0.39$ ;  $P<0.0001$ ). Neither eating behaviours nor BMI were significantly correlated with energy intake. However, cognitive dietary restraint and disinhibition were both significantly associated with daily energy density of the diet ( $r = -0.30$ ,  $P=0.002$ ;  $r = 0.21$ ,  $P=0.04$ , respectively). Because both cognitive dietary restraint and disinhibition were significantly

associated with energy density, regression analysis was also performed to determine the contribution of restraint, disinhibition and the interaction between restraint and disinhibition to the variance in energy density. Results showed that only cognitive dietary restraint tended ( $P=0.06$ ) to be an independent predictor of energy density. Neither disinhibition ( $P=0.97$ ) nor the interaction between restraint and disinhibition ( $P=0.51$ ) was a predictor of energy density.

The whole sample of women was separated according to median values of the distribution of restraint and disinhibition levels. Accordingly, four subgroups were compared: low restraint–low disinhibition; low restraint–high disinhibition; high restraint–low disinhibition; high restraint–high disinhibition. It is shown in Table 1 that women with low restraint and low disinhibition scores were characterised by a higher BMI than women from the other three groups ( $P<0.05$ ). Also, women with low restraint and high disinhibition gained significantly more weight since the age of 20 years than women characterised by low restraint and low disinhibition and than both subgroups of women with a higher restraint. Women with low restraint and high disinhibition also had a higher BMI at the age of 20 years than women with low restraint and low disinhibition ( $P<0.05$ ). It is also shown in Table 1 that both groups of women with high restraint level had a lower energy density than the two groups of women with lower restraint level. Although all women were weight stable at the time of investigation, some of them had already experienced dieting in the past. It was found that women with low restraint and low disinhibition scores were less susceptible to have been dieting in the past than women from the other three groups.

Table 2 presents differences in consumption of food and drinks according to restraint and disinhibition levels. Women with high restraint and low disinhibition were different from women with lower restraint for some dietary variables. In fact, they had a lower consumption of red meat and processed meat than women with low restraint and high disinhibition. They also presented higher intakes of whole grains than both groups of women with low restraint (i.e. low

**Table 1.** Dietary characteristics according to the level of dietary restraint and disinhibition in 112 postmenopausal women (Mean values and standard deviations)

	↓ Restraint/ ↓ disinhibition (n 18)		↓ Restraint/ ↑ disinhibition (n 32)		↑ Restraint/ ↓ disinhibition (n 38)		↑ Restraint/ ↑ disinhibition (n 23)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Age (years)	55.7	4.8	56.9	4.5	57.1	4.3	56.3	3.6
Age at menopause (years)	48.2	5.1	48.2	6.4	49.5	6.4	47.2	5.6
BMI (kg/m <sup>2</sup> )	25.7 <sup>b</sup>	4.9	32.6 <sup>a</sup>	6.8	26.3 <sup>b</sup>	5.0	28.8 <sup>b</sup>	3.8
Body weight (kg)	67.4 <sup>b</sup>	16.2	82.7 <sup>a</sup>	17.9	67.8 <sup>b</sup>	13.8	72.4 <sup>b</sup>	10.7
Body weight at 20 years (kg)	55.8 <sup>b</sup>	13.9	59.0 <sup>a</sup>	9.4	55.1 <sup>b</sup>	7.5	58.1 <sup>b</sup>	6.8
Change in body weight since age 20 years (kg/year)	0.30 <sup>b</sup>	0.19	0.65 <sup>a</sup>	0.49	0.35 <sup>b</sup>	0.36	0.43 <sup>b</sup>	0.26
Proportion of dieters (%)		33 <sup>a</sup>		78 <sup>b</sup>		71 <sup>b</sup>		75 <sup>b</sup>
Energy intake (kJ)	8370	2178	8852	1981	7721	1503	7867	1952
Protein (% of energy)	17.3	4.2	18.0	2.1	18.0	3.1	19.4	3.0
Carbohydrates (% of energy)	48.3	6.1	47.5	6.3	49.8	6.7	47.7	5.3
Fat (% of energy)	33.9	5.7	35.0	4.4	32.3	6.1	33.0	5.0
Number of meals (per d)	5.2	1.4	5.1	1.3	5.5	1.6	5.6	1.5
Daily food intake (g)	2525	617	2639	591	2838	612	2790	805
Daily energy density (kJ/g)	3.38 <sup>a</sup>	0.70	3.46 <sup>a</sup>	0.80	2.81 <sup>b</sup>	0.67	2.94 <sup>b</sup>	0.68

<sup>a,b</sup> Values with unlike superscript letters are significantly different ( $P<0.05$ ).

**Table 2.** Differences in consumption of food and drinks according to restraint and disinhibition levels (*n* 112) (Mean values and standard deviations)

Food and drinks	↓ Restraint/ ↓ disinhibition ( <i>n</i> 18)		↓ Restraint/ ↑ disinhibition ( <i>n</i> 32)		↑ Restraint/ ↓ disinhibition ( <i>n</i> 38)		↑ Restraint/ ↑ disinhibition ( <i>n</i> 24)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Dairy products	1.6	1.3	1.7	0.8	1.9	0.8	2.1	1.0
Partly skimmed/skimmed	1.0	0.9	1.0	0.7	1.3	0.8	1.4	0.8
Whole	0.7	0.6	0.7	0.6	0.7	0.6	0.7	0.5
Meat								
Red meat	1.2 <sup>a,b</sup>	0.8	1.3 <sup>a</sup>	0.8	0.8 <sup>b</sup>	0.6	1.2 <sup>a,b</sup>	0.8
Processed meat	0.6 <sup>b</sup>	0.8	1.1 <sup>a</sup>	1.2	0.5 <sup>b</sup>	0.6	0.8 <sup>a,b</sup>	0.9
Poultry	0.8	0.8	1.1	1.3	0.7	0.8	0.8	1.0
Fish	0.9	1.1	1.0	1.3	1.3	1.4	1.0	1.1
Substitutes	0.4	0.5	0.5	0.5	0.5	0.6	0.4	0.5
Grain products	4.4	2.1	4.4	1.2	4.8	1.7	3.5	1.7
Refined grains	2.5 <sup>a</sup>	1.5	2.3 <sup>a,b</sup>	1.7	1.7 <sup>a,b</sup>	1.3	1.5 <sup>b</sup>	1.5
Whole grains	1.9 <sup>b</sup>	1.5	2.1 <sup>b</sup>	2.1	3.1 <sup>a</sup>	1.5	2.0 <sup>b</sup>	1.3
Fruits	3.1	2.3	2.4	1.2	3.0	1.0	2.9	1.5
Fruit juice	1.1	1.0	0.7	0.8	0.9	0.8	1.3	1.1
Vegetables	2.5	1.4	3.3	1.4	3.1	1.3	3.2	1.2
Fat added	3.2	2.9	6.0	3.9	5.7	3.3	4.9	3.1
Desserts	0.7 <sup>b</sup>	0.6	1.3 <sup>a</sup>	1.1	0.7 <sup>a,b</sup>	0.6	0.7 <sup>a,b</sup>	0.5
Snacks	0.2	0.3	0.4	0.6	0.2	0.4	0.2	0.3
Soft drinks	0.1 <sup>b</sup>	0.2	0.8 <sup>a</sup>	1.1	0.2 <sup>b</sup>	0.2	0.3 <sup>b</sup>	0.5
Regular	0.1	0.2	0.2	0.5	0.1	0.2	0.1	0.3
Diet	0.0 <sup>b</sup>	0.0	0.6 <sup>a</sup>	1.1	0.1 <sup>b</sup>	0.2	0.2 <sup>b</sup>	0.4

<sup>a,b</sup> Mean values with unlike superscript letters are significantly different ( $P < 0.05$ ).

restraint–low disinhibition and low restraint–high disinhibition). In addition, women with high restraint and low disinhibition consumed fewer diet soft drinks than women with low restraint and high disinhibition.

Women with high restraint and high disinhibition also showed some differences when compared with women with lower restraint level. In fact, they had lower consumption of refined grains than women with low restraint and low disinhibition and lower consumption of diet soft drinks than women with low restraint and high disinhibition. When the two groups of women with high restraint values were directly compared with each other it was found that only the consumption of whole grains differed between the two groups (women with high restraint and low disinhibition having a higher consumption).

Low restraint combined with either high or low disinhibition differentially influenced dietary intake. Specifically, women with both low restraint and low disinhibition consumed less processed meat, less dessert and fewer soft drinks than women with low restraint and high disinhibition.

Subjects were also compared according to their susceptibility to hunger score (high *v.* low). Women characterised by a low susceptibility to hunger score had significantly lower total energy intakes than women with a higher score (7695 (SD 1685) *v.* 8648 (SD 1982) kJ respectively;  $P = 0.007$ ) (Table 3). No significant differences were observed for macronutrient partition, energy density and number of meals per d. However, energy partition at breakfast and lunch tended to be significantly different between the two groups of women (breakfast: 32.6 (SD 26.9) % of daily energy for low *v.* 24.8 (SD 19.1) % for high susceptibility to hunger,  $P = 0.08$ ; lunch: 23.2 (SD 11.7) % of daily energy for low *v.* 27.1 (SD 9.8) % for high susceptibility to hunger,  $P = 0.056$ ).

Consumption of meat substitutes (i.e. legumes, tofu, nuts, seeds) ( $P = 0.04$ ), desserts ( $P = 0.009$ ) and diet soft drinks

( $P = 0.04$ ) were significantly higher and consumption of fruit juices was significantly lower ( $P = 0.03$ ) in subjects with a high susceptibility to hunger score than in women with lower susceptibility to hunger (Table 4).

## Discussion

The main purpose of the present study was to examine how eating behaviours were associated with dietary profile, meal pattern and energy density in healthy postmenopausal

**Table 3.** Dietary variables according to the level of susceptibility to hunger in postmenopausal women (*n* 112)

(Mean values and standard deviations)

	Susceptibility to hunger			
	Low ( <i>n</i> 55)		High ( <i>n</i> 57)	
	Mean	SD	Mean	SD
Energy intake (kJ)	7695	1685	8648*	1982
Protein (% of energy)	18.3	3.9	18.0	2.1
Carbohydrates (% of energy)	48.7	6.4	48.2	6.0
Fat (% of energy)	32.7	5.3	34.3	5.4
Number of meals (per d)	5.2	1.4	5.6	1.6
Daily food intake (g)	2657	677	2781	633
Daily energy density (kJ/g)	3.01	0.73	3.22	0.80

\* Mean value was significantly different from that of the group with low susceptibility to hunger ( $P \leq 0.05$ ).

**Table 4.** Differences in daily consumption of food and drinks according to the level of susceptibility to hunger in postmenopausal women (*n* 112) (Mean values and standard deviations)

Food and drink (portions per d)	Low susceptibility to hunger ( <i>n</i> 55)		High susceptibility to hunger ( <i>n</i> 57)		<i>P</i>
	Mean	SD	Mean	SD	
Dairy products	1.8	1.0	1.9	1.0	0.6
Partly skimmed/ skimmed	1.2	0.9	1.2	0.8	0.9
Whole	0.6	0.5	0.8	0.6	0.3
Meat					
Red meat	1.6	1.5	1.8	1.5	0.5
Processed meat	0.7	0.8	0.8	1.0	0.7
Poultry	0.7	0.9	1.0	1.3	0.2
Fish	1.2	1.3	1.0	1.3	0.5
Substitutes	0.3	0.4	0.6	0.6	0.04
Grain products	4.2	1.7	4.4	1.6	0.4
Refined grains	1.9	1.5	2.0	1.4	0.8
Whole grains	2.3	1.5	2.4	1.3	0.6
Fruits	3.0	1.4	2.7	1.5	0.3
Fruit juice	1.2	1.1	0.8	0.7	0.03
Vegetables	3.2	1.5	3.3	1.5	0.8
Fat added	4.9	2.9	5.8	3.9	0.2
Desserts	0.7	0.6	1.1	0.9	0.009
Snacks	0.2	0.3	0.3	0.5	0.07
Soft drinks	0.2	0.6	0.5	0.8	0.08
Regular	0.1	0.4	0.1	0.2	0.09
Diet	0.1	0.5	0.4	0.7	0.04

women. The present results showed that women characterised by a high restraint level (presenting either high or low disinhibition) had a diet with a lower energy density as compared with women with lower restraint level. Differences in the consumption of red meat and processed meat, whole grains, refined grains and diet soft drinks were also observed between women with high restraint compared with those with lower restraint. Contrary to our hypothesis, the presence of disinhibition in women with high restraint did not seem to have a major impact on the overall pattern of food and drink consumption.

Our data also revealed that there was no significant association between eating behaviours and energy intake. Although some studies have shown that highly restrained individuals are more susceptible to report a lower consumption of energy<sup>(25,26,39–41)</sup>, a lower energy intake has not been systematically reported in highly restrained eaters<sup>(42)</sup>. The lack of consistency in reported associations between dietary restraint and actual physiological energy restriction might be explained by the fact that dietary restraint is a heterogeneous concept and can also reflect a state of perceived deprivation (eating less than wanted without creating energy deficit)<sup>(43)</sup>. In addition, it is recognised that restraint could be imposed either by those who are currently dieting to lose weight or in those who have successfully lost weight and wish to maintain their weight status. Also some restrained eaters imposed restraint simply because they do not want to gain weight<sup>(10)</sup>. It is also possible that highly restrained subjects are more likely to underreport their dietary intake<sup>(44)</sup>. However, the significant negative association between cognitive dietary restraint and BMI in the present study suggests that women with higher cognitive dietary restraint have an energy intake which is more accurately adapted to actual energy needs.

The present results also showed that women with low restraint and high disinhibition had the highest BMI and the more important body-weight gain since the age of 20 years, which suggests persistent positive energy balance. From the present results, it is difficult to determine whether the gain in body weight was a factor responsible for the determination of eating behaviours or whether eating behaviours had an impact on body-weight variation. Furthermore, we can not assume that eating behaviours measured after menopause are representative of eating behaviours measured at the age of 20 years since previous studies have shown that eating behaviours are likely to change over time<sup>(31,45)</sup>.

Although eating behaviours were not significantly associated with energy intake, the present results showed that cognitive dietary restraint was negatively and disinhibition was positively associated with energy density of the diet. Furthermore, regression analyses revealed that only cognitive dietary restraint tended to be a significant predictor of energy density. This is concordant with the fact that women with high restraint, irrespective of disinhibition level, had a diet with lower energy density than women with lower restraint level. The lower energy density of the diet of women with high restraint might be one of the factors explaining the significant association between higher cognitive dietary restraint and lower BMI. In fact, dietary energy density has been suggested as a determinant of energy intake and of body weight<sup>(46–48)</sup>.

With regard to food and drink consumption, we found some differences between women with high *v.* low cognitive dietary restraint. The lower consumption of red meat and processed meat and the higher consumption of whole grains in women with high restraint and low disinhibition when compared with women with lower restraint level are suggestive of a healthier dietary pattern. In fact, lower consumption of red meat and increased consumption of whole grains are components of the Alternate Healthy Eating Index that has been previously described as a dietary pattern associated with lower risk of major chronic diseases<sup>(49)</sup>. However, the lower consumption of diet soft drinks in women with high cognitive restraint is somehow discordant with previous studies suggesting that restrained eaters are more likely to avoid high-energy food items<sup>(50)</sup> and to prefer food generally labelled as 'low-calorie'<sup>(50)</sup>. On the other hand, the lower consumption of diet soft drinks might be an additional indicator of an overall healthier diet. In fact, it has been recently shown by Dhingra *et al.*<sup>(51)</sup> that the incidence of the metabolic syndrome was increased in middle-aged adults consuming at least one soft drink per d, regardless of whether it was regular or diet.

Contrary to our hypothesis, the presence of disinhibition did not seem to influence to a large extent the association between cognitive dietary restraint and food and drink consumption. In fact, the only difference between women with high restraint–low disinhibition and women with high restraint–high disinhibition was a higher whole grain intake in women with low disinhibition. Therefore, it appears that the presence of cognitive dietary restraint might be a more important determinant of food and drink consumption than disinhibition. This is concordant with regression analysis showing that cognitive dietary restraint tended to be a significant predictor of energy density whereas neither disinhibition nor the interaction between restraint and disinhibition predicted energy density. However, from the present results it can be postulated that the impact of

disinhibition on food pattern might vary according to cognitive dietary restraint. In fact, among women with lower restraint, the presence of disinhibition was associated with the consumption of more processed meat, more desserts and more diet soft drinks which can overall explain, at least partially, the increased BMI observed in women with low restraint and high disinhibition when compared with the other three groups of women tested.

The relationship between meal pattern and eating behaviours has not been thoroughly investigated so far. It has been reported that subjects with either high or low dietary restraint eat similar numbers of meals per d, but restrained eaters consumed more snacks<sup>(28)</sup>. In the present study, no significant difference in variables related to meal pattern was noted between groups of women separated on the basis of eating behaviour scores. However, women characterised by a low susceptibility to hunger consumed less energy and tended to consume a larger proportion of energy at breakfast than women with high susceptibility to hunger. It has been previously reported that the proportion of intake in the morning was negatively correlated with total daily intake<sup>(52)</sup>. De Castro also reported that human subjects demonstrated less satiety from a given amount of food later in the day than earlier<sup>(53)</sup>, suggesting that a larger proportion of total intake in the morning is related to a smaller total intake. According to the present results, women with low susceptibility to hunger score consumed less energy on a daily basis and had a lower body weight. Therefore, we could hypothesise that women with high susceptibility to hunger could benefit from eating a larger amount in the morning when the satiating value of food is higher and, thus, this may contribute to regulate their energy intake and probably their body weight. However, we agree that changes in daily meal partition may be difficult to achieve and maintain in a real-life context. Further studies will be needed to understand the interrelationship between susceptibility to hunger, meal pattern and body-weight regulation.

### Conclusion

The present results suggest associations between eating behaviours, energy and nutrient intakes, dietary food choices and meal pattern in healthy postmenopausal women. In the present study, postmenopausal women with higher cognitive dietary restraint appear to have a healthier dietary profile while the presence of disinhibition and susceptibility to hunger were rather associated with a less healthy dietary profile. Although nutritional advice *per se* is a key component in weight management, results from the present study suggest that intervention aiming at modifying eating behaviours may also influence food choices that will in turn impact on the regulation of energy balance and body weight.

### Acknowledgements

J. G. is a recipient of a studentship from the Fonds de la Recherche en Santé du Québec (FRSQ). V. P. is a recipient of a studentship from FRSQ and A. L. is a recipient of a studentship from the Canadian Institute of Health Research and the FRSQ. The present study was supported by the Canada

Institutes of Health Research (MOP-37957) and by the Heart and Stroke Foundation of Canada.

The authors would like to express their gratitude to the subjects for their excellent collaboration and to the staff of the Lipid Research Centre, the Physical Activity Sciences Laboratory, and the Diabetes Research Unit for their contribution to the present study. We especially want to thank Louise Corneau MSc RD, Nancy Gilbert MSc RD, Rolande Couture, Danielle Aubin, Fanny Therrien, Marie Tremblay and George Cousineau for their help in the collection and analysis of the data.

None of the authors had a personal or professional conflict of interest.

J. G. performed data analysis and drafted the manuscript. M. E. P. and V. P. participated in data collection. A. L. participated in data analysis. S. J. W., A. N., J. B. and S. L. conceived the study, and participated in its design and coordination.

### References

1. Ledoux M (1998) Poids corporel. In *Équité Sociale et de Santé 1998 (Body Weight: Social and Health Questions)*, chapter 8. Québec: Institut de la Statistique du Québec.
2. Flegal KM, Carroll MD, Ogden CL & Johnson CL (2002) Prevalence and trends in obesity among US adults, 1999–2000. *JAMA* **288**, 1723–1727.
3. Tjepkema M & Shields M (2005) *Nutrition Findings from Canadian Community Health Survey Adult Obesity in Canada: Measured Height and Weight*. Ottawa: Statistics Canada.
4. World Health Organization (1997) *Obesity: Preventing and Managing the Global Epidemic. Report on a WHO Consultation on Obesity*. Geneva: World Health Organization.
5. Kissebah AH (1996) Intra-abdominal fat: is it a major factor in developing diabetes and coronary artery disease? *Diabetes Res Clin Pract* **30**, Suppl., 25–30.
6. Fondation des Maladies du Coeur du Canada (2005) Les maladies cardiovasculaires et accidents vasculaires cérébraux au Canada (Cardiovascular illnesses and cerebral vascular accidents in Canada). 4-7-2005. Ottawa, Canada: Fondation des Maladies du Coeur du Canada. www.fmcoeur.ca
7. Eaker E, Chesebro JH, Sacks FM, Wenger NK, Whisnant JP & Winston M (1994) Special report: cardiovascular disease in women. Special writing group. *Heart Dis Stroke* **3**, 114–119.
8. Mosca L, Manson JE, Sutherland SE, Langer RD, Manolio T, Barrett-Connor E & Writing Group (1997) Cardiovascular disease in women: a statement for healthcare professionals from the American Heart Association. *Circulation* **96**, 2468–2482.
9. Wing RR & Phelan S (2005) Long-term weight loss maintenance. *Am J Clin Nutr* **82**, Suppl. 1, 222S–225S.
10. Weiss EC, Galuska DA, Khan LK & Serdula MK (2006) Weight-control practices among U.S. adults, 2001–2002. *Am J Prev Med* **31**, 18–24.
11. Stunkard AJ & Messick S (1985) The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J Psychosom Res* **29**, 71–83.
12. Williamson DA, Martin CK, York-Crowe E, Anton SD, Redman LM & Han H (2007) Measurement of dietary restraint: validity tests of four questionnaires. *Appetite* **48**, 183–192.
13. Drummond S, Dixon K, Griffin J & De Looy A (2004) Weight loss on an energy-restricted, low-fat, sugar-containing diet in overweight sedentary men. *Int J Food Sci Nutr* **55**, 279–290.

14. Dykes J, Brunner EJ, Martikainen PT & Wardle J (2004) Socio-economic gradient in body size and obesity among women: the role of dietary restraint, disinhibition and hunger in the Whitehall II study. *Int J Obes* **28**, 262–268.
15. Provencher V, Drapeau V, Tremblay A, Despres JP & Lemieux S (2003) Eating behaviors and indexes of body composition in men and women from the Quebec family study. *Obes Res* **11**, 783–792.
16. Lindroos AK, Lissner L, Mathiassen ME, Karlsson J, Sullivan M, Bengtsson C & Sjöström L (1997) Dietary intake in relation to restrained eating, disinhibition, and hunger in obese and non-obese Swedish women. *Obes Res* **5**, 175–182.
17. Carmody TP, Brunner RL & St Jeor ST (1995) Dietary helplessness and disinhibition in weight cyclers and maintainers. *Int J Eat Disord* **18**, 247–256.
18. Lawson OJ, Williamson DA, Champagne CM, DeLany JP, Brooks ER, Howat PM, Wozniak PJ, Bray GA & Ryan DH (1995) The association of body weight, dietary intake, and energy expenditure with dietary restraint and disinhibition. *Obes Res* **3**, 153–161.
19. Hays NP, Bathalon GP, McCrory MA, Roubenoff R, Lipman R & Roberts SB (2002) Eating behavior correlates of adult weight gain and obesity in healthy women aged 55–65 y. *Am J Clin Nutr* **75**, 476–483.
20. Westenhoefer J, Stunkard AJ & Pudel V (1999) Validation of the flexible and rigid control dimensions of dietary restraint. *Int J Eat Disord* **26**, 53–64.
21. McLean JA, Barr SI & Prior JC (2001) Cognitive dietary restraint is associated with higher urinary cortisol excretion in healthy premenopausal women. *Appetite* **73**, 7–12.
22. Borg P, Fogelholm M & Kukkonen-Harjula K (2004) Food selection and eating behaviour during weight maintenance intervention and 2-y follow-up in obese men. *Int J Obes Relat Metab Disord* **28**, 1548–1554.
23. Tuschl RJ, Platte P, Laessle RG, Stichler W & Pirke KM (1990) Energy expenditure and everyday eating behavior in healthy young women. *Am J Clin Nutr* **52**, 81–86.
24. French SA, Jeffery RW & Wing RR (1994) Food intake and physical activity: a comparison of three measures of dieting. *Addict Behav* **19**, 401–409.
25. Moreira P, de Almeida MD & Sampaio D (2005) Cognitive restraint is associated with higher intake of vegetables in a sample of university students. *Eat Behav* **6**, 229–237.
26. Rideout CA, McLean JA & Barr SI (2004) Women with high scores for cognitive dietary restraint choose foods lower in fat and energy. *J Am Diet Assoc* **104**, 1154–1157.
27. Contento IR, Zybert P & Williams SS (2005) Relationship of cognitive restraint of eating and disinhibition to the quality of food choices of Latina women and their young children. *Prev Med* **40**, 326.
28. Kirkley BG, Burge JC & Ammerman A (1988) Dietary restraint, binge eating and dietary behavior patterns. *Int J Eat Disord* **7**, 771–778.
29. French SA, Jeffery RW & Forster JL (1994) Dieting status and its relationship to weight, dietary intake, and physical activity changes over two years in a working population. *Obes Res* **2**, 135–144.
30. Westterterp-Plantenga MS, Wijckmans-Duijsens NE, Verboeket-van de Venne WP, de Graaf K, het Hof KH & Weststrate JA (1998) Energy intake and body weight effects of six months reduced or full fat diets, as a function of dietary restraint. *Int J Obes Relat Metab Disord* **22**, 14–22.
31. Drapeau V, Provencher V, Lemieux S, Despres JP, Bouchard C & Tremblay A (2003) Do 6-y changes in eating behaviors predict changes in body weight? Results from the Quebec Family Study. *Int J Obes Relat Metab Disord* **27**, 808–814.
32. Beiseigel JM & Nickols-Richardson SM (2004) Cognitive eating restraint scores are associated with body fatness but not with other measures of dieting in women. *Appetite* **43**, 47–53.
33. Schakel SF, Sievert YA & Buzzard IM (1988) Sources of data for developing and maintaining a nutrient database. *J Am Diet Assoc* **88**, 1268–1271.
34. Goulet J, Nadeau G, Lapointe A, Lamarche B & Lemieux S (2004) Validity and reproducibility of an interviewer-administered food frequency questionnaire for healthy French-Canadian men and women. *Nutr J* **3**, 3–13.
35. Ledikwe JH, Blanck HM, Khan LK, Serdula MK, Seymour JD, Tohill BC & Rolls BJ (2005) Dietary energy density determined by eight calculation methods in a nationally representative United States population. *J Nutr* **135**, 273–278.
36. Lohman TG, Roche AF & Martorell R (1988) The Airlie (VA) consensus conference. In *Anthropometric Standardization Reference Manual*, pp. 39–80 [TG Lohman, AF Roche and R Martorell, editors]. Champaign, IL: Human Kinetics Publishers.
37. Laessle RG, Tuschl RJ, Kotthaus BC & Pirke KM (1989) A comparison of the validity of three scales for the assessment of dietary restraint. *J Abnorm Psychol* **98**, 504–507.
38. Lluch A (1995) *Identification des Conduites Alimentaire par Approches Nutritionnelles et Psychométriques: Implications Thérapeutiques et Préventives dans l'Obésité Humaines (Identification of Eating Patterns by Nutritional and Psychometric Approaches: Implications for the Prevention and Treatment of Human Obesity)*. PhD Thesis, University of Nancy 1, France.
39. de Lauzon B, Romon M, Deschamps V, Lafay L, Borys JM, Karlsson J, Ducimetière P, Charles MA & Fleurbaix Laventie Ville Sante Study Group (2004) The Three-Factor Eating Questionnaire-R18 is able to distinguish among different eating patterns in a general population. *J Nutr* **134**, 2372–2380.
40. Klesges RC, Isbell TR & Klesges LM (1992) Relationship between dietary restraint, energy intake, physical activity, and body weight: a prospective analysis. *J Abnorm Psychol* **101**, 668–674.
41. de Castro JM (1995) The relationship of cognitive restraint to the spontaneous food and fluid intake of free-living humans. *Physiol Behav* **57**, 287–295.
42. Tepper BJ, Trail AC & Shaffer SE (1996) Diet and physical activity in restrained eaters. *Appetite* **27**, 51–64.
43. Lowe MR & Levine AS (2005) Eating motives and the controversy over dieting: eating less than needed *versus* less than wanted. *Obes Res* **13**, 797–806.
44. Bathalon GP, Tucker KL, Hays NP, Vinken AG, Greenberg AS, McCrory MA & Roberts SB (2000) Psychological measures of eating behavior and the accuracy of 3 common dietary assessment methods in healthy postmenopausal women. *Am J Clin Nutr* **71**, 739–745.
45. Rizvi SL, Stice E & Agras WS (1999) Natural history of disordered eating attitudes and behaviors over a 6-year period. *Int J Eat Disord* **26**, 406–413.
46. Yao M & Roberts SB (2001) Dietary energy density and weight regulation. *Nutr Rev* **59**, 247–258.
47. Poppitt SD & Prentice AM (1996) Energy density and its role in the control of food intake: evidence from metabolic and community studies. *Appetite* **26**, 153–174.
48. Kant AK & Graubard BI (2005) Energy density of diets reported by American adults: association with food group intake, nutrient intake, and body weight. *Int J Obes Relat Metab Disord* **29**, 950–956.
49. McCullough ML, Feskanich D, Stampfer MJ, Giovannucci EL, Rimm EB, Hu FB, Spiegelman D, Hunter DJ, Colditz GA & Willett WC (2002) Diet quality and major chronic disease risk in men and women: moving toward improved dietary guidance. *Am J Clin Nutr* **76**, 1261–1271.

50. Laessle RG, Tuschl RJ, Kotthaus BC & Pirke KM (1989) Behavioral and biological correlates of dietary restraint in normal life. *Appetite* **12**, 83–94.
51. Dhingra R, Sullivan L, Jacques PF, Wang TJ, Fox CS, Meigs JB, Di'Agostino RB, Gaziano JM & Vasan RS (2007) Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation* **116**, 480–488.
52. de Castro JM (2004) The time of day of food intake influences overall intake in humans. *J Nutr* **134**, 104–111.
53. de Castro JM (1987) Circadian rhythms of the spontaneous meal pattern, macronutrient intake, and mood of humans. *Physiol Behav* **40**, 437–446.

## Appendix 1

### Food list

**Vegetables.** Cruciferous vegetables, orange vegetables, dark green vegetables, tomatoes, other vegetables, garlic, potatoes.

**Fruits.** Citrus fruits, other raw fruits, fruit sauce, dried fruits.

**Fruit juices.** 100 % fruit juices.

**Legumes.** Legumes, tofu.

**Nuts.** Peanut butter (hydrogenated or not), other nut butter, nuts and seeds.

**Whole-grain bread and cereals.** Bread, crackers, bagels, English muffins, pita bread, homemade muffins, pasta, brown rice, whole-grain couscous, breakfast cereals, hot cereals, granola bars, other (linseeds, wheat germ), croissants, waffles, pancakes.

**Refined bread and cereals.** Bread, crackers, bagels, English muffins, pita bread, homemade muffins, pasta, rice, couscous, breakfast cereals, hot cereals, granola bars, other (linseeds, wheat germ), croissants, waffles, pancakes.

**Skimmed and partly skimmed dairy products.** Milk (1 or 2 % fat), soya milk, sour cream (less than 5 % fat), cottage cheese (1 or 2 % fat), yoghurt (skimmed, 1 or 2 % fat), frozen milk or yoghurt (less than 5 % fat), milk dessert (skimmed, 1 or 2 % fat).

**Whole dairy products.** Milk (3.25 % fat), yoghurt (2 % fat or more), regular cottage or ricotta cheese, cheese, cream, sour cream (5 % fat or more), ice cream.

**Processed meat.** Delicatessen, sausages, headcheese (brawn), bacon.

**Red meat.** Beef, pork, veal, lamb, hamburgers, other red meat.

**Organ meat.** Liver, kidney, liver pâté, etc.

**Poultry.** Chicken, turkey, duck, other white meat not mentioned above.

**Fish and seafoods.** Fresh, frozen and canned fish, shrimps, scallops, etc.

**Other food categories.** Fries.

**Eggs.** Eggs, recipes (quiche, omelette).

**Saturated and trans fat.** Butter, lard, shortening, hydrogenated margarine.

**Unhydrogenated fat.** Unhydrogenated margarine, olive oil, rapeseed oil, other oils, olives, mayonnaise, salad dressing (regular or light).

**Fast food.** Poutine (French fries, cheese curds and gravy), chicken (nuggets or fried), fish (nuggets or fried), meat pie.

**Pizza.** All dressed, cheese, vegetarian.

**Snacks.** Popcorn, salted crackers, pretzels, chips (crisps).

**High-energy drinks.** Regular soft drinks, fruit punch.

**Low-energy drinks.** Diet soft drinks, other drinks low in energy (diet fruit drinks), excluding water.

**Alcohol.** Beer, wine, spirits.

**Other beverages.** Water, coffee, tea, herbal tea.

**Soup.** Soup, cream.

**Dessert, sweets.** Chocolate, candy, cookies, pie, cake, doughnuts, bars.

**Side dishes.** Sauce (ketchup, mustard, etc), jam, syrup, honey, sugar.

### Notes

For dairy products we have established that a portion was equivalent to one cup of milk or enriched soya beverages, 50 g cheese or 175 g yoghurt.

For red meat and processed meat, poultry, or fish, one portion was equivalent to 1 ounce (about 30 g).

For the meat substitutes component we have established that a portion was equivalent to half a cup of legumes, quarter a cup of nuts or seeds or 100 g tofu.

For grains products we have established that a portion was equivalent to one slice of bread, half a cup of pasta, rice or couscous, 30 g cereal.

For fruits and vegetables we have established that a portion was equivalent to half a cup or one medium fruit or vegetable (fruit and vegetable juices included).

For added fat a portion was equivalent to one teaspoon.

For sweets, we have established that a portion was equivalent, for example, to 1/12 of cake, 1/6 of pie or one regular chocolate bar.

For soft drinks a portion was equivalent to a cup.

For snacks a portion was equivalent to one cup of pop maize or ten saltine crackers, pretzels, chips (crisps) or nachos.