

Comparative Studies of the Dust around Red Supergiant and Oxygen-Rich Asymptotic Giant Branch Stars in the Local Universe

B. A. Sargent¹, S. Srinivasan², A. Speck³, K. Volk⁴, F. Kemper², W. Reach⁵, E. Lagarde⁶, J.-P. Bernard^{7,8}, I. McDonald⁹, M. Meixner⁴, G. C. Sloan¹⁰ and O. Jones⁴

¹Center for Imaging Science and Laboratory for Multiwavelength Astrophysics, Rochester Institute of Technology, 54 Lomb Memorial Drive, 17-3195, Rochester, NY 14623, USA, email: baspci@rit.edu

²Academia Sinica Institute of Astronomy and Astrophysics, P.O. Box 23-141, Taipei 10617, Taiwan

³Department of Physics and Astronomy, University of Missouri, Columbia, MO 65211, USA

⁴Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218, USA

⁵Universities Space Research Association, MS 232-11, Moffett Field, CA 94035, USA

⁶Laboratoire Lagrange, UMR7293, Université Côte d'Azur, CNRS, Observatoire de la Côte d'Azur, Boulevard de l'Observatoire, 06304 Nice Cedex 4, France

⁷CNRS, IRAP, 9 Av. Colonel Roche, BP 44346, F-31028 Toulouse Cedex 4, France

⁸Université de Toulouse, UPS-OMP, IRAP, F-31028 Toulouse Cedex 4, France

⁹Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, Alan Turing Building, Oxford Road, Manchester M13 9PL, UK

¹⁰Cornell Center for Astrophysics and Planetary Science, Cornell University, Ithaca, NY 14853, USA

Abstract. We analyze the dust emission features seen in *Spitzer Space Telescope* Infrared Spectrograph (IRS) spectra of red supergiant (RSG) and oxygen-rich asymptotic giant branch (AGB) stars in the Large Magellanic Cloud and Small Magellanic Cloud galaxies and in various Milky Way globular clusters. The spectra come from the *Spitzer* Legacy program SAGE-Spectroscopy (PI: F. Kemper), the *Spitzer* program SMC-Spec (PI: G. Sloan), and other archival *Spitzer*-IRS programs. The broad 10 and 20 micron emission features attributed to amorphous dust of silicate composition seen in the spectra show evidence for systematic differences in the centroid of both emission features between O-rich AGB and RSG populations. Radiative transfer modeling using the GRAMS grid of models of AGB and RSG stars suggests that the centroid differences are due to differences in dust properties. We investigate differences in dust composition, size, shape, etc that might be responsible for these spectral differences. We explore how these differences may arise from the different circumstellar environments around RSG and O-rich AGB stars and assess effects of varying metallicity (LMC versus SMC versus Milky Way globular cluster) and other properties (mass-loss rate, luminosity, etc.) on the dust originating from these stars. BAS acknowledges funding from NASA ADAP grant NNX13AD54G.

Keywords. stars: AGB and post-AGB, (stars:) circumstellar matter




Figure 1. Shown is the dust model of IRAS 04544-6849, an oxygen-rich (O-rich) AGB star in the LMC, according to Jones *et al.* (2012). The data is from pid 3583 (PI: T. Onaka), AOR #11219968, with the spectrum reduced according to the methods described by Kemper *et al.* (2010). An O-rich GRAMS model from Sargent *et al.* (2011) has been fit to its optical/infrared spectral energy distribution, the stellar photosphere has been subtracted from the *Spitzer* Infrared Spectrograph spectrum, and the residuals are the black points in this plot being fit by the model (solid orange line). The model is of similar style to those of Sargent *et al.* (2009), except the components with size given in the legend are for spherical dust grains whose optical constants are those of the glassy silicates of cosmic composition from Speck *et al.* (2015).

References

- Jones, O. *et al.* 2012, *MNRAS*, 427, 3209
 Kemper, F. *et al.* 2010, *PASP*, 122, 683
 Sargent, B. A. *et al.* 2009, *ApJ*, 690, 1193
 Sargent, B. A., Srinivasan, S., & Meixner, M. 2011, *ApJ*, 728, 93
 Speck, A. K., Pitman, K. M., & Hofmeister, A. M. 2015, *ApJ*, 809, 65