

## Suicidology and suicide prevention

### EPP1389

#### Managing ethical issues in community-based adolescent self-harm research. ethical practice in an adolescent self-harm preventative intervention research project with pupils in secondary schools in wales.

R. Parker\*

Decipher, Cardiff University, Cardiff, United Kingdom

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1547

**Introduction:** Adolescent self-harm in Europe is a major public health challenge and shares a risk continuum with suicide. Recent research demonstrates a strong risk correlation with attempting suicide in the community-based adolescent self-harm population group, similar to research findings for populations in health setting-based research. In the UK increasing rates in hospital admissions represent the “tip of the iceberg” within the community, with potentially two thirds of the population group not accessing health services for support. This brings many health risks, including an increased suicide risk. Finding a solution to these issues requires a preventative intervention approach for young people, including community-based delivery to address service access barriers. Secondary schools are posited as key settings where this type of support could be delivered. But emerging UK research demonstrates challenges in completing adolescent self-harm research with young people in schools.

**Objectives:** This paper centres on a current adolescent self-harm preventative intervention research project in Cardiff University. It focuses on ethical research practice in community-based adolescent self-harm research.

**Methods:** This paper appraises some of the core ethical issues, challenges and their management in completing adolescent self-harm prevention intervention research in secondary school settings in Wales. It also provides an overview of the project’s innovative safety protocol design.

**Results:** This project was successful in managing the potential risks to the school-based adolescent research participants.

**Conclusions:** This work helps address some of the current research barriers to completing adolescent self-harm prevention intervention research in schools, to facilitate shared solutions to the urgent public health challenge of adolescent self-harm.

**Keywords:** adolescent self-harm; preventative intervention research; secondary schools; ethics in research practice

### EPP1388

#### The relationship between physical pain and suicidal thoughts and behaviors in adolescents: A meta-analysis

M. De Filippi<sup>1</sup>, M. Riganese<sup>1</sup>, E. Salmè<sup>1</sup>, F. Madeddu<sup>1</sup> and R. Calati<sup>1,2\*</sup>

<sup>1</sup>Psychology, University of Milano-Bicocca, Milan, Italy and <sup>2</sup>Adult Psychiatry, Nîmes University Hospital, Nîmes, France

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1548

**Introduction:** Suicide is the third leading cause of death in adolescents (15-19 years). Physical pain is an important risk factor for suicidal thoughts and behaviors, especially in a delicate phase as adolescence.

**Objectives:** Several studies investigated this association and the aim of this meta-analysis was to synthesize data in literature about this topic in adolescents (11-20 years).

**Methods:** We started from a systematic review published by Hinze and colleagues in 2019 and we searched on PubMed (January 2019-June 2020) studies comparing rates of suicidal outcomes (Suicidal Ideation, Suicide Plan, Attempt, Self-Harm: SI, SP, SA, SH) in individuals with any type of physical pain (head, back, neck, chest, stomach, abdomen, muscle, joint, arthritis) vs. those without it. Data were analyzed with Comprehensive Meta-Analysis software (CMA, version 3).

**Results:** Of the 16 included studies, eleven focused on SI (68.8%), six (37.5%) on SA, four (25%) on SH and two (12.5%) on SP. Adolescents with physical pain were more likely to report SI ( $p < .001$ ), SH ( $p < .001$ ), SA ( $p = .004$ ) and SP ( $p = .006$ ). In all analyses, the between study heterogeneity was high. The presence of publication bias has been detected in SI ( $k \geq 10$ ).

**Conclusions:** Results are in line with previous literature on this topic. Future research should investigate the specific impact of: acute vs. chronic pain; different types and intensities of pain; planned vs. impulsive action and therefore suicidal intent; role of psychological factors (in particular sensitivity and tolerance to physical pain).

**Keywords:** Suicide; Suicidal Thoughts and Behaviors; adolescents; Physical Pain

### EPP1389

#### Physical pain – suicidality association in adults: A meta-analysis

M. Riganese<sup>1</sup>, M. De Filippi<sup>1</sup>, E. Salmè<sup>1</sup>, F. Madeddu<sup>1</sup> and R. Calati<sup>1,2\*</sup>

<sup>1</sup>Psychology, University of Milano-Bicocca, Milan, Italy and

<sup>2</sup>Psychiatry, Nîmes University Hospital, Nîmes, France

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1549

**Introduction:** Multiple epidemiologic and clinical studies have explored the relationship between physical pain and suicidal thoughts and behaviours.

**Objectives:** The aim of this meta-analysis was to provide an update of the data already present in literature about this specific association in adults.

**Methods:** Starting from a meta-analysis published by Calati and colleagues in 2015, 28 studies were included in this work. After searching on Pubmed (until March 2020), data were extracted from articles comparing the rates of current and lifetime suicidal thoughts and behaviours (death wish, suicidal ideation, suicidal planning, suicide attempt and suicide death: DW, SI, SP, SA, and SD) in adults with any type of physical pain and in individuals who did not report this condition. Data were analysed using Cochrane Collaboration Review Manager software (RevMan, version 5.4).

**Results:** Although high between-study heterogeneity was detected in most analyses, results suggested that adults with physical pain are more likely to report any form of suicidal outcome, except for death by suicide, compared to individuals not affected by pain. No