

ON THE THETA DIVISOR OF $SU(r, 1)$

SONIA BRIVIO AND ALESSANDRO VERRA

Abstract. Let $SU(r, 1)$ be the moduli space of stable vector bundles, on a smooth curve C of genus $g \geq 2$, with rank $r \geq 3$ and determinant $\mathcal{O}_C(p)$, $p \in C$; let \mathcal{L} be the generalized theta divisor on $SU(r, 1)$. In this paper we prove that the map $\phi_{\mathcal{L}}$, defined by \mathcal{L} , is a morphism and has degree 1.

§0. Introduction

Let C be a smooth, irreducible, complex, projective curve, of genus $g \geq 2$. Let $SU(r, d)$ denotes the moduli space of semistable vector bundles with rank r and fixed determinant $L \in \text{Pic}^d(C)$. $SU(r, d)$ is an irreducible projective variety of dimension $(r^2 - 1)(g - 1)$, (see [S] and [N-R]), its Picard variety is free cyclic, see [D-N], the ample generator \mathcal{L} is called the generalized theta divisor of $SU(r, d)$. Let $\phi_{\mathcal{L}}: SU(r, d) \rightarrow |\mathcal{L}|^*$ be the map associated to the theta divisor: if $r = 2$, then $\phi_{\mathcal{L}}$ is an embedding, see [Be1], [L],[B-V1], [vG-I] for d even, [D-R], [Be2] and [B-V2] for d odd. In this paper, we will assume $r \geq 3$ and we will consider $SU(r, 1)$, where $L = \mathcal{O}_C(p)$ and p is a given point of C , our first result is the following:

THEOREM 0.0.1. *For any curve C of genus $g \geq 2$: $\deg(\phi_{\mathcal{L}}) = 1$, the linear system $|\mathcal{L}|$ on $SU(r, 1)$ is base points free, i.e. the map $\phi_{\mathcal{L}}$ is a morphism.*

As a second result we prove the following:

THEOREM 0.0.2. *For any curve C of genus $g \geq 2$, we have $\deg(\phi_{\mathcal{L}}) = 1$.*

The paper is organized as follows. The first section is devoted to proving theorem (0.0.1). In section 2, we study rank r -bundles with $r + 1$

Received December 7, 1999.

2000 Mathematics Subject Classification: 14 D20.

The authors were partially supported by the European Science Project "AGE (Algebraic Geometry in Europe)", contract n. ERBCHRXCT940557.

sections extending the rank 2 case dealt in [B-V2]. Fix a line bundle $l \in \text{Pic}^g(C)$: we can identify the spaces $SU(r, 1)$ and $SU(r, O_C(p + rl))$, let $E \in SU(r, O_C(p + rl))$, assume that $h^0(E) = r + 1$ and the natural map $w_E: \wedge^r H^0(E) \rightarrow H^0(\det E)$ is injective, then $\text{Im } w_E$ is a $(r + 1)$ -dimensional subspace of $H^0(O_C(p + rl))$. This allow us to define a map

$$g_l: SU(r, O_C(p + rl)) \rightarrow G_l(r + 1, H^0(O_C(p + rl))),$$

we prove that g_l is a birational map and it is defined by a linear system in $|\mathcal{L}|$. In section 3, we prove theorem (0.0.2). Actually, we perform a non empty open subset $\mathcal{U} \subset SU(r, 1)$ such that the restriction $\phi_{\mathcal{L}|_{\mathcal{U}}}$ is an embedding. \mathcal{U} is naturally defined as the set of bundles ξ for which exists $l \in \text{Pic}^g(C)$, s.t. g_l is biregular at the point $E = \xi(l)$. If $r = 2$, in [B-V2] we proved that actually $\mathcal{U} = SU(2, 1)$, which allows us to conclude that \mathcal{L} is very ample. If $r \geq 3$, actually \mathcal{U} can be a proper subset of $SU(r, 1)$, (see lemma (3.2.1)), this unable us to extend completely the result of rank 2.

Finally, we would like to remember that rank 2 vector bundles with 3 sections were useful also in proving that $\phi_{\mathcal{L}}$ is an embedding at singular points of $SU(2)$, see [I-vG].

0.1. Notations.

We reserve the notation ξ for points of $SU(r, 1)$; with some abuse, the same notation will be used for the vector bundle corresponding to ξ . For a vector bundle ξ of degree d and rank r we denote by $\mu := \frac{d}{r}$ the slope of ξ . We say that ξ is semistable iff for every proper subbundle $\eta \subset \xi$ we have $\mu(\eta) \leq \mu(\xi)$, it is stable iff the inequality is strict. Given two vector bundles ξ, η on C , they are said complementary if $\chi(\xi \otimes \eta) = 0$.

We recall that there exists a Poincaré family on $SU(r, 1)$, see [N-R], i.e. a vector bundle U on $SU(r, 1) \times C$ such that $U|_{\xi \times C} \simeq \xi$, for any $\xi \in SU(r, 1)$. Let as usual π_i denote the natural projections of $SU(r, 1) \times C$ onto factors. Note that if U is a Poincaré bundle, then for any $A \in \text{Pic}(SU(r, 1))$, $U \otimes \pi_1^* A$ is a Poincaré bundle too. Actually there exists a unique Poincaré bundle U on $SU(r, 1) \times C$ with the further following property, (see [Ra]):

$$\det U|_{SU(r, 1) \times \{x\}} \simeq \mathcal{L},$$

where \mathcal{L} is the theta divisor of $SU(r, 1)$. Following [Ra], we will call such a bundle U the universal bundle.

§1. On the base points of the theta divisor

1.0.

Let θ be an effective divisor of degree $g - 1$ on C , θ defines a natural isomorphism

$$(1) \quad f_\theta: SU(r, 1) \rightarrow SU(r, r(g - 1) - 1)$$

sending ξ to $\xi^*(\theta)$. Let $(\xi, \eta) \in SU(r, 1) \times SU(r, r(g - 1) - 1)$ we have

$$(2) \quad \chi(\xi \otimes \eta) = 0,$$

hence the subset

$$(3) \quad \hat{\Theta}_\xi := \{\eta \in SU(r, r(g - 1) - 1) \mid h^0(\xi \otimes \eta) > 0\}$$

is either $SU(r, r(g - 1) - 1)$ or a theta divisor of $SU(r, r(g - 1) - 1)$, see [D-N].

LEMMA 1.0.1. *Let $U_\xi \subset \hat{\Theta}_\xi$ be the locus of points η such that each non zero morphism $u: \eta^* \rightarrow \xi$ is a monomorphism. Then U_ξ is a non empty open subset.*

Proof. Let \mathcal{F} be a family of stable vector bundles on $S \times C$, let $U: \mathcal{F}^* \rightarrow \pi_2^*\xi$ be a non zero morphism of vector bundles. It is enough to show that the locus Δ of points $s \in S$ such that U_s is not a monomorphism is closed. This is immediate because Δ is the projection of the degeneracy locus of U . The non emptiness follows from the exact sequence

$$(4) \quad 0 \rightarrow \xi(-\theta) \rightarrow \xi \rightarrow O_\theta \otimes \xi \rightarrow 0$$

where $\eta^* = \xi(-\theta)$. □

LEMMA 1.0.2. *We have: $\dim U_\xi \leq (r^2 - 1)(g - 1) - 1$.*

Proof. Let $\eta \in U_\xi$: then there exists an exact sequence as follows

$$(5) \quad 0 \rightarrow \eta^* \rightarrow \xi \rightarrow O_D \rightarrow 0,$$

where D is a divisor in the linear system $|\det(\xi) \otimes \det(\eta)|$, that is $|r\theta|$. Let's consider the natural rational map

$$(6) \quad V_D: \text{Hom}(\xi, O_D) \rightarrow SU(r, r(g - 1) - 1)$$

which associates to any epimorphism $v: \xi \rightarrow O_D$ the sheaf $(\ker v)^*$. Then any $\eta \in U_\xi$ belongs to the image of the above map V_D . The group of automorphisms $\text{Aut}(O_D)$ naturally acts on $\text{Hom}(\xi, O_D)$; the action is faithful on points v which are epimorphisms, moreover $\text{Aut}(O_D)$ contains the torus $\mathbf{C}^{*\text{deg } D}$. Since $\text{Hom}(\xi, O_D) \simeq \xi^* \otimes O_D$, it follows that the dimension of the image of the previous map at V_D is bounded by

$$(r - 1) \text{deg } D = (r^2 - 1)(g - 1) - (r - 1)(g - 1).$$

On the other hand, if $r \geq 3$ we have $\dim |D| = (r - 1)(g - 1) - 1$. Let $\mathcal{D} \subset C \times |D|$ be the universal divisor, then U_ξ is contained in the image of the natural map

$$(7) \quad \pi_{2*}((\pi_1^* \xi) \otimes O_{\mathcal{D}}) \rightarrow SU(r, r(g - 1) - 1).$$

By the previous count the dimension of this image is at most $(r^2 - 1)(g - 1) - 1$. □

Let $\xi \in SU(r, 1)$, with the above notations, we define

$$(8) \quad \Theta_\xi := f_\theta^* \hat{\Theta}_\xi.$$

PROPOSITION 1.0.1. *For any $\xi \in SU(r, 1)$, Θ_ξ is actually a theta divisor on $SU(r, 1)$.*

Proof. It is enough to prove the assertion for $\hat{\Theta}_\xi$. Note that $\hat{\Theta}_\xi$ is either a theta divisor on $SU(r, r(g - 1) - 1)$ or it is $SU(r, r(g - 1) - 1)$. Let's assume that $\hat{\Theta}_\xi = SU(r, r(g - 1) - 1)$, then by the preceding lemma U_ξ is an open subset of $SU(r, r(g - 1) - 1)$: but this is impossible because $\dim U_\xi < \dim SU(r, r(g - 1) - 1)$. This implies the claim. □

1.1.

Fix $\theta \in C^{(g-1)}$, the previous remarks allow us to define a map

$$(9) \quad \phi_\theta: SU(r, 1) \rightarrow |\mathcal{L}| \simeq \mathbf{P}^n$$

just sending ξ to the divisor Θ_ξ . From the previous proposition follows immediately that ϕ_θ is a morphism.

LEMMA 1.1.1. *We have: $\phi_\theta^* O_{\mathbf{P}^n}(1) = \mathcal{L}$.*

Proof. Let $\xi_0 \in SU(r, 1)$ be a point which is not a point of ramification of ϕ_θ , set $\eta := f_\theta(\xi_0) = \xi_0^*(\theta)$. Let H be the hyperplane of \mathbf{P}^n consisting of divisors passing through ξ_0 . We have:

$$\phi_\theta^* H = \{\xi \in SU(r, 1) : \eta \in \hat{\Theta}_\xi\} = \{\xi \in SU(r, 1) : h^0(\xi \otimes \eta) > 0\};$$

and set theoretically this is a divisor in the linear system $|\mathcal{L}|$. Actually, $\xi_0 \in \phi_\theta^* H$ and since it is not a point of ramification, we have $\phi_\theta^* H = \mathcal{L}$. □

As an immediate consequence we have

PROPOSITION 1.1.1. *The map associated to the theta divisor $\phi_{\mathcal{L}}: SU(r, 1) \rightarrow |\mathcal{L}|^*$ is a morphism.*

§2. Bundles with $r + 1$ sections

2.1. Definition

Let $(\xi, l) \in SU(r, 1) \times \text{Pic}^g(C)$. We say that (ξ, l) satisfies condition (*) if the following three properties hold:

- (i) $h^0(\xi(l)) = r + 1$,
- (ii) $\xi(l)$ is globally generated,
- (iii) the determinant map $w_{\xi,l} : \wedge^r H^0(\xi(l)) \rightarrow H^0(\det \xi(l))$ is injective.

We will set

$$(10) \quad X_l := \{\xi \in SU(r, 1) / (\xi, l) \text{ satisfies } (*)\}.$$

2.2. Remark

Assume that a pair $(\xi, l) \in SU(r, 1) \times \text{Pic}^g(C)$ satisfies properties (i) and (ii), then it satisfies (iii) too. First of all, note that since $h^0(\xi(l)) = r + 1$, every vector of $\wedge^r H^0(\xi(l))$ is indecomposable. So assume that $v \neq 0$ is in the kernel of the map $w_{\xi,l}$, then $v = s_1 \wedge s_2 \wedge \dots \wedge s_r$, with $s_i \in H^0(\xi(l))$, $i = 1 \dots r$. Then the sections s_1, \dots, s_r would generate a subbundle $\eta \subset \xi(l)$ with the following properties: $rk\eta = s \leq r - 1$, $h^0(\eta) \geq r$, and η is globally generated too. This implies $rk\eta = r - 1$, $h^0(\eta) = r$ and the following commutative diagram

$$(11) \quad \begin{array}{ccccccc} 0 & \longrightarrow & (\det \eta)^{-1} & \longrightarrow & H^0(\eta) \otimes \mathcal{O}_C & \xrightarrow{e} & \eta & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{O}_C(p + rl)^{-1} & \longrightarrow & H^0(\xi(l)) \otimes \mathcal{O}_C & \longrightarrow & \xi(l) & \longrightarrow & 0, \end{array}$$

which implies $\text{deg } \eta \geq 1 + rg$, which contradicts the stability of $\xi(l)$.

We will show later that actually for any $l \in \text{Pic}^g(C)$, X_l is a non empty open subset of $SU(r, 1)$.

2.3. Definition

Let $l \in \text{Pic}^g(C)$, we will consider the Grassmannian

$$(12) \quad G_l = G(r + 1, H^0(O_C(p + rl)))$$

of $(r + 1)$ dimensional subspaces of $H^0(O_C(p + rl))$. If (ξ, l) satisfies assumptions (i) and (iii) then the image of the determinant map $w_{\xi,l}$ is a $(r + 1)$ dimensional subspace of $H^0(O_C(p + rl))$, let's denote it by

$$W := \text{Im } w_{\xi,l}.$$

This defines a map

$$(13) \quad g_l: SU(r, 1) \rightarrow G_l(r + 1, H^0(O_C(p + rl)))$$

by sending ξ to the point of the Grassmannian corresponding to the subspace $W \hookrightarrow H^0(O_C(p + rl))$.

Note that there is a canonical isomorphism $\wedge^r H^0(\xi(l)) \simeq H^0(\xi(l))^*$, which induces an inclusion

$$(14) \quad w'_\xi: H^0(\xi(l))^* \hookrightarrow H^0(O_C(p + rl)),$$

whose image is again W . Assume now that $\xi(l)$ is globally generated too, then we have an exact sequence

$$(15) \quad 0 \rightarrow O_C(p + rl)^{-1} \rightarrow H^0(\xi(l)) \otimes O_C \rightarrow \xi(l) \rightarrow 0,$$

and its dual

$$(16) \quad 0 \rightarrow \xi(l)^* \rightarrow H^0(\xi(l))^* \otimes O_C \rightarrow O_C(p + rl) \rightarrow 0;$$

passing to cohomology we have

$$(17) \quad 0 \rightarrow H^0(\xi(l)^*) \rightarrow H^0(\xi(l))^* \xrightarrow{\pi} H^0(O_C(p + rl)) \rightarrow \dots,$$

since $\xi(l)$ is stable, then $H^0(\xi(l)^*) = 0$, so we can conclude that π is injective. We claim that $\text{Im } \pi = W$, so that we can identify the maps π and w'_ξ .

2.4.

Let $W \in G_l(r + 1, H^0(O_C(p + rl)))$, assume that $|W|$ is base point free. Then we can consider the evaluation map $e: W \otimes O_C \rightarrow O_C(p + rl)$, which is surjective, so its kernel is a rank r vector bundle, let's define

$$(18) \quad E_W := (\text{Ker } e)^*.$$

We have $\det E_W = O_C(p + rl)$, moreover we have the following exact sequence

$$(19) \quad 0 \rightarrow O_C(p + rl)^{-1} \rightarrow W^* \otimes O_C \rightarrow E_W \rightarrow 0,$$

so we can conclude that E_W is generated by $(r + 1)$ global sections spanning the subspace $\text{Im}(W^* \hookrightarrow H^0(E_W))$. Passing to cohomology, we have

$$(20) \quad 0 \rightarrow W^* \rightarrow H^0(E_W) \rightarrow H^1(O_C(p + rl)^{-1}) \rightarrow W^* \otimes H^1(O_C) \rightarrow \dots;$$

note that $h^0(E_W) = r + 1$ if and only $H^0(E_W) \simeq W^*$, that is the following multiplication map is an isomorphism

$$(21) \quad \mu_W: W \otimes H^0(\omega_C) \rightarrow H^0(\omega_C \otimes O_C(p + rl)).$$

We have the following results:

LEMMA 2.4.1. *Let E be a rank r vector bundle with $h^0(E) = \chi(E) = r + 1$, which is globally generated, then E is stable.*

Proof. By Riemann Roch theorem we have $\deg(E) = 1 + rg$, and $\mu(E) = g + \frac{1}{r}$. Assume there exists a destabilizing subbundle $F \subset E$ with $rk(F) = s \leq r - 1$ and $\mu(F) \geq g + \frac{1}{r}$. This implies $\deg(F) \geq 1 + sg$ and $\chi(F) \geq s + 1$. Since E is generated by $r + 1$ global sections spanning $H^0(E)$, then $h^0(F) = s + 1$ and F is globally generated too. So we have a commutative diagramm

$$(22) \quad \begin{array}{ccccccc} 0 & \longrightarrow & (\det F)^{-1} & \longrightarrow & H^0(F) \otimes O_C & \xrightarrow{e} & F \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \det E^{-1} & \longrightarrow & H^0(E) \otimes O_C & \xrightarrow{e} & E \longrightarrow 0, \end{array}$$

from the inclusion $(\det F)^{-1} \hookrightarrow \det E^{-1}$ we have $sg + 1 \geq rg + 1$, which is impossible. This concludes the proof.

Assume that for a subspace W the map μ_W is an isomorphism, then by preceding lemma E_W is stable, so that $E_W(-l) = \xi \in SU(r, 1)$. Moreover, (ξ, l) satisfies conditions (i),(ii) so that the map g_l is defined at the point ξ and we actually have $g_l(\xi) = W$. We would remark that the exact sequence

$$(23) \quad 0 \rightarrow \xi(l)^* \rightarrow W \otimes O_C \rightarrow O_C(p + rl) \rightarrow 0$$

is just the pull-back of the Euler sequence

$$(24) \quad 0 \rightarrow \Omega_{\mathbf{P}^{r-1}}(1) \rightarrow W \otimes O_{\mathbf{P}^{r-1}} \rightarrow O_{\mathbf{P}^{r-1}}(1) \rightarrow 0$$

under the morphism $f: C \rightarrow \mathbf{P}^{r-1} = \mathbf{P}(W^*)$ defined by $|W|$. Hence it turns out that

$$(25) \quad \xi(l) \simeq f^*T_{\mathbf{P}^{r-1}}(-1).$$

Let's define the following subsets of G_l :

$$(26) \quad B_l = \{W \in G_l: |W| \text{ has base points} \}$$

and D_l as the set of W such that the multiplication map

$$(27) \quad \mu_W: W \otimes H^0(\omega_C) \rightarrow H^0(\omega_C \otimes O_C(p + rl))$$

is not surjective. Note that $\forall l$, we have $B_l \subset D_l$. Moreover, we have the following fact:

LEMMA 2.4.2. *For any $l \in \text{Pic}^g(C)$, D_l is a Cartier divisor on G_l .*

Proof. For more details see also [B], th.(0.0.1).

There exists a homomorphism between vector bundles $\mu: \mathcal{G} \rightarrow \mathcal{F}$ such that at the point $W \in G_l$ is actually the multiplication map

$$(28) \quad \mu_W: W \otimes H^0(\omega_C) \rightarrow H^0(\omega_C \otimes O_C(p + rl));$$

so that D_l is actually the degeneracy locus of μ . From Thom-Porteous's formula it is either a Cartier divisor or $D_l = G_l$. Actually we show that there exists $W \notin D_l$.

Claim: let $r \geq 1$, for any line bundle L of degree $rg + 1$, base points free and non special, there exists a subspace $W \subset H^0(L)$ s.t. μ_W is surjective.

We will prove the claim by recurrence on r . Note that if $r = 1$, and L is a non special base points free line bundle of degree $g + 1$, the assertion

follows from the base points free pencil trick, see [A-C-G-H]. Assume that the claim is true for degree $1 + (r - 1)g$. Let L be any line bundle of degree $1 + rg$: choose $x_1, \dots, x_g \in C$ with the following properties:

- i) $x_1 + \dots + x_g$ is non special,
- ii) $L(-x_1 - \dots - x_g)$ is base points free and non special.

By the induction hypothesis there exists an r dimensional subspace $\bar{W} \subset H^0(L(-x_1 - \dots - x_g))$ for which $\mu_{\bar{W}}$ is surjective. Choose a global section $s \in H^0(L)$, such that $s(x_i) \neq 0$, for $i = 1, \dots, g$, and define the subspace

$$W := \langle \bar{W}, s \rangle.$$

By i), we can find independent global sections $\omega_1, \omega_2, \dots, \omega_g$ such that $\omega_i(x_j) \neq 0$ if and only if $j = i$; let $f_i = \mu_W(s \otimes \omega_i)$, then it is easy to see that f_1, \dots, f_g are independent global sections of $H^0(L \otimes \omega_C)$. This implies the following commutative diagram

$$(29) \quad \begin{array}{ccc} \bar{W} \otimes H^0(\omega_C) & \xrightarrow{\mu_{\bar{W}}} & H^0(L(-x_1 - \dots - x_g) \otimes \omega_C) \\ \downarrow & & \downarrow \\ W \otimes H^0(\omega_C) & \xrightarrow{\mu_W} & H^0(L \otimes \omega_C) \\ \downarrow & & \downarrow \\ \langle s \rangle \otimes H^0(\omega_C) & \xrightarrow{\mu_s} & \langle f_1, \dots, f_g \rangle \end{array}$$

Since both $\mu_{\bar{W}}$ and μ_s are surjective, we can conclude that μ_W is surjective too.

As an immediate consequence of the lemma we have that X_l is a non empty open subset of $SU(r, 1)$: in fact if W is a point in $G_l - D_l$, then by the previous arguments $E_W(-l) = \xi \in X_l$. Moreover the map $h_l: G_l - D_l \rightarrow X_l$ sending W to $E_W(-l)$ is actually the inverse map of g_l .

2.5.

Let's consider the Pluecker embedding of the grassmannian G_l :

$$(30) \quad p_l: G_l(r + 1, H^0(O_C(p + rl))) \hookrightarrow \mathbf{P}^N = \mathbf{P}(\wedge^{r+1} H^0(O_C(p + rl)))$$

and look at the composition map

$$(31) \quad p_l \cdot g_l: SU(r, 1) \rightarrow \mathbf{P}^N,$$

we have the following result:

PROPOSITION 2.5.1. *Let $l \in \text{Pic}^g(C)$,*

- (1) $g_l: SU(r, 1) \rightarrow G_l(r + 1, H^0(O_C(p + rl)))$ *is a birational map, the restriction $g_l|_{X_l}: X_l \rightarrow G_l - D_l$ is biregular;*
- (2) *the rational map $p_l \cdot g_l: SU(r, 1) \rightarrow \mathbf{P}^N$ is defined by $N + 1$ independent global sections of $H^0(\mathcal{L})$, where \mathcal{L} is the generalized theta divisor on $SU(r, 1)$.*

Proof. (1) Let $l \in \text{Pic}^g(C)$, note that we can identify the two moduli spaces $SU(r, 1)$ and $SU(r, O_C(p + rl))$ via the natural isomorphism sending $\xi \rightarrow \xi(l)$. Let \mathcal{U}_l be the universal bundle on $SU(r, O_C(p + rl)) \times C$, let as usual π_i , with $i = 1, 2$, denote the natural projections. We recall that

$$(32) \quad \det \mathcal{U}_l|_{SU(r, O_C(p+rl)) \times x} \simeq \mathcal{L},$$

moreover $\det \mathcal{U}_l|_{\xi(l) \times C} \simeq O_C(p + rl)$, so that we can conclude that

$$(33) \quad \det \mathcal{U}_l \simeq \pi_2^* O_C(p + rl) \otimes \pi_1^* \mathcal{L}.$$

We will consider, on $SU(r, O_C(p + rl))$, the torsion free sheaf $\pi_{1*} \mathcal{U}_l$, whose fibre at the point ξ is $H^0(\xi(l))$. Let's consider the following open subset of $SU(r, O_C(p + rl))$

$$(34) \quad V_l := \{\xi(l): h^0(\xi(l)) = r + 1\},$$

then $\pi_{1*} \mathcal{U}_l|_{V_l}$ is a vector bundle of rank $r + 1$. There is a natural map between sheaves on $SU(r, O_C(p + rl)) \times C$, see [H],

$$(35) \quad E: \pi_1^*(\pi_{1*} \mathcal{U}_l) \rightarrow \mathcal{U}_l,$$

let's consider the map $\wedge^r E$

$$(36) \quad \wedge^r E: \pi_1^*(\wedge^r \pi_{1*} \mathcal{U}_l) \rightarrow \wedge^r \mathcal{U}_l = \det \mathcal{U}_l,$$

and tensor this map with $\pi_1^* \mathcal{L}^{-1}$, so we have

$$(37) \quad \pi_1^*(\wedge^r(\pi_{1*} \mathcal{U}_l) \otimes \mathcal{L}^{-1}) \rightarrow \pi_2^* O_C(p + rl).$$

Finally let's push down this map on $SU(r, O_C(p + rl))$, by using the projecting formula and recalling that $\pi_{1*} O_{SU(r, O_C(p+rl)) \times C} \simeq O_{SU(r, O_C(p+rl))}$, we will have the following map

$$(38) \quad G: \wedge^r(\pi_{1*} \mathcal{U}_l) \otimes \mathcal{L}^{-1} \rightarrow \pi_{1*} \pi_2^* O_C(p + rl).$$

Note that $\pi_{1*}\pi_2^*O_C(p + rl)$ is the trivial bundle on $SU(r, O_C(p + rl))$ with fibre $H^0(O_C(p + rl))$, moreover at the point $\xi(l)$ G is actually the determinant map

$$(39) \quad w_\xi: \wedge^r H^0(\xi(l)) \rightarrow H^0(O_C(p + rl)).$$

If $g_l(\xi)$ is defined, then $(\text{Im } G)_{\xi(l)} = g_l(\xi)$ and this shows that g_l is a rational map. Moreover, let $U_l \subset SU(r, O_C(p + rl))$ the set of points $\xi(l)$ satisfying properties (i) and (iii), then $X_l \subset U_l$ and the restriction $G|_{U_l}$ is an injection of vector bundle, and $\text{codim } U_l \geq 2$.

Since $\dim SU(r, 1) = \dim G_l = (r^2 - 1)(g - 1)$, and moreover both $SU(r, 1)$ and G_l are smooth and irreducible, then by Zariski's main theorem it is enough to show that $g_l|_{X_l}$ is injective, but this follows from the preceding section.

(2) Since $\wedge^r(\pi_{1*}\mathcal{U}_l) \simeq \pi_{1*}\mathcal{U}_l^* \otimes \det(\pi_{1*}\mathcal{U}_l)$, $G|_{U_l}$ gives the following injection

$$(40) \quad (\pi_{1*}\mathcal{U}_l)^* \otimes \det(\pi_{1*}\mathcal{U}_l) \otimes \mathcal{L}^{-1} \hookrightarrow H^0(O_C(p + rl)) \otimes O_{SU(r,1)},$$

which is actually the pull back of the universal subbundle \mathcal{W} on G_l , via the map $g_l|_{U_l}$. Since the Pluecker map p_l of G_l is defined by the line bundle $\det W^*$, we can conclude that

$$(41) \quad (p_l \cdot g_l)^*(O_{\mathbf{P}^N(1)}) \simeq \det(g_l^*W^*).$$

We will prove that actually $g_l^*W^* \simeq \pi_{1*}\mathcal{U}_l$ and $\det \pi_{1*}\mathcal{U}_l = \mathcal{L}$.

Let's consider again the natural map of sheaves

$$(42) \quad E: \pi_1^*(\pi_{1*}\mathcal{U}_l) \rightarrow \mathcal{U}_l,$$

the restriction at $\xi(l) \times C$ is actually the evaluation map: assume that $\xi(l) \in X_l$, then $E|_{\xi(l) \times C}$ is surjective and $(\ker E)|_{\xi(l) \times C} \simeq O_C(p + rl)^{-1}$. Let's consider the set $V \subset SU(r, O_C(p + rl)) \times C$ of pairs $(\xi(l), x)$ with $\xi(l) \in X_l$: we have

$$(43) \quad (\text{Ker } E)|_V = \pi_2^*O_C(p + rl)^{-1} \otimes \pi_1^*B,$$

with $B \in \text{Pic}(SU(r, O_C(p + rl)))$. Look at the following exact sequences on V :

$$(44) \quad 0 \rightarrow \ker E|_V \rightarrow \pi_1^*(\pi_{1*}\mathcal{U}_l)|_V \rightarrow \mathcal{U}_l|_V \rightarrow 0,$$

$$(45) \quad 0 \rightarrow \mathcal{U}_l^*|_V \rightarrow \pi_1^*(\pi_{1*}\mathcal{U}_l)^*|_V \rightarrow (\text{Ker } E)^*|_V \rightarrow 0,$$

by pushing down to $SU(r, O_C(p + rl))$ we obtain an injective map Π :

$$(46) \quad \Pi: (\pi_{1*}\mathcal{U}_l)^*_{|V} \rightarrow \pi_{1*}(\text{Ker } E)^*_{|V},$$

where $\pi_{1*}(\text{Ker } E)^*_{|V} = \pi_{1*}\pi_2^*O_C(p+rl) \otimes B$. Note that by construction, Π turns out to be the restriction to V of the above map G , so we can conclude that actually $B = O_{SU(r, O_C(p+rl))}$ and $\det \pi_{1*}\mathcal{U}_l = \mathcal{L}$, and this concludes the proof. □

As an immediate consequence we have an alternative proof of the following well known result, see [N]:

PROPOSITION 2.5.2. *SU(r, 1) is a rational variety.*

§3. The main result

Let $\phi_{\mathcal{L}}: SU(r, 1) \rightarrow |\mathcal{L}|^*$ be the map associated to the theta divisor. By prop. (2.5.1) there exist s_0, \dots, s_N , independent global sections of $H^0(\mathcal{L})$ which define the rational map $p_l \cdot g_l$. Let V be the subspace spanned by them, we have a natural inclusion $V \hookrightarrow H^0(\mathcal{L})$, which induces a linear projection

$$(47) \quad \pi_l: |\mathcal{L}|^* \rightarrow \mathbf{P}(V^*) = \mathbf{P}^N$$

such that $g_l = \pi_l \cdot \phi_{\mathcal{L}}$. This allows us to prove that for any curve C of genus $g \geq 2$, the map $\phi_{\mathcal{L}}: SU(r, 1) \rightarrow \mathbf{P}^n$ has degree one.

3.1. Proof of theorem (0.0.2)

Actually, we will perform a non empty open subset \mathcal{U} of $SU(r, 1)$, such that the restriction of $\phi_{\mathcal{L}}$ to \mathcal{U} is actually injective, moreover we will prove that the tangent map $d(\phi_{\mathcal{L}})_{\xi}$ at a point ξ of \mathcal{U} is injective too.

Consider in $SU(r, 1) \times \text{Pic}^g(C)$ the set X containing pairs (ξ, l) satisfying property (*). We will denote by

$$(48) \quad \mathcal{U}: = p_1(X),$$

then \mathcal{U} is a non empty open subset of $SU(r, 1)$. First of all note that if $\xi \in \mathcal{U}$ the following set

$$(49) \quad \{l \in \text{Pic}^g(C) \mid (\xi, l) \text{ satisfies } (*)\}$$

is a non empty open subset of $\text{Pic}^g(C)$. Now let ξ_1 and ξ_2 be any two points of \mathcal{U} : then there exists l such that $(\xi_i, l) \in \mathcal{U}$, for $i = 1, 2$. For such an l ,

let $g_l: X_1 \rightarrow G_l$ be the rational map defined in (2.3), then by pr. (2.5.1) the restriction $g_l|_{X_l}$ is biregular and both ξ_1 and ξ_2 are in X_l . Now assume that $\phi_{\mathcal{L}}(\xi_1) = \phi_{\mathcal{L}}(\xi_2)$. Since $g_l = \pi_l \cdot \phi_{\mathcal{L}}$, then we have $g_l(\xi_1) = g_l(\xi_2)$. But $g_l|_{X_l}$ is injective, so we can conclude that $\xi_1 \simeq \xi_2$.

Assume now that $d(\phi_{\mathcal{L}})_{\xi}(v) = 0$ for a point $\xi \in \mathcal{U}$ and a tangent vector $v \in T_{SU(r,1),\xi}$. Let $l \in \text{Pic}^g(C)$ such that $(\xi, l) \in U$: then consider the rational map g_l , the linear projection π_l is defined at $\phi_{\mathcal{L}}(\xi)$, so we have

$$(50) \quad (d\pi_l)_{\phi_{\mathcal{L}}(\xi)} \cdot (d\phi_{\mathcal{L}})_{\xi} = (dg_l)_{\xi}.$$

Since $\xi \in X_l$ and $g_l|_{X_l}$ is biregular, then $(dg_l)_{\xi}(v) = 0$, hence $v = 0$, and $(d\phi_{\mathcal{L}})_{\xi}$ is injective. This concludes the proof.

3.2.

For $r \geq 3$, \mathcal{U} may be a proper subset of $SU(r, 1)$, that is there exist bundles ξ such that for any $l \in \text{Pic}^g(C)$ we have $\xi(l) \notin X_l$.

Let E be a semistable bundle on C of rank r , for any $l \in \text{Pic}^g(C)$ we have $h^0(E(l)) \geq \max(0, \chi(E(l)))$; actually there exists an open subset $U \subset \text{Pic}^g(C)$ such that for $l \in U$ this value is constant, following Raynaud, let's denote it by $h^0(E(l_{gen}))$, (see [R]). If $r \leq 2$ or $r = 3$ and the curve is general, then Raynaud proved that for any bundle we have $h^0(E(l_{gen})) = \max(0, \chi(E(l)))$; for $r \geq 4$ he showed the existence of bundles which do not satisfy this property, we will call such bundles Raynaud bundles, see [R].

Let $\eta \in SU(r)$: for any non zero morphism $\lambda \in \text{Hom}(\eta, C_p)$ the sheaf $\ker \lambda$ is actually a vector bundle on C with $\det \ker \lambda = O_C(-p)$:

$$(51) \quad 0 \rightarrow \ker \lambda \rightarrow \eta \xrightarrow{\lambda} C_p \rightarrow 0.$$

We claim that if η is stable then $\ker \lambda$ is stable too. In fact, if $\alpha \subset \ker \lambda \subset \eta$ is a destabilizing subbundle of $\ker \lambda$, then $\mu(\alpha) = \frac{d}{s} \geq \frac{-1}{r}$, with $s \leq r - 1$: this implies $d \geq 0$ and contradicts the stability of η . Let's define

$$(52) \quad \xi := \ker \lambda^*,$$

we can conclude that $\xi \in SU(r, O_C(p))$, and fits into the exact sequence

$$(53) \quad 0 \rightarrow \eta^* \rightarrow \xi \xrightarrow{v} C_p \rightarrow 0.$$

In the above notations, we can prove the following fact:

LEMMA 3.2.1. *If $\eta^* \in SU(r)$ is a stable Raynaud bundle, then for any $l \in \text{Pic}^g(C)$, we have $\xi(l) \notin X_l$.*

Proof. Consider the exact sequence (53) and tensor with $l \in \text{Pic}^g(C)$,

$$(54) \quad 0 \rightarrow \eta^*(l) \rightarrow \xi(l) \xrightarrow{v_l} C_p \rightarrow 0,$$

passing to cohomology, we can consider the following commutative diagram

$$(55) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^0(\eta^*(l)) & \longrightarrow & H^0(\xi(l)) & \xrightarrow{\bar{v}_l} & C \\ & & \downarrow & & \downarrow e_p & & \downarrow \\ & & (\eta^*(l))_p & \longrightarrow & \xi(l)_p & \xrightarrow{\bar{v}_{l,p}} & C_p \longrightarrow 0 \end{array}$$

Since η^* is a Raynaud bundle, then $h^0(\eta^*(l)) \geq r+1$ for any $l \in \text{Pic}^g(C)$, this implies that either $h^0(\xi(l)) \geq r+2$ for any $l \in \text{Pic}^g(C)$, or $h^0(\xi(l)) = r+1$ for l generic, and moreover \bar{v}_l is the zero map. In this case, $\text{Im } e_p \subset \text{Ker}(v_{p,l})$ for any l , which implies that $\xi(l)$ is not globally generated at p for any l . So we can conclude that $\xi \notin X_l$ for any $l \in \text{Pic}^g(C)$, and \mathcal{U} is a proper subset of $SU(r, O_C(p))$. □

REFERENCES

[A-C-G-H] E. Arbarello, M. Cornalba, P.A. Griffiths and J. Harris, *Geometry of Algebraic curves*, Springer verlag, Berlin, 1985.

[Be1] A. Beauville, *Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions theta*, Bull. Soc.Math.France, **116** (1988), 431–448.

[Be2] ———, *Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions theta II*, Bull. Soc. Math.France, **119** (1991), 259–291.

[B] S.Brivio, *On the degeneracy locus of a map of vector bundles on Grassmannian Varieties*, Preprint (1999).

[B-V1] S. Brivio and A. Verra, *The theta divisor of $SU_C(2)^s$ is very ample if C is not hyperelliptic*, Duke Math. J., **82** (1996), 503–552.

[B-V2] ———, *On the theta divisor of $SU(2,1)$* , Int. J. math., **10, 8** (1998), 925–942.

[D-N] I.M. Drezet and M.S. Narasimhan, *Groupes de Picard des variétés des modules des fibrés semistable sur les courbes algebriques*, Invent.Math., **97** (1989), 53–94.

[D-R] U.V. Desale and S. Ramanan, *Classificationn of vector bundles of rank two on hyperelliptic curves*, Invent. Math., **38** (1976), 161–185.

[H] R. Hartshorne, *Algebraic Geometry*, Springer verlag, New York, 1977.

[I-vG] E.I zadi and L. van Geemen, *The tangent space to the moduli space of vector bunldes on a curve and the singular locus of the theta divisor of the Jacobian*, Preprint (1997).

- [L] Y. Laszlo, *A propos de l'espace des modules de fibres de rang 2 sur une courbe*, Math. Ann., **299** (1994), 597–608.
- [N-R] M.S. Narasimhan and S. Ramanan, *Moduli of vector bundles on a compact Riemann surface*, Ann.Math., **89** (1969), 19–51.
- [N] P.E. Newstead, *Rationality of moduli spaces of vector bundles over an algebraic curve*, Math. Ann., **215** (1975), 251–268. Correction, ibidem, **249**, (1980), 281–282.
- [Ra] S. Ramanan, *The moduli spaces of vector bundles over an algebraic curve*, Math. Ann., **200** (1973), 69–84.
- [R] M. Raynaud, *Sections des fibrés vectoriels sur une courbe*, Bull.Soc.math. France, **110** (1982), 103–125.
- [S] C.S. Seshadri, *Fibrés vectoriels sur les courbes algébriques*, Astérisque, **96** (1982), 3–50.

Sonia Brivio
Dipartimento di Matematica
Università di Pavia
via Abbiategrasso
209 - 27100 Pavia
Italy
`Brivio@dimat.unipv.it`

Alessandro Verra
Dipartimento di Matematica
Università di Roma Tre
largo S. Leonardo Murialdo 1 - 00146
Roma
Italy
`Verra@matrm3.mat.uniroma3.it`