

- independent risk factor for invasive aspergillosis and zygomycosis in patients with hematologic malignancy. *Infect Control* 1987;8:71-75.
45. Sherertz RJ, Belani A, Kramer BS, et al. Impact of air filtration on nosocomial *Aspergillus* infections. Unique risk of bone marrow transplant recipients. *Am J Med* 1987;83:709-718.
 46. Rhame FS. Prevention of nosocomial aspergillosis. *J Hosp Infect* 1991;18(suppl A):466-472.
 47. Rhame FS, Streifel AJ, Kersey JH Jr, et al. Extrinsic risk factors for pneumonia in the patient at high risk of infection. *Am J Med* 1984;76:42-52.
 48. Brachman PS. Nosocomial infection-airborne or not? In: Brachman PS, Eickhoff TC, eds. In: *Proceedings of the International Conference on Nosocomial Infections*. Chicago, IL: American Hospital Association; 1971:189-192.
 49. Kundsinn RB. Documentation of airborne infection during surgery. *Ann NY Acad Sci* 1980;353:255-261.
 50. Ayliffe GA. Role of the environment of the operating suite in surgical wound infection. *Rev Infect Dis* 1991;13(suppl):S800-S804.
 51. Haley RW, Tenney JH, Lindsey JO II, et al. How frequent are outbreaks of nosocomial infection in community hospitals? *Infect Control* 1985;6:233-236.
 52. Wenzel RP, Thompson RL, Landry SM, et al. Hospital acquired infections in intensive care unit patients: an overview with emphasis on epidemics. *Infect Control* 1983;4:371-375.
 53. Jarvis WR. Nosocomial outbreaks: the Centers for Disease Control's Hospital Infections Program experience, 1980-1990. *Am J Med* 1991;91(suppl 3B):101S-106S.
 54. Eickhoff TC. Microbiological sampling: hospitals. *JAHA*. 1970;44:86-87.

Whirlpool Bath Source of Outbreak of Legionnaires' Disease Aboard Cruise Ship

by Gina Pugliese, RN, MS
Medical News Editor

As of August 10, 1994, a total of 14 passengers aboard the cruise ship *Horizon* had Legionnaires' disease (LD) confirmed by either sputum culture, detection of antigens of *Legionella pneumophila* serogroup 1 (LP₁) in urine by radioimmunoassay, or fourfold rise in antibody titers between acute- and convalescent-phase serum specimens. Under investigation are 28 other passengers with pneumonia that occurred

within 2 weeks after sailing aboard the cruise ship. Cases have occurred from separate week-long cruises between April 30 and July 9, 1994.

A case-control study revealed a strong association between exposure to whirlpool baths and illness. Cultures taken from a sand filter, used for recirculation of whirlpool water, yielded an isolate of LP₁; this isolate and the clinical isolate had matching monoclonal antibody subtyping patterns. Various interventions were completed, including hyperchlorination of the ship's pota-

ble water supply, removal of the whirlpool filters, and discontinuation of the whirlpool baths.

Additional recommendations to reduce the risk of transmission from whirlpool baths aboard cruise ships will be the subject of a special meeting scheduled to be held this fall. Information about the meeting is available from CDC's National Center for Environmental Health: (404) 488-7093.

FROM: Update: outbreak of Legionnaires' disease associated with a cruise ship. *MMWR* 1994;43(31):574-575.