

# Salt intake and dietary sources of salt on weekdays and weekend days in Australian adults

Caryl Nowson<sup>1</sup>, Karen Lim<sup>1</sup>, Mary-Ann Land<sup>2</sup>, Jacqui Webster<sup>2</sup>, Jonathan E Shaw<sup>3</sup>, John Chalmers<sup>2</sup>, Victoria Flood<sup>4</sup>, Mark Woodward<sup>2,5</sup> and Carley Grimes<sup>1,\*</sup>

<sup>1</sup>Institute for Physical Activity and Nutrition (IPAN), School of Exercise & Nutrition Sciences, Deakin University, Locked Bag 20000, Waurn Ponds, Geelong, VIC 3220, Australia; <sup>2</sup>George Institute for Global Health, University of New South Wales, Sydney, Australia; <sup>3</sup>Clinical Diabetes and Epidemiology, Baker IDI Heart & Diabetes Institute, Melbourne, Australia; <sup>4</sup>Faculty of Health Sciences, University of Sydney and Western Sydney Local Health District, Sydney, Australia; <sup>5</sup>The George Institute for Global Health, University of Oxford, Oxford, UK

Submitted 8 August 2017: Final revision received 22 November 2017: Accepted 6 December 2017: First published online 1 February 2018

## Abstract

**Objective:** To assess if there is a difference in salt intake (24 h urine collection and dietary recall) and dietary sources of salt (Na) on weekdays and weekend days.

**Design:** A cross-sectional study of adults who provided one 24 h urine collection and one telephone-administered 24 h dietary recall.

**Setting:** Community-dwelling adults living in the State of Victoria, Australia.

**Subjects:** Adults ( $n$  598) who participated in a health survey (53.5% women; mean age 57.1 (95% CI 56.2, 58.1) years).

**Results:** Mean (95% CI) salt intake (dietary recall) was 6.8 (6.6, 7.1) g/d and 24 h urinary salt excretion was 8.1 (7.8, 8.3) g/d. Mean dietary and 24 h urinary salt (age-adjusted) were 0.9 (0.1, 1.6) g/d ( $P=0.024$ ) and 0.8 (0.3, 1.6) g/d ( $P=0.0017$ ), respectively, higher at weekends compared with weekdays. There was an indication of a greater energy intake at weekends (+0.6 (0.02, 1.2) MJ/d,  $P=0.06$ ), but no difference in Na density (weekday: 291 (279, 304) mg/MJ; weekend: 304 (281, 327) mg/MJ;  $P=0.360$ ). Cereals/cereal products and dishes, meat, poultry, milk products and gravy/sauces accounted for 71% of dietary Na.

**Conclusions:** Mean salt intake (24 h urine collection) was more than 60% above the recommended level of 5 g salt/d and 8–14% more salt was consumed at weekends than on weekdays. Substantial reductions in the Na content of staple foods, processed meat, sauces, mixed dishes (e.g. pasta), convenience and takeaway foods are required to achieve a significant consistent reduction in population salt intake throughout the week.

**Keywords**  
Salt  
Sodium  
Sodium chloride  
Diet  
Weekend  
Urinary sodium  
Australia

It is recognised that reducing average salt intake from current high levels (8–9 g/person per d), and specifically reducing the salt (Na) content of processed foods, should have a significant impact on reducing the number of myocardial infarctions and strokes in Australia<sup>(1)</sup>. Australia is a member of the WHO Member States who have agreed on a voluntary global non-communicable disease target of a 30% relative reduction in mean population intake of salt, with the aim of achieving less than 5 g salt/d (approximately 2 g Na/d) by 2025<sup>(2)</sup>.

Utilising 24 h dietary recall methodology, the 2011–12 Australian Health Survey indicated an average salt intake of 7.1 g/d for males and 5.4 g/d for females<sup>(3)</sup> which is lower than the average salt intake of 9.2 g/d for males and 6.7 g/d for females as measured by 24 h urine collections in over 300 people in a recent population-based study<sup>(4)</sup>. The difference in estimated intake of salt between the two

methods may relate to the inability of dietary recall to capture the amount of discretionary salt used and the restricted range of foods included in food databases, which prevents analysis that captures all the variations in salt content of different brands of manufactured foods. Dietary assessment does, however, provide valuable information on the relative contributions of different foods to total Na intake, as in developed countries manufactured foods have been estimated to contribute at least 70% of the daily amount of salt consumed<sup>(5–7)</sup>. Data from the Australian Health Survey in adults, using a 24 h dietary recall, indicated that the major dietary sources of Na are cereal-based products and dishes (25%; mainly from the mixed dishes where cereal is the major ingredient, e.g. pasta dishes), cereal and cereal products (18%; mainly bread) and meat and poultry (18%; mainly processed meat and mixed dishes)<sup>(3)</sup>. In addition, it is recognised that

\*Corresponding author: Email carley.grimes@deakin.edu.au

restaurant meals, takeaway foods and convenience foods contain relatively high concentrations of Na<sup>(8)</sup>. In New Zealand an assessment of the commonly consumed fast foods in the national dietary survey of 2008/9 indicated that the majority exceeded the Na targets set by the UK Food Standards Agency in 2012<sup>(9)</sup>. In addition many discretionary foods (which are energy-dense and nutrient-poor), such as processed meats and sausages, savoury pastries and pies, commercially fried foods, potato chips, crisps and other fatty and/or salty snack foods, contain high levels of added salt<sup>(10)</sup>. The recent 2011–12 Australian Health Survey indicated that, on average, just over one-third (35%) of total daily energy consumed was reported to be from discretionary foods<sup>(3)</sup>. Therefore, the frequency of consumption of these discretionary foods is likely to impact on the usual level of salt (Na) intake.

It is known that dietary patterns alter on weekends and holidays and it may be that more salt is consumed due to a different selection of food products. There is an indication that people are more likely to eat away from home and use pre-prepared convenience foods and takeaway foods at weekends<sup>(11)</sup>. In the Australian 1995 National Nutrition Survey, adults were found to have higher intakes of muscle meat and sausages on weekends<sup>(12)</sup> and in the USA, weekend consumption has been associated with increased energy intake and poorer diet quality<sup>(13)</sup>. Given that Na intakes are well above dietary recommendations it is useful to consider if dietary Na intake differs between weekdays and weekends as this information can be used to tailor interventions that seek to alter eating habits and/or utilise Na reformulation targets. Therefore, we aimed to compare salt intake (measured by 24 h urine collection and dietary recall) and dietary sources of salt on weekdays and weekend days in a sample of Australian adults.

## Methods

Participants were recruited from a sample of 3653 people (38% response rate) participating in the Victorian Health Monitor survey (VHM; May 2009 to April 2010)<sup>(14)</sup>, for which the recruitment strategy and methods have previously been documented<sup>(4)</sup>. In 2011, a subgroup of 3487 people who participated in the VHM survey, lived within 100 km of a commercial pathology centre and who had a valid current address were invited to participate by letter in the present salt study. Participation involved provision of a 24 h urine collection, additional information on discretionary salt use and self-reported body weight, and participation in a telephone-administered interview providing a 24 h dietary recall over the telephone (utilising visual aids used in the previous VHM survey to estimate serving sizes). Of the 3487 adults who were sent letters of invitation to participate in the salt survey, 1003 people registered interest and provided telephone contact details.

Of those who registered interest, 697 people were selected at random to participate to meet the sampling size criteria. This resulted in participation of 605 people (326 women, 279 men), representing 17% of those who received invitations to participate. All participants provided written informed consent.

Participants recorded the start and end times for their urine collection and reported any missed urine collection during the 24 h period. Urine collections were considered valid if the collection time fell between 14 and 31 h and all urinary excretion results were standardised to a 24 h period. The 24 h urine samples were considered 'week-end' collections if the first void of the collection was on a Saturday or Sunday morning. The urine samples were received, processed and analysed by the Australian accredited pathology service centres of Healthscope and Dorevitch. Urinary Na and K concentrations were determined using ion-selective electrodes and urinary creatinine concentration was determined from the kinetic Jaffe reaction using the Advia 2400 Clinical Chemistry System (Siemens). Assessment of completeness of urine collections was made by assessing the predicted 24 h urine excretion of creatine based on age and gender<sup>(15)</sup>. In addition, criteria based on low urine volume (<500 ml), reported missed collections and extreme statistical outliers, which are similar to those previously utilised, were used to identify probable over- and under-collection of 24 h samples<sup>(16)</sup>. Of the 605 participants, seven urine collections were excluded due to likely errors in collection: three women with 24 h creatinine <4 mmol/d, one man with 24 h creatinine >3 SD below the mean, two women with 24 h urine volume <500 ml and one woman who reported missing more than one void during the collection period, resulting in the final sample of 598 participants.

### *Demographic, anthropometric and dietary data*

Demographic data and height measurements were obtained as part of the VHM survey. Height was measured to the nearest 0.1 cm without shoes using a stadiometer. Annual household income and highest level of education of the participant were self-reported in VHM and used as indicators of socio-economic status. Participants provided information on their current body weight, together with information on knowledge and attitudes to salt intake and salt use during cooking and at the table, during a telephone interview.

All participants received a shopping voucher to the value of \$AU 20 or a double-pass movie ticket on completion of each 24 h urine collection, together with a summary of their individual results.

Participants completed one 24 h diet recall via telephone which was scheduled within 3 d of completing their 24 h urine collection (mean elapsed time = 5.7 (SD 6.3) d; median = 3 (interquartile range 1–8) d; range = 1–31 d). A five-pass method was used to record food and beverage intakes on the day prior to the interview based on the

method validated by the US Department of Agriculture<sup>(17,18)</sup>. All participants were provided with a food model booklet to assist with estimating portion sizes<sup>(19)</sup>. The 24 h diet recalls were considered 'weekend' intakes if participants recalled the foods and beverages consumed on a Saturday or Sunday.

Low energy reporters were identified following the Goldberg cut-off method (ratio of energy intake (excluding dietary fibre) to BMR <0.9)<sup>(20)</sup>, with BMR estimated using the Schofield equation for the specific gender and age<sup>(21)</sup>.

Foods and beverages reported by each participant in his/her 24 h diet recall were entered into FoodWorks 7 software (Xyris Software, Brisbane, Queensland, Australia) and nutrient intakes were estimated using the Australian food composition database AUSNUT 2007<sup>(22)</sup>. Dietary intake did not include discretionary salt or dietary supplements. We have presented the mean intakes of Na and other nutrients of the group, the proportion of energy sourced from macronutrients, Na and K density (milligrams per megajoule of energy), and the molar Na:K ratio. The food groupings specified in the AUSNUT 2007<sup>(22)</sup> database were used to estimate the food contributions to dietary Na intake. The amount of Na (milligrams) contributed by each food group was divided by the mean dietary Na intake of the study sample, multiplied by 100 and presented as a percentage.

### Statistical analysis

Participant characteristics are presented as mean and SD, mean and 95% CI, or *n* and %. Comparisons were made using independent *t* tests and  $\chi^2$  tests for continuous and categorical variables, respectively. Subgroup analyses by gender were completed to assess gender differences. To compare bivariate, continuous associations, we used Pearson's correlation. Age-adjusted residuals were derived from regression analysis for men and women separately. A *P* value of <0.05 was considered significant, throughout. Data were analysed using the statistical software package Stata/SE version 14.1.

### Results

There were 598 participants, about half of whom were women (Table 1). Twenty-five women and twenty-nine men were identified as having potentially implausibly low energy intakes (9.2% of sample). The twenty-five female under-reporters had energy intakes 47% lower than the rest of the women (4474 kJ/d compared with 8449 kJ/d) and the twenty-nine male under-reporters had energy intakes 48% lower than the rest of the men (5693 kJ/d compared with 11031 kJ/d). A sensitivity analysis was conducted which included these fifty-four potential under-reporters, which did not change the results in any meaningful way: mean salt intake of the sample was 7.1 (SD 0.1) g/d when these fifty-four participants were excluded and

6.8 (SD 0.1) g/d when they were included (*P*=0.283). Importantly, there was no difference in the proportion of under-reporters on weekend days compared with weekdays (under-reporters: 9.3% on weekdays *v.* 8.0% on weekend days;  $\chi^2$  *P*=0.661), therefore all participants were included.

The average salt intake assessed from the 24 h diet recall was approximately 16% lower than that assessed from the mean 24 h urine excretion (Tables 1 and 2). There was a significant modest correlation between dietary intake and urinary excretion of salt (*r*=0.343, *P*<0.001). Men reported consuming more energy (+2.5 MJ/d; 24 h recall) than women and men reported and excreted more salt than women, and these gender differences in energy intake and salt excretion remained even after adjustment for body weight (reported dietary salt intake: men 7.6 (95% CI 7.2, 8.1) g/d *v.* women 6.1 (95% CI 5.7, 6.5) g/d, *P*<0.001; 24 h salt excretion: men 9.0 (95% CI 8.6, 9.3) g/d *v.* women 7.3 (95% CI 6.9, 7.6) g/d, *P*<0.001).

The mean 24 h salt excretion in this group was 8.1 (95% CI 7.8, 8.3) g/d. Participants who performed their urine collection on a weekday compared with a weekend day were on average 4.0 (95% CI 1.9, 6.1) years older (weekday: 58.3 (95% CI 57.2, 59.4) years *v.* weekend day: 54.3 (95% CI 52.5, 56.1) years), but there were no differences by gender or body weight. There were no differences in the age, gender, body weight or BMI of participants who reported weekday intakes in their 24 h recall compared with those who reported weekend day intakes. Overall, reported dietary salt intake and 24 h urinary excretion (adjusted for age) were higher on weekend days (Saturday and Sunday) compared with weekdays (Table 3). The mean reported dietary salt intake was 0.9 g/d (14%) higher on weekend days and age-adjusted urinary salt excretion was 0.8 g/d (8%) higher on weekend days compared with weekdays. When split by gender, the difference in salt excretion between weekdays and weekend days was no longer statistically significant. There was an indication of a 6% greater intake of energy on weekend days (*P*=0.06) but Na density did not differ between weekdays and weekends. For men only, there was a 16% greater reported salt intake on weekend days, but no difference in energy intake.

The major food sources of Na were cereals and cereal products (19%) which included breads and bread rolls (13%). This was followed by meat and poultry (16%), then cereal-based products and dishes (15%), milk products (11%) and gravy and sauces (11%), which in total accounted for 71% of dietary Na. Most of the Na within the meat and poultry group (contributing 16% of total Na) came from processed meats (8%) including ham (4%) and bacon (2%). Within the cereal-based product group, the major sources of Na were savoury rice-based dishes (2%; fried rice, risotto and sushi), pizza (1%) and pasta and noodle dishes (1%). Of the cheeses, combined regular and

**Table 1** Demographic characteristics, dietary electrolyte intake and 24 h urinary excretion of electrolytes in community-dwelling adults (*n* 598) living in the State of Victoria, Australia, 2011\*,†

	Overall ( <i>n</i> 598)		Women ( <i>n</i> 320)		Men ( <i>n</i> 278)		P‡
	<i>n</i> or Mean	% or 95% CI	<i>n</i> or Mean	% or 95% CI	<i>n</i> or Mean	% or 95% CI	
Women	320	53.5					
Age (years)	57.1	56.2, 58.1	56.7	55.4, 57.9	57.7	56.2, 59.1	0.309
Weight (kg)§	76.2	75.0, 77.5	69.6	68.0, 71.2	83.8	82.3, 85.4	<0.001
BMI (kg/m <sup>2</sup> )§	26.6	26.2, 27.0	26.3	25.7, 26.9	27.0	26.6, 27.5	0.062
Urinary Na (mmol/d)	137.9	133.4, 142.5	118.4	113.4, 123.4	160.3	153.2, 167.5	<0.001
Urinary salt (g/d)	8.1	7.8, 8.3	6.9	6.6, 7.2	9.4	8.9, 9.8	<0.001
Urinary K (mmol/d)	77.8	75.8, 79.7	70.7	68.3, 73.1	85.8	83.0, 88.7	<0.001
Urinary Na:K (mmol)	1.86	1.80, 1.93	1.78	1.69, 1.86	1.96	1.87, 2.06	0.003
Diet Na (mmol/d)	116	111, 121	103	97, 109	131	124, 138	<0.001
Diet K (mmol/d)	93	91, 96	87	84, 90	100	96, 105	<0.0001
Dietary Na:K (mmol)	1.4	1.3, 1.4	1.3	1.2, 1.4	1.4	1.3, 1.5	0.039
Weekend urine sample	179	29.9	89	27.8	90	32.4	0.224
Weekend diet recall	127	21.2	65	20.3	62	22.3	0.553
Metropolitan Melbourne	366	61.2	210	65.6	156	56.1	0.017
Age range							
< 40 years	65	10.9	37	11.6	28	10.1	0.707
40–60 years	261	43.7	142	44.4	119	42.8	
≥ 60 years	272	45.5	141	44.1	131	47.1	
Annual household income							
≤ \$AU 40 000	156	26.1	94	29.4	62	22.3	0.211
\$AU 40 000–70 000	144	24.1	74	23.1	70	25.2	
> \$AU 70 000	265	44.3	133	41.6	132	47.5	
Don't know/refused	33	5.5	19	5.9	14	5.0	
Highest level of education							
No tertiary education	346	57.9	192	60.0	154	55.4	0.255
With tertiary education	252	42.1	128	40.0	124	44.6	

\*Categorical variables presented as *n* and %; continuous variables presented as mean and 95% CI.

†Electrolyte excretion from one 24 h urine collection; dietary intake from one 24 h diet recall.

‡*P* value for the difference between women and men. Gender differences in categorical variables compared using the  $\chi^2$  test; gender differences in continuous variables compared using the independent-samples *t* test.

§Self-reported weight.

|| Grams of salt = mmol Na  $\times$  23  $\times$  2.54/1000.**Table 2** Nutrient intakes of community-dwelling adults (*n* 598) living in the State of Victoria, Australia, 2011\*

Nutrient	Overall ( <i>n</i> 598)		Women ( <i>n</i> 320)		Men ( <i>n</i> 278)	
	Mean	95% CI	Mean	95% CI	Mean	95% CI
Energy (MJ/d)	9.2	8.9, 9.4	8.1	7.9, 8.4	10.5	10.0, 1.1
Protein (%E)	19.1	18.6, 19.5	19.1	18.6, 19.7	19.0	18.2, 19.7
Fat (%)	32.4	31.7, 33	33.2	32.3, 34.2	31.3	30.5, 32.2
Saturated fat (%E)	12.0	11.7, 12.4	12.1	11.6, 12.5	12.0	11.5, 12.4
Carbohydrate (%E)	41.8	41.1, 42.6	41.6	40.5, 42.6	42.1	41, 43.3
Alcohol (%E)	3.0	2.6, 3.4	2.4	1.9, 2.8	3.8	3.2, 4.4
Protein (g/d)	104	101, 107	93	89, 96	118	112, 123
Total fat (g/d)	84	81, 86	76	73, 80	92	87, 97
Saturated fat (g/d)	31	30, 33	28	26, 30	35	33, 38
Carbohydrate (g/d)	244	236, 252	213	204, 221	280	267, 292
Na (mg/d)	2664	2555, 2773	2367	2230, 2504	3007	2842, 3171
Salt (g/d)	6.8	6.6, 7.1	6.1	5.7, 6.4	7.7	7.3, 8.1
Na density (mg/MJ)	294	283, 305	295	279, 311	293	278, 308
K (mg/d)	3631	3528, 3735	3387	3261, 3513	3912	3750, 4075
K density (mg/MJ)	408	398, 418	427	413, 441	386	372, 400

%E, percentage of energy.

\*Dietary intake from one 24 h diet recall.

reduced-fat natural, traditional cheese (category includes cheddar, feta, mozzarella) contributed the most Na (4%) followed by processed cheese (2%). Overall, the gravies and savoury sauces group provided 8% of total Na and, within this group, the major contributor to Na was savoury sauces (6%), of which soya sauce contributed to 3%.

Other foods contributing  $\geq 1\%$  of Na intake included stock cubes (2%).

When the contribution of different food groups to total dietary Na consumed on weekdays and weekends was assessed, there was some indication, although not significant, of a greater contribution of gravies and savoury sauces,

**Table 3** Dietary and urinary sodium/salt on weekdays compared with weekend days in community-dwelling adults (n 598) living in the State of Victoria, Australia, 2011

	Overall (n 598)						Women (n 320)						Men (n 278)					
	Weekday (n 471)			Weekend (n 127)			Weekday (n 255)			Weekend (n 65)			Weekday (n 216)			Weekend (n 62)		
	Mean	95% CI	P*, †	Mean	95% CI	P*, †	Mean	95% CI	P*, †	Mean	95% CI	P*, †	Mean	95% CI	P*, †	Mean	95% CI	P*, †
24 h Dietary recall	9.1	8.8, 9.4	0.060	9.7	9.1, 10.3	0.060	8.1	7.8, 8.3	0.187	8.5	7.9, 9.1	0.187	10.3	9.9, 10.8	0.032	10.9	10.0, 11.8	0.270
Energy intake (MJ/d)	291	279, 304	0.360	304	281, 327	0.360	296	278, 314	0.819	292	262, 321	0.819	286	270, 302	0.099	316	280, 353	0.099
Na density (mg/MJ)	6.6	6.3, 6.9	0.024	7.5	6.8, 8.2	0.024	6.0	5.6, 6.4	0.389	6.4	5.6, 7.2	0.389	7.4	7.0, 7.9	0.032	8.7	7.6, 9.9	0.032
Dietary salt (g/d)‡																		
	Weekday (n 419)			Weekend (n 179)			Weekday (n 231)			Weekend (n 89)			Weekday (n 188)			Weekend (n 90)		
Salt excretion	Mean	95% CI		Mean	95% CI		Mean	95% CI		Mean	95% CI		Mean	95% CI		Mean	95% CI	
Urinary salt (g/d)‡, §	10.4	10.1, 10.7	0.017	11.2	10.6, 11.7	0.017	8.4	8.1, 8.8	0.113	9.0	8.4, 9.6	0.113	13.2	12.8, 13.6	0.173	13.9	13.0, 14.7	0.173

\*Unpaired *t* test for difference weekday v. weekend.†*F* test specifying unequal variance.

‡Grams of salt = mmol Na × 23 × 2.54/1000.


§Urinary salt adjusted for age (unstandardised predicted value).

processed meat, mixed dishes where cereal is a major ingredient, seasonings and stock cubes to total Na on weekend days compared with weekdays, together with a reduced contribution from plain breads (Fig. 1). There was some indication for more Na to be consumed from the meat group on weekends (+149 mg,  $P=0.053$ ), with 96 mg Na more from processed meat ( $P=0.088$ ), compared with weekdays (Table 4).

## Discussion

Overall dietary salt intake was 8–14% higher on weekends (Saturday and Sunday) compared with weekdays as measured by 24 h excretion or 24 h dietary recall, respectively. The top five major dietary sources of Na in this sample were similar to those found in the national 2011–12 Australian Health Survey<sup>(3)</sup>. Both surveys indicated that more than 70% of dietary Na (74% in the national survey and 71% in the present survey) was derived from five major food group categories: cereal-based products and dishes; cereals and cereal products; meat and poultry and game products and dishes; milk products and dishes; and sauces. Most of the Na within the meat and poultry group came from processed meats, including ham and bacon. This highlights the importance of ensuring that clear Na reduction reformulation targets are specified for these foods if the Australian population is to achieve a 30% reduction in Na intake by 2025. It has been estimated that a 35% reduction in total population salt intake could be achieved by a 36% reduction in the Na content of packaged foods in conjunction with a 40% reduction in discretionary salt use and the Na content of foods consumed away from home. This range of reductions would result in the population meeting the WHO target for a 30% reduction in dietary salt<sup>(23)</sup>. Currently, progress in reducing the Na content of key food items in Australia has been slow. A recent analysis of the change in the Na content of pasta sauces available and the impact of the Food and Health Dialogue Na target for this product indicated that the average level of Na was above the 2012 UK target and that manufacturers would need to reformulate pasta sauce products well beyond the Food and Health Dialogue target to achieve a meaningful reduction in Na content<sup>(24)</sup>. Furthermore, although there appears to have been a small fall in the Na content of fast foods produced by large fast-food chains between 2009 and 2012<sup>(8)</sup>, the average Na content per serving remained high at 1.5 g salt/serving.

Although there was an indication, which did not reach significance, of a slightly greater intake of energy (7%) on weekend days compared with weekdays, Na density did not differ at weekends. Men particularly appeared to consume more salt on weekend days (18% greater dietary salt intake compared with weekdays). Therefore, it appears that the higher salt intake at weekends could be due to a small increase in food intake combined with a small shift in the range of processed foods eaten. Although



**Fig. 1** Main food sources of sodium (salt) by weekday (■) and weekend day (□) in the diet of community-dwelling adults ( $n$  598; 53.5% women; mean age 57.1 (95% CI 56.2, 58.1) years) living in the State of Victoria, Australia, 2011. Percentage contribution (with 95% CI represented by horizontal bars) of food groups\* to sodium intake by type of day. \*Foods are categorised according to AUSNUT 2007 (3-digit code level). Only foods contributing  $\geq 1\%$  are presented

individually not significant, there were small shifts in specific food groups with more Na being derived from gravies and savoury sauces, processed meat, mixed cereal dishes, seasonings and stock cubes on weekend days compared with weekdays. As these flavouring and seasoning agents (gravies, savoury sauces, stock cubes) typically contain the highest concentrations of Na in foods, there is significant scope for manufacturers to gradually reduce the Na content of these products and significantly reduce Na intakes on weekends and holidays. Soya sauce is one such product which, although only small amounts may be used in one day, on average contributes 3% of salt intake.

The comparability of these results, using two different methodologies, confirms within our selected sample population group that those reporting weekend dietary intake and performing 24 h urine collections at the weekend had a greater intake, resulting in 0.9 g salt/d higher dietary intake and 0.8 g salt/d from 24 h urinary excretion. These results are similar to data from the US National Health and Nutrition Examination Survey (NHANES) where on average adults consumed 0.5 g more salt on a Saturday compared with a weekday<sup>(13)</sup>. Generally people appear to consume less healthy foods at weekends<sup>(25)</sup> and may be more likely to consume more discretionary foods. In NHANES, there was an increase in energy from discretionary foods at weekend days which contributed to greater intakes of total fat, saturated fat, sugar and Na<sup>(13)</sup>. We found some indication that on weekend days, more Na came from processed meats in

contrast to Na coming from fresh meat consumption, which tended to fall.

Although we did not collect information on which foods were eaten outside the home, there is evidence from the USA that eating fast foods and eating in full-service restaurants are more prevalent at weekends<sup>(13)</sup>. In addition, it has been found that eating at both fast-food and full-service restaurants is associated with increased intakes of total energy, total fat, saturated fat, cholesterol and Na (0.75 and 1.0 g salt/d, respectively)<sup>(26)</sup>. In US high-school children, a 10-year longitudinal study found that frequency of fast-food intake predicted Na intake, such that those consuming fast foods at least four times weekly had a 5% higher daily intake of Na compared with those consuming none<sup>(27)</sup>. The average amount of Na contained in fast-food products from large food outlet chains in Australia has been reported to be 605 mg Na (1.5 g salt) per serving<sup>(8)</sup> and there is huge scope to reduce the Na content of many of these high-Na foods, which people are more likely to consume on a regular basis at weekends. Salt does serve some technological functions in food, such as dough development in bread and water binding and preservation in meats; however, for many food products, the amount of salt added generally exceeds that needed to enhance taste<sup>(28)</sup>.

The strength of the current study is that we collected both an objective measure of Na intake (24 h urine excretion) together with information on dietary intake

**Table 4** Major food group sources of dietary sodium (*n* 598) in the diet of community-dwelling adults (*n* 598) living in the State of Victoria, Australia, 2011

Food group*	% contribution to daily Na intake	Weekend day Na intake (mg/d)		Weekday Na intake (mg/d)		P†
		Mean	95% CI	Mean	95% CI	
Cereals and cereal products	19.3	531	459, 603	510	471, 549	0.618
<i>Regular breads, and bread rolls (plain/unfilled/untopped)</i>	<i>13.4</i>	<i>361</i>	<i>295, 427</i>	<i>357</i>	<i>323, 390</i>	<i>0.901</i>
<i>Breakfast cereals and bars, unfortified and fortified varieties</i>	<i>2.3</i>	<i>70</i>	<i>40, 101</i>	<i>60</i>	<i>49, 71</i>	<i>0.519</i>
<i>English-style muffins, flat breads, savoury &amp; sweet breads</i>	<i>2.0</i>	<i>67</i>	<i>32, 102</i>	<i>49</i>	<i>35, 63</i>	<i>0.346</i>
Meat, poultry, game products	15.8	539	397, 682	390	341, 440	0.053
<i>Processed meat</i>	<i>7.7</i>	<i>282</i>	<i>178, 385</i>	<i>186</i>	<i>150, 223</i>	<i>0.088</i>
<i>Sausages, frankfurts, saveloys</i>	<i>2.3</i>	<i>71</i>	<i>21, 121</i>	<i>57</i>	<i>36, 78</i>	<i>0.613</i>
<i>Muscle meat</i>	<i>2.1</i>	<i>58</i>	<i>29, 88</i>	<i>55</i>	<i>39, 70</i>	<i>0.840</i>
Cereal-based products and dishes	14.6	439	308, 571	377	323, 431	0.386
<i>Mixed dishes where cereal is the major ingredient</i>	<i>5.2</i>	<i>180</i>	<i>84, 276</i>	<i>129</i>	<i>91, 166</i>	<i>0.328</i>
<i>Pastries</i>	<i>3.3</i>	<i>90</i>	<i>41, 139</i>	<i>87</i>	<i>58, 116</i>	<i>0.926</i>
<i>Savoury biscuits</i>	<i>2.1</i>	<i>71</i>	<i>26, 116</i>	<i>53</i>	<i>41, 66</i>	<i>0.467</i>
Milk products and dishes	10.7	311	254, 367	279	253, 305	0.279
<i>Cheese</i>	<i>5.7</i>	<i>174</i>	<i>124, 224</i>	<i>146</i>	<i>122, 169</i>	<i>0.290</i>
<i>Dairy milk (cow, sheep, goat)</i>	<i>3.5</i>	<i>94</i>	<i>78, 109</i>	<i>92</i>	<i>84, 101</i>	<i>0.862</i>
Savoury sauces and condiments	10.6	327	210, 444	269	222, 317	0.368
<i>Gravies and savoury sauces</i>	<i>8.3</i>	<i>253</i>	<i>142, 363</i>	<i>214</i>	<i>168, 260</i>	<i>0.522</i>
<b><i>Soya sauce</i></b>	<b><i>3.0</i></b>	<b><i>107</i></b>	<b><i>27, 188</i></b>	<b><i>73</i></b>	<b><i>44, 103</i></b>	<b><i>0.432</i></b>
Miscellaneous	6.4	217	140, 294	157	115, 198	0.183
<i>Herbs, spices, seasonings and stock cubes</i>	<i>4.6</i>	<i>163</i>	<i>88, 238</i>	<i>111</i>	<i>73, 150</i>	<i>0.230</i>
Soup	4.6	112	48, 177	124	80, 168	0.772
<i>Dry soup mix</i>	<i>2.4</i>	<i>57</i>	<i>2, 111</i>	<i>66</i>	<i>32, 99</i>	<i>0.789</i>
Vegetable products and dishes	4.3	137	97, 177	108	94, 122	0.182
Fish and seafood products/dishes	3.5	86	55, 118	94	75, 114	0.673
Non-alcoholic beverages	2.0	52	38, 65	53	46, 60	0.851
Legume, pulse products/dishes	1.9	34	11, 57	56	38, 75	0.146
Fats and oils	1.9	57	42, 71	49	41, 56	0.316
Snack foods	0.9	15	4, 25	27	16, 37	0.112
Egg products and dishes	0.9	27	16, 39	23	17, 28	0.462
Alcoholic beverages	0.8	24	18, 31	22	18, 26	0.575
Confectionery and cereal/nut/fruit/seed bars	0.6	13	8, 19	18	14, 22	0.144
Seed and nut products and dishes	0.4	9	3, 15	12	9, 16	0.333
Dairy substitutes	0.4	8	2, 15	12	7, 17	0.279
Fruit products and dishes	0.3	8	5, 10	7	6, 8	0.497
Sugar products and dishes	0.1	4	1, 6	3	2, 3	0.379
Special dietary foods	0.1	1	-1, 3	2	0, 4	0.720

\*Food groups are presented at the 2-digit code level, with the 3-digit code level presented in italics and the 8-digit code level presented in bold italics (only for soya sauce).

†Unpaired *t* test.

over a 24 h period using robust standard methodology. Although the study included two different groups of people who collected their urine and reported their dietary intake on different days of the week (30 and 21%, respectively, performed their 24 h urine collection and reported their dietary intake at weekends), we did find that both urinary salt excretion and dietary intake were higher at weekends in this group with an average age of 57 years.

## Conclusions

Salt intake in in this population, assessed from 24 h urine collections, was between 35 and 62% above recommended levels of 5–6 g/d. Both urine excretion of salt and dietary information confirmed that salt intake was 8 and 14%, respectively, higher on weekend days, which

raises the long-term average of an individual's daily consumption of salt. It is also likely that the dietary habits and food choices at weekends, which contribute to a greater consumption of salt on weekend days, will be similar to those seen for holiday periods, leading to long-term higher intakes which over a lifetime are likely to increase cardiovascular risk. The main food groups that contributed more Na on weekends were processed meats and mixed dishes containing cereals (e.g. pasta, rice and pizza dishes). Substantial reductions in Na content of both staple foods and 'occasional' foods (convenience and takeaway foods which have high levels of Na) are required to achieve significant sustained reductions in population salt intake. This is particularly the case as eating out, takeaway and convenience foods appear to be becoming more of the 'usual' way of life in Australia, as well as other developed countries.

## Acknowledgements

**Acknowledgements:** The authors would like to acknowledge the contributions of the Victorian Department of Health and Human Services for provision of the VHM data and Baker IDI for collecting the initial VHM data. **Financial support:** The data collection and laboratory analyses were funded by an Australian Government National Health and Medical Research Council (NHMRC) partnership grant. Other partners included AWASH (Australian Division for World Action on Salt and Health), NSW Health, NSW Food Authority and the Australian Food and Grocery Council. J.W. is supported by a joint NHMRC Career Development Fellowship/Heart Foundation Future Leaders Fellowship (grant number 1082924) and undertakes additional work on salt reduction from the NHMRC, the WHO and VicHealth. C.G. is supported by an Alfred Deakin University postdoctoral fellowship. J.E.S. is supported by an NHMRC Research Fellowship. The funders had no role in the design, analysis or writing of this article. **Conflict of interest:** C.N. and C.G. are members of the World Action on Salt and Health (WASH) and AWASH Australian Action on Salt and Health; neither receives any financial support from these. C.N. has received remuneration from Meat and Livestock Australia, Nestlé Health Science and the Dairy Health Consortium as a consultant. These payments are unrelated to the submitted work. J.C. has received grants from the NHMRC of Australia as Chief Investigator for studies on dietary salt content and dietary salt reduction. J.W. is Director of the WHO Collaborating Centre on Population Salt Reduction. V.F. has received grants from Meat and Livestock Australia, Dairy Australia and the Grain and Legume Nutrition Council, for unrelated research work. **Authorship:** C.N. wrote the manuscript and oversaw the data analysis. V.F., M.W., M.-A.L., J.C., J.E.S. and J.W. contributed to the design and monitoring of the study. K.L. performed the data analysis and C.G. devised the data analysis approach together with C.N. C.G. took responsibility for the integrity of the data and the accuracy of the data analysis. All authors provided input into the final manuscript. **Ethics of human subject participation:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human participants were approved by the ethics committees of the Alfred Hospital (Project No. 421/10) and Deakin University (Project No. 2011-025). Written informed consent was obtained from all participants.

## References

- Goodall S, Gallego G & Norman R (2008) *Scenario Modelling of Potential Health Benefits Subsequent to the Introduction of the Proposed Standard for Nutrition, Health and Related Claims*. Sydney: Centre for Health Economics Research and Evaluation, University of Technology.
- World Health Organisation (2013) *Global Action Plan for the Prevention and Control of Noncommunicable Diseases 2013–2020*. Geneva: WHO.
- Australian Bureau of Statistics (2014) *Australian Health Survey: Nutrition First Results – Foods and Nutrients, 2011–12*. Catalogue no. 4364.0.55.007. Canberra: Commonwealth of Australia.
- Nowson C, Lim K, Grimes C *et al.* (2015) Dietary salt intake and discretionary salt use in two general population samples in Australia: 2011 and 2014. *Nutrients* **7**, 10501–10512.
- James PT, Ralph A & Sanchez-Castillo CP (1987) The dominance of salt in manufactured food in the sodium intake of affluent societies. *Lancet* **1**, 426–429.
- Mattes RD & Donnelly D (1991) Relative contributions of dietary sodium sources. *J Am Coll Nutr* **10**, 383–393.
- Harnack LJ, Cogswell ME, Shikany JM *et al.* (2017) Sources of sodium in US adults from 3 geographic regions. *Circulation* **135**, 1775–1783.
- Garcia J, Dunford EK, Sundstrom J *et al.* (2014) Changes in the sodium content of leading Australian fast-food products between 2009 and 2012. *Med J Aust* **200**, 340–344.
- Prentice CA, Smith C & McLean RM (2016) Sodium in commonly consumed fast foods in New Zealand: a public health opportunity. *Public Health Nutr* **19**, 958–966.
- National Health and Medical Research Council (2013) *Australian Dietary Guidelines*. Canberra: NHMRC.
- McCarthy S (2014) Weekly patterns, diet quality and energy balance. *Physiol Behav* **134**, 55–59.
- Australian Bureau of Statistics (1999) *National Nutrition Survey: Foods Eaten, Australia, 1995*. Catalogue no. 4804.0. Canberra: Commonwealth of Australia.
- An R (2016) Weekend–weekday differences in diet among US adults, 2003–2012. *Ann Epidemiol* **26**, 57–65.
- Department of Health (2013) *Victorian Health Monitor*. Melbourne: State Government of Victoria.
- Forni Ogna V, Ogna A, Vuistiner P *et al.* (2015) New anthropometry-based age- and sex-specific reference values for urinary 24-hour creatinine excretion based on the adult Swiss population. *BMC Med* **13**, 40.
- Land M-A, Webster J, Christoforou A *et al.* (2014) Salt intake assessed by 24-h urinary sodium excretion in a random and opportunistic sample in Australia. *BMJ Open* **4**, e003720.
- Conway JM, Ingwersen LA & Moshfegh AJ (2004) Accuracy of dietary recall using the USDA five-step multiple-pass method in men: an observational validation study. *J Am Diet Assoc* **104**, 595–603.
- Conway JM, Ingwersen LA, Vinyard BT *et al.* (2003) Effectiveness of the US Department of Agriculture 5-step multiple-pass method in assessing food intake in obese and nonobese women. *Am J Clin Nutr* **77**, 1171–1178.
- State Government of Victoria (2009) *4000 for Health Food Model Booklet*. Melbourne: Victorian Government Department of Human Services.
- Goldberg GR, Black AE, Jebb SA *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* **45**, 569–581.
- Schofield WN (1985) Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* **39**, Suppl. 1, 5–41.
- Food Standards Australia New Zealand (2013) AUSNUT 2007. <http://www.foodstandards.gov.au/science/monitoringnutrients/ausnut/Pages/ausnut2007.aspx> (accessed February 2015).
- Eyles H, Shields E, Webster J *et al.* (2016) Achieving the WHO sodium target: estimation of reductions required in the sodium content of packaged foods and other sources of dietary sodium. *Am J Clin Nutr* **104**, 470–479.
- Trevena H, Dunford E, Neal B *et al.* (2014) The Australian Food and Health Dialogue – the implications of the sodium

- recommendation for pasta sauces. *Public Health Nutr* **17**, 1647–1653.
25. Haines PS, Hama MY, Guilkey DK *et al.* (2003) Weekend eating in the United States is linked with greater energy, fat, and alcohol intake. *Obes Res* **11**, 945–949.
  26. An R (2016) Fast-food and full-service restaurant consumption and daily energy and nutrient intakes in US adults. *Eur J Clin Nutr* **70**, 97–103.
  27. Schmidt M, Affenito SG, Striegel-Moore R *et al.* (2005) Fast-food intake and diet quality in black and white girls: the National Heart, Lung, and Blood Institute Growth and Health Study. *Arch Pediatr Adolesc Med* **159**, 626–631.
  28. Hutton T (2002) Sodium technological functions of salt in the manufacturing of food and drink products. *Br Food J* **104**, 126–152.