

FINITE GROUPS WITH HEREDITARILY G -PERMUTABLE SCHMIDT SUBGROUPS

A. BALLESTER-BOLINCHES , S. F. KAMORNIKOV , V. PÉREZ-CALABUIG 
and V. N. TYUTYANOV

(Received 12 June 2023; accepted 29 June 2023; first published online 28 July 2023)

Abstract

A subgroup A of a group G is said to be hereditarily G -permutable with a subgroup B of G , if $AB^x = B^xA$ for some element $x \in \langle A, B \rangle$. A subgroup A of a group G is said to be hereditarily G -permutable in G if A is hereditarily G -permutable with every subgroup of G . In this paper, we investigate the structure of a finite group G with all its Schmidt subgroups hereditarily G -permutable.

2020 Mathematics subject classification: primary 20D10.

Keywords and phrases: finite group, Schmidt subgroup, hereditarily G -permutable subgroup.

1. Introduction

All groups considered in the paper are finite.

Recall that a group G is said to be a *minimal nonnilpotent group* or *Schmidt group* if G is not nilpotent and every proper subgroup of G is nilpotent. It is clear that every nonnilpotent group contains Schmidt subgroups, and their embedding has a strong structural impact (see, for example, [2, 3, 10]).

However, the following extensions of permutability turn out to be important in the structural study of groups and were introduced by Guo *et al.* in [6].

DEFINITION 1.1. Let A and B be subgroups of a group G .

- (1) A is said to be *G -permutable* with B if there exists some $g \in G$ such that $AB^g = B^gA$.
- (2) A is said to be *hereditarily G -permutable* with B (or *G - h -permutable* with B , for short) if there exists some $g \in \langle A, B \rangle$ such that $AB^g = B^gA$.

The second and third authors are supported by the BRFFR and RSF (Project F23RSF-237).

© The Author(s), 2023. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



- (3) A is said to be G -permutable in G if A is G -permutable with all subgroups of G .
 (4) A is said to be *hereditarily G -permutable* (or *G -h-permutable*, for short) in G if A is hereditarily G -permutable with all subgroups of G .

It is clear that permutability implies G -permutability but the converse does not hold in general as the Sylow 2-subgroups of the symmetric group of degree 3 show.

Our main goal here is to complete the structural study of groups in which every Schmidt subgroup of a group G is G -h-permutable. This study was started in [2] where we prove the following important fact.

THEOREM 1.2 [2, Theorem B]. *If every Schmidt subgroup of a group G is G -h-permutable in G , then G is soluble.*

Observe that the alternating group of degree 4 is a nonsupersoluble Schmidt group.

Let $p_1 > p_2 > \dots > p_r$ be the primes dividing $|G|$ and let P_i be a Sylow p_i -subgroup of G , for each $i = 1, 2, \dots, r$. Then we say that G is a *Sylow tower group of supersoluble type* if all subgroups $P_1, P_1P_2, \dots, P_1P_2 \dots P_{r-1}$ are normal in G . The class of all Sylow tower groups of supersoluble type is denoted by \mathfrak{D} .

Recall that if \mathfrak{F} is a nonempty class of groups and π is a set of primes, then \mathfrak{F}_π is the class of all π -groups in \mathfrak{F} . In particular, if p is a prime, then \mathfrak{R}_p is the class of all p -groups and $\mathfrak{D}_{\pi(p-1)}$ is the class of all Sylow tower groups G of supersoluble type such that every prime dividing $|G|$ also divides $p - 1$.

If G is a group, then $\text{Soc}(G)$ is the product of all minimal normal subgroups of G and $\Phi(G)$ is the Frattini subgroup of G , that is, the intersection of all maximal subgroups of G .

Our main goal here is to describe completely the groups G with trivial Frattini subgroup which have their Schmidt subgroups G -h-permutable.

THEOREM 1.3. *Let G be a group with $\Phi(G) = 1$. Assume that $\mathfrak{F} = LF(F)$ is the saturated formation locally defined by the canonical local definition F such that $F(p) = \mathfrak{R}_p \mathfrak{D}_{\pi(p-1)}$ for every prime p . If every Schmidt subgroup of G is G -h-permutable in G , then the following statements hold:*

- (1) $G = [\text{Soc}(G)]M$ is the semidirect product of $\text{Soc}(G)$ with an \mathfrak{F} -group M ;
- (2) if $\Phi(M) = 1$, then M is supersoluble.

We shall adhere to the notation and terminology of [1, 4].

2. Definitions and preliminary results

Our first lemma collects some basic properties of G -h-permutable subgroups which are very useful in induction arguments. Its proof is straightforward.

LEMMA 2.1. *Let A, B and K be subgroups of G with K normal in G . Then, the following statements hold.*

- (1) If A is G - h -permutable with B , then AK/K is G/K - h -permutable with BK/K in G/K .
- (2) If $K \subseteq A$, then A/K is G/K - h -permutable with BK/K in G/K if and only if A is G - h -permutable with B in G .
- (3) If A is G - h -permutable in G , then AK/K is G/K - h -permutable in G/K .
- (4) If $A \subseteq B$ and A is G - h -permutable in G , then A is B - h -permutable in B .

The following result describes the structure of Schmidt groups.

LEMMA 2.2 [5, 8]. *Let S be a Schmidt group. Then S satisfies the following properties:*

- (1) the order of S is divisible by exactly two prime numbers p and q ;
- (2) S is a semidirect product $S = [P]\langle a \rangle$, where P is a normal Sylow p -subgroup of S and $\langle a \rangle$ is a nonnormal Sylow q -subgroup of S and $\langle a^q \rangle \in Z(S)$;
- (3) P is the nilpotent residual of S , that is, the smallest normal subgroup of S with nilpotent quotient;
- (4) $P/\Phi(P)$ is a noncentral chief factor of S and $\Phi(P) = P' \subseteq Z(S)$;
- (5) $\Phi(S) = Z(S) = P' \times \langle a^q \rangle$;
- (6) $\Phi(P)$ is the centraliser $C_P(a)$ of a in P ;
- (7) if $Z(S) = 1$, then $|S| = p^m q$, where m is the order of p modulo q .

In what follows, $Sch(G)$ denotes the set of all Schmidt subgroups of a group G . Following [3], a Schmidt group with a normal Sylow p -subgroup will be called an $S_{(p,q)}$ -group.

The proof of Theorem 1.3 follows after a series of lemmas. They give us an interesting picture of the groups with supersoluble Schmidt subgroups.

LEMMA 2.3. *Let $\mathfrak{F} = \{H \mid Sch(H) \subseteq \mathfrak{U}\}$, where \mathfrak{U} is the class of all supersoluble groups. Then, \mathfrak{F} satisfies the following properties:*

- (1) if $G \in \mathfrak{F}$, then G is a Sylow tower group of supersoluble type; in particular, G is a soluble group;
- (2) \mathfrak{F} is a subgroup-closed saturated Fitting formation;
- (3) $\mathfrak{U} \subseteq \mathfrak{F}$;
- (4) $\mathfrak{F} = LF(F)$, where F is the canonical local definition such that $F(p) = \mathfrak{R}_p \mathfrak{D}_{\pi(p-1)}$ for every prime p .

PROOF. Statements (1), (2) and (3) follow from [7, Lemma 4 and Theorem 2].

Let $\mathfrak{H} = LF(F)$ be a local formation defined by the formation function F with $F(p) = \mathfrak{R}_p \mathfrak{D}_{\pi(p-1)}$ for every prime p . Assume that $\mathfrak{F} \not\subseteq \mathfrak{H}$. Let G be a group in $\mathfrak{F} \setminus \mathfrak{H}$ of minimal order. Since \mathfrak{F} is a saturated formation, it follows that G is a primitive soluble group. Let $N = Soc(G)$ be the unique minimal normal subgroup of G . Then $G/N \in \mathfrak{H}$. Since G is a Sylow tower group of supersoluble type and $C_G(N) = N$, we see that N is a Sylow p -subgroup of G , where p is the largest prime dividing $|G|$.

Let $q \in \pi(G)$ with $q \neq p$ and let Q be a Sylow q -subgroup of G . Since $N = C_G(N)$, it follows that PQ is not nilpotent. Hence, G has an $S_{(p,q)}$ -subgroup S , which is

supersoluble p -closed because $G \in \mathfrak{F}$. Then, by statements (4) and (5) of Lemma 2.2, $|S/Z(S)| = pq$ and therefore, by statement (7) of Lemma 2.2, q divides $p - 1$. Since G is a Sylow tower group of supersoluble type, it follows that

$$G/N = G/C_G(N) \in \mathfrak{D}_{\pi(p-1)},$$

and thus $G \in \mathfrak{S}$, which is a contradiction. Hence, $\mathfrak{F} \subseteq \mathfrak{S}$.

Assume that $\mathfrak{F} \neq \mathfrak{S}$, and let G be a group in $\mathfrak{S} \setminus \mathfrak{F}$ of minimal order. Since \mathfrak{S} is a saturated formation and $F(p)$ is a formation of soluble groups for all primes p , it follows that G is a primitive soluble group. Let N be a unique minimal normal subgroup of G . The choice of G yields $G \in \mathfrak{S}$ and $G/N \in \mathfrak{F}$. Since G is soluble, N is a p -group for some prime p , and from $G \in \mathfrak{S}$, it follows that

$$G/N = G/C_G(N) \in \mathfrak{R}_p \mathfrak{D}_{\pi(p-1)}.$$

We conclude that $G/N \in \mathfrak{D}_{\pi(p-1)}$ because $O_p(G/N) = 1$ by [4, Lemma A.13.6].

Let S be an $S_{(r,q)}$ -subgroup of G . If $r \neq p$, then S is contained in some Hall p' -subgroup H of G . Since $H \cong G/N \in \mathfrak{F}$, we see that $S \in \mathfrak{U}$. If $r = p$, then from $G/N \in \mathfrak{D}_{\pi(p-1)}$, it follows that q divides $p - 1$. Thus, by Lemma 2.2, $S \in \mathfrak{U}$. Consequently, every Schmidt subgroup of G is supersoluble, which is a contradiction. Hence, $\mathfrak{F} = \mathfrak{S}$. □

The following examples show that groups in Lemma 2.3 may not be supersoluble.

EXAMPLE 2.4. Let

$$Q = \langle a, b \mid a^4 = b^4 = 1, a^2 = b^2, b^{-1}ab = a^{-1} \rangle$$

be the quaternion group of order 8. Then G has a faithful and irreducible module A over the field of 5 elements of dimension 2. Let $G = [A]Q$ be the corresponding semidirect product. Then G is not supersoluble and $C = [A]\langle a \rangle$ and $D = [A]\langle b \rangle$ are supersoluble and normal subgroups of $G = CD$. By Lemma 2.3, $G \in \mathfrak{F} = \{H \mid \text{Sch}(H) \subseteq \mathfrak{U}\}$.

EXAMPLE 2.5. Assume that M is a nonabelian group of order 21. Then M has a faithful and irreducible module N over $\text{GF}(43)$, the field of 43 elements (see, for example, [4, Corollary B.11.8]). Consider the semidirect product $G = [N]M$. It is obvious that G is not supersoluble. By Lemma 2.3, $G \in \mathfrak{F} = \{H \mid \text{Sch}(H) \subseteq \mathfrak{U}\}$.

The following result is of interest although it is not needed for the proof of Theorem 1.3.

PROPOSITION 2.6. *Let $\mathfrak{F} = \{H \mid \text{Sch}(H) \subseteq \mathfrak{U}\}$. Then, for every $n \in \mathbb{N}$, there exists a group $G \in \mathfrak{F}$ of nilpotent length n .*

PROOF. Let $n \geq 2$ and let p_1, p_2, \dots, p_n be primes such that $p_1 < p_2 < \dots < p_n$ and p_i divides $p_j - 1$ for all $i < j$, where $i = 1, 2, \dots, n - 1, j = 2, \dots, n$. By Dirichlet's theorem, there exists an infinite set of primes of the form

$$p_1 p_2 \cdots p_{n_0} + 1,$$

where $n_0 \in \mathbb{N}$. Assume that p_{n+1} is one of them. It is obvious that p_i divides $p_{n+1} - 1$ for any $i = 1, 2, \dots, n$.

Assume that G_1 is a cyclic group of order p_1 . Assume that $i \geq 2$ and G_{i-1} is in \mathfrak{F} and of nilpotent length $i - 1$. By [4, Corollary B.11.8], G_{i-1} has a faithful and irreducible module V_{p_i} over the field of p_i elements. Let $G_i = [V_{p_i}]G_{i-1}$ be the corresponding semidirect product. Then $F(G_i) = V_{p_i}$ and hence the nilpotent length of G_i is equal to i . Furthermore, by Lemma 2.3, $G_i \in \mathfrak{F}$. In particular, G_n is an \mathfrak{F} -group of nilpotent length n . □

The following subgroup embedding property was introduced by Vasil’ev, Vasil’eva and Tyutyanov in [9].

DEFINITION 2.7. A subgroup H of a group G is said to be \mathbb{P} -subnormal in G if there exists a chain of subgroups

$$H = H_0 \subseteq H_1 \subseteq \dots \subseteq H_{n-1} \subseteq H_n = G$$

such that for every $i = 1, 2, \dots, n$, either $|H_i : H_{i-1}| \in \mathbb{P}$ or H_{i-1} is normal in H_i .

Note that \mathbb{P} -subnormality coincides with $\mathbb{K}\mathfrak{U}$ -subnormality (see [1, Ch. 6]) in the soluble universe.

LEMMA 2.8. Let A be a G -h-permutable subgroup of a soluble group G . Then, A is \mathbb{P} -subnormal in G . In particular, the supersoluble residual $A^{\mathfrak{U}}$ of A is subnormal in G .

PROOF. Let G be a group of smallest order for which the lemma is not true, and let L be a minimal normal subgroup of G . Since G is soluble, $|L| = p^n$ for some prime $p \in \pi(G)$ and $n \geq 1$. Suppose that $G = AL$. Then A is a maximal subgroup of G and $A \cap L = 1$. Let L_1 be a subgroup of prime order of L . Then, $AL_1^x = L_1^x A$ for some $x \in G$. Consequently, AL_1^x is a subgroup of G . Since A is maximal in G and $A \neq AL_1^x$, we see that $AL_1^x = G$. Because

$$|G : A| = |AL_1^x|/|A| = |L_1^x|/|A \cap L_1^x| = |L_1^x|,$$

we conclude that $|G : A| = p$ and then A is \mathbb{P} -subnormal in G , which is a contradiction. Hence, $G \neq AL$. Since $|AL| < |G|$, by Lemma 2.1, it follows that A is \mathbb{P} -subnormal in AL . By Lemma 2.1, AL/L is (G/L) -h-permutable in G/L , and from $|G/L| < |G|$, it follows that AL/L is \mathbb{P} -subnormal in G/L . In particular, AL is \mathbb{P} -subnormal in G by [1, Lemma 6.1.6]. However, then A is a \mathbb{P} -subnormal subgroup of G by [1, Lemma 6.1.7], which is a contradiction. Consequently, A is \mathbb{P} -subnormal in G . Applying [1, Lemma 6.1.9], we conclude that $A^{\mathfrak{U}}$ is subnormal in G . □

EXAMPLE 2.9. Let G be a group isomorphic to the alternating group of degree 6. Since G does not have maximal subgroups of prime index, the identity subgroup 1 of G is G -h-permutable but not \mathbb{P} -subnormal in G . Thus, the solubility of the group G in Lemma 2.8 is essential.

LEMMA 2.10. Let $G \in \mathfrak{F} = \{H \mid \text{Sch}(H) \subseteq \mathfrak{U}\}$. If $\Phi(G) = 1$ and every Schmidt subgroup of G is G -h-permutable in G , then G is supersoluble.

PROOF. We argue by induction on $|G|$. Let N be a minimal normal subgroup of G . Since G is soluble by Lemma 2.3, it follows that N is p -elementary abelian for some prime p . Since $\Phi(G) = 1$, it follows that $G = NM$ for some maximal subgroup M of G and $N \cap M = 1$.

Suppose that $NM_{p'}$ is p -nilpotent. Then $NM_{p'} \subseteq C_G(N)$. Then $G/C_G(N)$ is a p -group. Since $O_p(G/C_G(N)) = 1$ by [4, Lemma A.13.6], we have $N \subseteq Z(G)$. Then $G = N \times M$. Now, M belongs to \mathfrak{F} and $\Phi(M) \subseteq \Phi(G) = 1$ by [4, Theorem A.9.2]. By induction, M is supersoluble. Hence, G is supersoluble.

Assume that $NM_{p'}$ is not p -nilpotent. Consequently, $NM_{p'}$ contains a minimal non- p -nilpotent group X . By [1, Corollary 6.4.5], X is an $S_{\langle p,q \rangle}$ -subgroup $X = [P]Q$ and $P \subseteq N$. We can assume without loss of generality that $Q \subseteq M_{p'}$. Since the subgroup $[P]Q$ is G -h-permutable, we may assume that $([P]Q)M = PM$ is a subgroup of G . Consequently, $P = N$ and NQ is an $S_{\langle p,q \rangle}$ -subgroup G . By hypothesis, NQ is supersoluble. Hence, in view of Lemma 2.2, $|N/\Phi(N)| = p$ by Lemma 2.2. Since $\Phi(N) = 1$, it follows that $|N| = p$.

Consequently, we may assume that every minimal normal subgroup of G is cyclic. Then, by [4, Theorem A.10.6], $F(G)$ is a direct product of normal subgroups of G of prime order and so $G/C_G(F(G))$ is abelian. Since $C_G(F(G)) \subseteq F(G)$ by [4, Theorem A.10.6], it follows that $G/F(G)$ is abelian. In particular, G is supersoluble. \square

3. Proof of Theorem 1.3

Since G is soluble and $\Phi(G) = 1$, we conclude that $F(G) = \text{Soc}(G)$ and $G = [\text{Soc}(G)]M$ for some subgroup M of G , that is, $\text{Soc}(G) \cap M = 1$ by [4, Theorem A.10.6].

Let S be a Schmidt subgroup of M . Suppose that S is an $S_{\langle p,q \rangle}$ -subgroup. Then, by hypothesis, S is G -h-permutable in G . Consequently, by Lemma 2.8, S is \mathbb{P} -subnormal in G and S^{ll} is subnormal in G . In view of Lemma 2.2, we see that either $S^{\text{ll}} \neq 1$ is a p -subgroup of S or $S^{\text{ll}} = 1$. Assume that $S^{\text{ll}} \neq 1$. Then,

$$S^{\text{ll}} \subseteq F(G) \cap M = 1,$$

which is a contradiction. Therefore, every Schmidt subgroup of M is supersoluble.

By Lemma 2.3, it follows that $M \in \mathfrak{F} = LF(F)$, where F is the formation function given by $F(r) = \mathfrak{R}_r \mathfrak{D}_{\pi(r-1)}$ for any prime r .

By Lemma 2.10, M is supersoluble provided that $\Phi(M) = 1$.

References

- [1] A. Ballester-Bolinches and L. M. Ezquerro, *Classes of Finite Groups*, Mathematics and Its Applications, 584 (Springer, Dordrecht, 2006).
- [2] A. Ballester-Bolinches, S. F. Kamornikov, V. N. Tyutyayov and V. Pérez-Calabuig, ‘Finite groups with G -permutable Schmidt subgroups’, *Mediterr. J. Math.* **20**(174) (2023), 1–12.
- [3] A. Ballester-Bolinches, S. F. Kamornikov and X. Yi, ‘Finite groups with σ -subnormal Schmidt subgroups’, *Bull. Malays. Math. Sci. Soc.* **45** (2022), 2431–2440.

- [4] K. Doerk and T. Hawkes, *Finite Soluble Groups*, De Gruyter Expositions in Mathematics, 4 (Walter de Gruyter and Co., Berlin, 1992).
- [5] J. A. Gol'fand, 'On groups all of whose subgroups are special', *Doklady Akad. Nauk SSSR* **60** (1948), 1313–1315 (in Russian).
- [6] W. Guo, K. P. Shum and A. N. Skiba, 'X-quasinormal subgroups', *Sib. Math. J.* **48**(4) (2007), 593–605.
- [7] V. S. Monakhov, 'Finite groups with a given set of Schmidt subgroups', *Math. Notes* **58** (1995), 1183–1186.
- [8] O. J. Schmidt, 'Über Gruppen, deren sämtliche Teiler spezielle Gruppen sind', *Mat. Sb.* **31** (1924), 366–372.
- [9] A. F. Vasil'ev, T. I. Vasil'eva and V. N. Tyutyaynov, 'On the finite groups of supersoluble type', *Sib. Math. J.* **51**(6) (2010), 1004–1012.
- [10] V. A. Vedernikov, 'Finite groups with subnormal Schmidt subgroups', *Algebra Logic* **46**(6) (2007), 363–372.

A. BALLESTER-BOLINCHES, Department of Mathematics,
Guangdong University of Education,
Guangzhou, Guangdong 510303, PR China
e-mail: Adolfo.Ballester@uv.es

S. F. KAMORNIKOV,
Francisk Skorina State Gomel University,
Gomel, Belarus
e-mail: sfkamornikov@mail.ru

V. PÉREZ-CALABUIG, Departament de Matemàtiques,
Universitat de València, Dr. Moliner, 50, 46100 Burjassot,
València, Spain
e-mail: Vicent.Perez-Calabuig@uv.es

V. N. TYUTYANOV,
Gomel Branch of International University 'MITSO', Gomel,
Belarus
e-mail: vtutanov@gmail.com