

VOL. VIII.—1894.

THE
JOURNAL OF LARYNGOLOGY,
RHINOLOGY AND OTOTOLOGY;

AN ANALYTICAL RECORD OF CURRENT LITERATURE

RELATING TO

THE THROAT, NOSE AND EAR.

PUBLISHED MONTHLY.

London:

F. J. REBMAN, 11, ADAM STREET, STRAND.

AMERICAN AGENTS:

THE F. A. DAVIS COMPANY, 1914-16, CHERRY STREET, PHILADELPHIA, PA.

ENTERED AT STATIONERS HALL

THE JOURNAL OF LARYNGOLOGY, RHINOLOGY, AND OTOLOGY.

Founded in 1887 by MORELL MACKENZIE *and* NORRIS WOLFENDEN.

EDITED BY

R. NORRIS WOLFENDEN, M.D. *Cantab.*

JOHN MACINTYRE, M.B. *Glasgow.*

DUNDAS GRANT, M.A., M.D., F.R.C.S. *Eng.*

WITH THE ASSISTANCE OF

DR. SANDFORD (*Cork*), DR. MILLIGAN (*Manchester*),

AND WITH THE CO-OPERATION OF

DRS. BARON (*Bristol*), R. BOTEY (*Barcelona*),

CARTAZ (*Paris*), BRYSON DELAVAN (*New York*),

DRAISPUL (*St. Petersburg*), GRAZZI (*Florence*), GUYE (*Amsterdam*),

HICGUET (*Brussels*), MIDDLEMASS HUNT (*Liverpool*),

VALERIUS IDELSON (*Berne*), JOAL (*Paris*), A. B. KELLY (*Glasgow*),

KARWOWSKI (*Warsaw*), LACOCARRET (*Toulouse*), LAKE (*London*),

LICHTWITZ (*Bordeaux*), JOHN N. MACKENZIE (*Baltimore*),

G. W. MAJOR (*Montreal*), PROF. MASSEI (*Naples*),

MEYJES (*Amsterdam*), E. MEYER (*Berlin*), MICHAEL (*Hamburg*),

HOLGER MYGIND (*Copenhagen*), W. PORTER (*St. Louis*),

ROBERTSON (*Newcastle*), SAJOUS (*Philadelphia*), SENDZIAK (*Warsaw*),

SOTA (*Seville*), AND ST. CLAIR THOMSON (*London*).

CONTENTS.

Association Reports:—

International Medical Congress—(continued).

Pachydermia Laryngis. Dr. A. DAMIENO.	Page 383
Syphilo-tuberculosis of the Larynx. Dr. A. DAMIENO.	384
Treatment of Nasal Hemorrhage. Dr. I. DIONISIO.	385
Erysipelas of the Larynx. Dr. FASANO.	387
Ventriloquism. Drs. FLATAU and GUTZMANN.	389
Croup and Croupous Diseases treated by Pilocarpin. Dr. SZIKLAI.	390
Latee Empyema of the Maxillary Sinus. Dr. GRADEWIGNO.	397
Caustic Treatment in the Removal of Carious Ossicles. Dr. FERRERI.	399
Magnifying Aural Speculum. Dr. DUNDAS GRANT.	404
Aural Forceps. Dr. DUNDAS GRANT.	405
The Physiology of the Middle Ear. Dr. BRACCHI.	405
Auscultation of Noises in the Ear. Dr. MADEUF.	409
Treatment of Purulent Otitis in the Scrofulous. Dr. ISAIA.	410
Deaf-Mutism up to Reception in Institutions. Dr. FLATAU.	412
Treatment of Deafness due to Sclerosis. Dr. BOSIO.	414
New Auditory Intermitent Paracusis. Dr. COZZOLINO.	415

American Laryngological Association—

Presidential Address.	417
Nasal Polypus and Etmoiditis. Dr. CASSELDERRY.	419
Papillary Hypertrophy of Nasal Mucous Membrane. Dr. WRIGHT.	423
Metallic Electrodes in Nasal and Post-Nasal Disease. Dr. RICE.	424
Results of Cutting Operations on the Nasal Septum. Dr. FRENCH.	425
Early Operation in Diseases of the Antrum. Dr. DALY.	427
Compound Follicular Odontoma involving the Antrum. Dr. DE ROALDES.	428
Extraction of a Cockle Burr from the Larynx. Dr. SHIELDS.	429
Cases of Laryngeal Neoplasm. Dr. C. H. KNIGHT.	430
Cases of Laryngeal Malignant Neoplasm. Dr. BLISS.	430
Case of Laryngeal Omy by a New Method. Dr. SWAIN.	430
Exudative Pharyngitis. Dr. GLASGOW.	431
Singers' Nodes. Dr. KNIGHT.	432
Sarcoma of the Tonsil. Dr. WATSON.	434
Early Diagnosis of Malignant Tumours. Dr. GLEITSMANN.	434
Neurasthenic Throats. Dr. P. POUCHER.	436
Clinic Reports. Dr. F. INGALS.	436
Etiology of Rheumatic Affectations due to Tonsillar Diseases. Dr. WAGNER.	437

Laryngological Society of London—

Lupus of Nose and Larynx. Dr. GRANT.	439
Probable Epithelioma of Larynx. Dr. GRANT.	440
Paralysis of Thyro-Arytenoid Muscle. Dr. GRANT.	440
Removal of Thyroid for Graves' Disease. Mr. LAKE.	441
Lupus of Pharynx and Larynx. Dr. SEMON.	441
Doubtful Malignant Disease of Larynx. Dr. SEMON.	443
Sequel to Obscure Ulceration of Pharynx in arrested Tubercular Disease. } Dr. SEMON.	444
Fixation of Left Vocal Cord. Dr. SPICER.	445
Unilateral Laryngitis. Mr. STEWART.	445
Carcinoma Laryngis. Mr. STEWART.	445
Congenital Abnormality of Larynx. Dr. BALL.	446
Intra-Laryngeal Syringe. Dr. W. WILLIAMS.	446
Sarcoma of Nose and Tonsils. Dr. BOND.	446
Epithelioma of Cervical Glands. Dr. BOND.	447

Laryngological Society of Berlin—

Creolin in Phthisis. Dr. SCHEIDEWALDT.	447
Tussis Nervosa. Dr. ROSENBERG.	447
Vascular Polypi. Dr. HEINMANN.	447
Septal Polypi. Dr. ALEXANDER.	447
Gummata of Turbinate. Dr. SCHEINMANN.	448
Cleft Palate. Dr. HERZFELD.	448
Malignant Growth of the Nose. Dr. HATAN.	449
Diphtheria and Lacunar Tonsillitis. Dr. MEYER.	449
Vascular Polypi. Dr. SCHEIER.	449
Suppuration of Sphenoidal Sinus. Dr. HERZFELD.	449

Abstracts.—Diphtheria.

Mouth, Pharynx, &c.	449
Nose and Naso-Pharynx.	454
Larynx.	456
Thyroid.	458
Ears.	463

Reviews. — GOLDSCHMIDT: Tuberculosis.

BRESGEN: Headache in Nasal Disease.	467
ROSENBERG: Diseases of the Throat and Larynx.	464

Notes.

	468
	469

GUIDE TO ADVERTISEMENTS.

	PAGE		PAGE
Allen & Hanburys	iv., v.	Godfrey & Cooke	x.
Bullock & Co.	xiii.	Ingram & Royle	ix.
Burroughs, Wellcome & Co.	vi., viii., xiv.	Lambert Pharmacal Co.	ii.
Central London Throat Hospital	xv.	Kebman	x., xi., xii., xiii.

Excellence in Pharmacy.

HAZELINE.

THE following Report by Dr. SHOEMAKER, an eminent United States authority taken from his work "Materia Medica and Therapeutics," thoroughly describes the distinct utility of this product, and is of special interest, since it emanates from the country where the Hazeline is distilled:—

"HAZELINE is the name of an excellent preparation made by BURROUGHS WELLCOME & CO., of London, from the twigs of Witch Hazel. The active principles of the plant are contained in Hazeline, which is a useful embrocation in cases of contusions, sprains, and chronic rheumatism. It is applied upon absorbent cotton for the relief of hæmorrhoids, and to allay the pain and smarting of burns, eczema, and erysipelas. Diluted with water, it may be beneficially injected into the rectum for internal hæmorrhoids. As a gargle or inhalation it is useful in coryza, ozæna, and pharyngitis. Hazeline is highly commended by the British Medical Association's Committee on Therapeutics. Hazeline, being prepared from the fresh green twigs, contains all the valuable volatile principles of the plant Witch Hazel, and is much more uniform and reliable in its action than are the tinctures, fluid extracts, &c., prepared from the dry bark."

Hazeline supplied to the Medical Profession by all Chemists in $\frac{1}{4}$ -lb. and 1-lb. bottles, at 1s. 2d. and 3s. 6d. each.

"EUCALYPTIA."

"EUCALYPTIA" is extensively employed for inhalation, and acts as a stimulating expectorant. It may be inhaled in combination with Ammonium Chloride vapour from the Vereker Chloride of Ammonium Inhaler, or dry with the "Pinol Eucalyptia" Dry Inhaler or Menthol Pocket Inhaler. Such inhalations afford excellent relief in catarrh, ozæna, cold in the head, sore throat, aphonia, asthma, bronchitis, and phthisis. A good method for using "Eucalyptia" in steam inhalations is to drop 30 minims of the oil on the sponge of a Wallich's Improved Inhaler, place in the jug a pint of boiling water, and inhale from the mouthpiece. The steam is drawn through the sponge without difficulty and in passing through becomes thoroughly saturated with the "Eucalyptia," which is conveyed into the lungs and air-passages.

"Eucalyptia" supplied to the Medical Profession by all Chemists in 2-oz. and 1-lb. bottles, at 1s. 3d. and 6s. each.

PINOL

Is the pure essential oil distilled from the needles of the Pinus Pumilio. It possesses superior antiseptic properties, and its agreeable odour clearly distinguishes it from the class of ordinary pine oils. Its ozonising properties have been recognised in its use and effect in dry inhalations in coryza, hay fever, quinsy, aphonia, and phthisis and it has been successfully used in oily atomisations in connection with Paroleine. On account of its pleasant odour it is one of the most agreeable of disinfectants for sprinkling about the sick room.

Supplied to the Medical Profession in $\frac{1}{2}$ -oz. and 1-oz. bottles, at 1s. 3d. and 2s. each.

PURE TEREBENE (B., W. & CO.)

A PURE Terebene, free from all irritating admixtures, and possessing an agreeable odour of Pine, prepared by repeated distillation of pure oil of turpentine in the presence of sulphuric acid. Terebene (B., W. & Co.) is a powerful expectorant; it has been used with much success in asthma, and even as a palliative in phthisis. Terebene, for pulmonary affections, may be inhaled from a dry inhaler, such as the "Pinol Eucalyptia" Dry Inhaler, or when desired may be inhaled through Wallich's Improved Steam Inhaler.

Supplied to the Medical Profession in 1-oz., 2-oz., 8-oz., and 16-oz. bottles, at 9d., 1s., 2s., and 3s. 9d. each.

SPECIMENS SENT TO MEDICAL MEN FREE ON REQUEST.

BURROUGHS, WELLCOME & CO., Manufacturing Chemists.
SNOW HILL BUILDINGS, LONDON.

Cable and Telegraphic Address—"BURCOME, LONDON."