

# Hyperlipidaemia and its risk factors in the Guangxi Bai Ku Yao and Han populations

Yin Ruixing<sup>1,\*</sup>, Yang Dezhai<sup>2</sup>, Li Shuquan<sup>2</sup>, Chen Yuming<sup>1</sup>, Yang Hanjun<sup>3</sup>,  
Feng Qiming<sup>4</sup>, Pan Shangling<sup>5</sup>, Lin Weixiong<sup>2</sup>, Tan Jing<sup>6</sup> and Li Yiyang<sup>1</sup>

<sup>1</sup>Department of Cardiology, Institute of Cardiovascular Diseases, The First Affiliated Hospital, Guangxi Medical University, 22 Shuangyong Road, Nanning 530021, Guangxi, People's Republic of China: <sup>2</sup>Department of Molecular Biology, Medical Scientific Research Center, Guangxi Medical University, Nanning, People's Republic of China: <sup>3</sup>The Health Bureau of Nandan County, Guangxi Zhuang Autonomous Region, Hechi, People's Republic of China: <sup>4</sup>Department of Health Statistics, Public Health School, Guangxi Medical University, Nanning, People's Republic of China: <sup>5</sup>Department of Pathophysiology, School of Premedical Sciences, Guangxi Medical University, Nanning, People's Republic of China: <sup>6</sup>Department of Internal Medicine, Nandan County People's Hospital, Guangxi Zhuang Autonomous Region, Hechi, People's Republic of China

Submitted 20 November 2007: Accepted 19 May 2008: First published online 23 July 2008

## Abstract

**Objective:** To compare the differences in hyperlipidaemia prevalence and its risk factors between the Guangxi Bai Ku Yao and Han populations.

**Design:** Cross-sectional study of hyperlipidaemia.

**Setting:** Both populations were from Lihu and Baxu villages in Nandan County, Guangxi Zhuang Autonomous Region, People's Republic of China.

**Subjects:** A total of 1170 healthy subjects of Bai Ku Yao and 1173 participants of Han Chinese aged 15–89 years were surveyed by a stratified randomized cluster sampling. Information on demographic, dietary and lifestyle characteristics was collected by standard questionnaires. Blood pressure, height, weight, waist circumference, serum lipids and apolipoproteins were measured, and BMI (kg/m<sup>2</sup>) was calculated as weight divided by the square of height.

**Results:** The prevalence rates of hypercholesterolaemia, hypertriglyceridaemia and hyperlipidaemia in Bai Ku Yao and Han were 12.4% *v.* 26.2% ( $P < 0.001$ ), 15.0% *v.* 14.8% ( $P > 0.05$ ) and 24.4% *v.* 33.9% ( $P < 0.001$ ), respectively. Hyperlipidaemia was positively correlated with BMI, waist circumference, total energy and total fat intakes, and negatively associated with physical activity and total dietary fibre intake in Bai Ku Yao ( $P < 0.05$  to 0.001). Hyperlipidaemia was positively associated with age, alcohol consumption, BMI, waist circumference, total energy and total fat intakes, and inversely correlated with physical activity and total dietary fibre intake in Han ( $P < 0.05$  to 0.001).

**Conclusions:** The prevalence of hypercholesterolaemia and hyperlipidaemia was significantly lower in the Bai Ku Yao than in the Han population, which might result from different dietary habits, lifestyle choices and physical activity level, as well as genetic factors between the two ethnic groups.

**Keywords**  
Lipids  
Apolipoproteins  
Hyperlipidaemia  
Prevalence  
Risk factors

Coronary artery disease has a multifactorial origin, including hereditary and acquired risk factors which may be the direct cause of the disease or merely associated with it<sup>(1,2)</sup>. Disorders in lipid metabolism such as high levels of plasma or serum total cholesterol (TC)<sup>(3)</sup>, TAG<sup>(4)</sup>, LDL cholesterol (LDL-C)<sup>(5)</sup> and apo B<sup>(6)</sup>, and low levels of HDL cholesterol (HDL-C)<sup>(7)</sup>, play a relevant role in the progression of atherosclerosis<sup>(8)</sup>; thus the laboratory assessment of lipids is of fundamental importance to diagnose, treat and prevent this condition<sup>(9)</sup>. Although a great deal of research effort has been focused on the determinants of these lipid phenotypes in different ethnic

groups and diseases<sup>(10–12)</sup>, very few data are available about the prevalence of hyperlipidaemia and its risk factors in isolated minority groups.

There are fifty-six ethnic groups in China. Han is the largest group and Yao is the eleventh largest minority among the fifty-five minority groups according to population size. Bai Ku Yao (White-trouser Yao), an isolated subgroup of the Yao minority, is so named because all of the men wear white knee-length knickerbockers. The population size is about 30 000. Because of their isolation from other ethnic groups, special customs and cultures including their clothing, intra-ethnic marriages, dietary

habits, and corn wine and rum intakes are still completely preserved to the present day. Little is known about the prevalence of hyperlipidaemia and its risk factors in this population. Therefore, the present study was undertaken to compare the differences in hyperlipidaemia prevalence and its risk factors between the Guangxi Bai Ku Yao and Han populations.

## Methods

### Subjects

The subjects of Bai Ku Yao residing in Lihu and Baxu villages in Nandan County, Guangxi Zhuang Autonomous Region, People's Republic of China, were surveyed by a stratified randomized cluster sampling. First, teams (resident communities; inhabitants) were selected randomly from the two villages. Then, sexes in each team were separated for the survey. Finally, the sampled resident sections were determined from the local population registers. A total of 1230 subjects from sixteen teams were asked to participate in the study and 1193 subjects actually participated. The response rate was 97.0%. Twenty-three persons (1.9%) with a history or evidence of other diseases were excluded from the analysis. A total of 1170 subjects were included in the present study. The ages of the subjects ranged from 15 to 85 years, with an average age of 37.5 (SD 16.7) years. There were 588 males (50.3%) and 582 females (49.7%). All subjects were peasants. The age distribution was as follows: <20 years, 185 people (15.8%); 20–29 years, 261 people (22.3%); 30–39 years, 239 people (20.4%); 40–49 years, 210 people (18.0%); 50–59 years, 139 people (11.9%); 60–69 years, eighty-three people (7.1%);  $\geq 70$  years, fifty-three people (4.5%). The subjects accounted for 3.9% of the total Bai Ku Yao population.

During the same period, a total of 1235 subjects of Han Chinese from fifteen teams were asked to participate in the study by the same method and 1205 subjects completed the survey. The response rate was 97.6%. Thirty-two people (2.7%) with a history or evidence of other diseases were excluded. A total of 1173 subjects were included in the present study. The mean age of the subjects was 38.3 (SD 16.8) years (range 15 to 89 years). There were 589 men (50.2%) and 584 women (49.8%). All of them were also peasants. The age distribution was as follows: <20 years, 180 persons (15.3%); 20–29 years, 241 persons (20.5%); 30–39 years, 251 persons (21.4%); 40–49 years, 208 persons (17.7%); 50–59 years, 139 persons (11.8%); 60–69 years, eighty-five persons (7.2%);  $\geq 70$  years, sixty-nine persons (5.9%).

All study subjects were essentially healthy and had no evidence of any chronic illness, including hepatic, renal or thyroid disease, heart attack or myocardial infarction, stroke, congestive heart failure or diabetes mellitus. They were not taking medications known to affect serum lipid

levels (lipid-lowering drugs such as statins or fibrates, beta-blockers, diuretics or hormones). The present study was approved by the Ethics Committee of the First Affiliated Hospital, Guangxi Medical University. Informed consent was obtained from all subjects after they received a full explanation of the study.

### Epidemiological survey

The survey was carried out using internationally standardized methods, following a common protocol<sup>(13)</sup>. Information on demographics (age, gender, residential area), socio-economic status (education level achieved, marital status, annual household income), cigarette smoking, alcohol consumption and physical activity was collected with standardized questionnaires. The 24 h dietary recall method was used to determine the dietary intakes of each subject<sup>(14)</sup>. Detailed descriptions of all foods, beverages and supplements consumed during the 24 h period before the interview, including the quantity, cooking method and brand names, were recorded by a chief physician. The interviewer used food models and pictures depicting portion sizes and followed a standardized protocol for determining the weight of food consumed. The intakes of macronutrients from the ingredients were determined by using the 2002 Chinese Food Composition Table<sup>(15)</sup>. Although a 24 h dietary recall may be inaccurate when diets are highly variable, the Bai Ku Yao diet is consistent throughout the year and among individuals because of the Bai Ku Yao's reliance on a limited number of locally available food items. Overall physical activity was ascertained with the use of a modified version of the Harvard Alumni Physical Activity Questionnaire<sup>(16)</sup>, which included questions about the number of hours per day (mean of a regular weekday and a regular weekend day) spent sleeping and in sedentary, light, moderate and vigorous activities; the interviewer ensured that the total time added up to 24 h. The alcohol information included questions about the number of grams of rice wine, wine, beer or liquor consumed during the preceding 12 months. At the physical examination, several parameters including height, weight and waist circumference were measured. Sitting blood pressure was measured three times with the use of a mercury sphygmomanometer after a rest of at least 15 min, and the average of the three measurements was used for the level of blood pressure. Systolic blood pressure was determined by the first Korotkoff sound and diastolic blood pressure by the fifth Korotkoff sound. Body weight, to the nearest 50 g, was measured using a portable balance scale. Subjects were weighed without shoes and in a minimum of clothing. Height was measured, to the nearest 0.5 cm, using a portable steel measuring device. From these two measurements BMI (kg/m<sup>2</sup>) was calculated. Waist circumference was measured with a non-stretchable measuring tape, at the level of the smallest area of the waist, to the nearest 0.1 cm.

### **Measurements of lipids and apolipoproteins**

A venous blood sample was drawn from an antecubital vein in all subjects after an overnight fast. The blood was transferred into glass tubes and allowed to clot at room temperature. Immediately following clotting serum was separated by centrifugation for 15 min at 3000 rpm. The levels of TC, TAG, HDL-C and LDL-C in samples were determined enzymatically using commercially available kits: Tcho-1, TG-LH (RANDOX Laboratories Ltd, Crumlin, UK), Cholestest N HDL and Cholestest LDL (Daiichi Pure Chemicals Co., Ltd, Tokyo, Japan), respectively. Serum apo A1 and apo B levels were measured by an immunoturbidimetric assay (RANDOX Laboratories Ltd). All determinations were performed with an autoanalyser (type 7170A; Hitachi Ltd, Tokyo, Japan) in the Clinical Science Experiment Center of the First Affiliated Hospital, Guangxi Medical University<sup>(17,18)</sup>.

### **Diagnostic criteria**

The normal values of serum TC (mmol/l), TAG (mmol/l), HDL-C (mmol/l), LDL-C (mmol/l), apo A1 (g/l), apo B (g/l) and the ratio apo A1:apo B in the Clinical Science Experiment Center were 3.10–5.17, 0.56–1.70, 0.91–1.81, 2.70–3.20, 1.00–1.78, 0.63–1.14 and 1.00–2.50, respectively. The individuals with TC > 5.17 mmol/l and/or TAG > 1.70 mmol/l were defined as hyperlipidaemic<sup>(17,18)</sup>. Hypertension was diagnosed according to the criteria of the 1999 WHO/International Society of Hypertension guidelines for the management of hypertension<sup>(19,20)</sup>. The diagnostic criteria of overweight and obesity were according to the Cooperative Meta-analysis Group of China Obesity Task Force. Normal weight, overweight and obesity were defined as BMI < 24, 24–28 and > 28 kg/m<sup>2</sup>, respectively<sup>(21)</sup>.

### **Statistical analysis**

The data were organized and analysed using Excel XP (Microsoft, Seattle, WA, USA) and the SPSS for Windows statistical software package version 11.5.0 (SPSS Inc., Chicago, IL, USA). Means and standard deviations as well as frequency distributions of participant characteristics were calculated. The difference in mean values of parameters between Bai Ku Yao and Han was tested by Student's unpaired *t* test. The difference in percentage values was tested by the  $\chi^2$  test. The influences of sex, age, physical activity, BMI, waist circumference, alcohol consumption and cigarette smoking were adjusted for in the statistical analyses. In order to evaluate the risk factors for hyperlipidaemia, unconditional logistic regression analysis was also performed in the combined population of Bai Ku Yao and Han, the Bai Ku Yao and the Han, respectively. The backward multiple logistic regression method was used to select the risk factors significantly associated with hyperlipidaemia. Total intake of each nutrient was summed over all foods consumed. The Matlab 5.0 software (The MathWorks, Inc.,

Natick, MA, USA) was used for processing these procedures by the method of multiplication of matrix<sup>(22)</sup>. A *P* value of less than 0.05 was considered statistically significant.

## **Results**

### **Comparison of general characteristics between Bai Ku Yao and Han**

The demographic, dietary and other lifestyle characteristics of Bai Ku Yao and Han are shown in Table 1. The level of physical activity and the intakes of carbohydrate, vegetal protein and total dietary fibre were higher in Bai Ku Yao than in Han ( $P < 0.001$  for all), whereas educational level, height, weight, BMI, waist circumference, blood pressure levels including systolic, diastolic and pulse pressure, hypertension and the intakes of total energy, total fat, total protein, dietary cholesterol and salt were higher in Han than in Bai Ku Yao ( $P < 0.05$  to  $0.001$ ). In addition, there were also significant differences in the staple and subsidiary foods and drinks between the two ethnic groups. For the great majority of Bai Ku Yao people, corn (gruel or tortillas) was the staple food and rice, soya, buckwheat, sweet potato and pumpkin products were the subsidiary foods all the year round. Approximately 90% of the beverages comprised corn wine and rum that they brewed themselves, the alcohol content is about 15% (v/v). The subjects of Bai Ku Yao were also accustomed to drinking hempseed soup. In contrast, rice was the staple food and corn, broomcorn, potato and taro products were the subsidiary foods in Han Chinese. About 90% of the beverages comprised rice wine with alcohol content of about 30% (v/v). There were no significant differences in the age structure, the percentages of subjects who consumed alcohol or smoked cigarettes, or the ratio of men to women between the two ethnic groups ( $P > 0.05$ ).

### **Prevalence of hyperlipidaemia in Bai Ku Yao and Han**

As shown in Table 2, the prevalence of high TC (hypercholesterolaemia), LDL-C, apo B and apo A1:apo B ratio was lower in Bai Ku Yao than in Han ( $P < 0.05$  to  $0.001$ ), whereas the prevalence of low HDL-C and apo A1 was higher in Bai Ku Yao than in Han ( $P < 0.001$  for each). There was no significant difference in the prevalence of high TAG (hypertriacylglycerolaemia) between the two ethnic groups ( $P > 0.05$ ). In the subjects with hypercholesterolaemia and hypertriacylglycerolaemia, there were thirty-six subjects in Bai Ku Yao (3.1%) and eighty-three subjects in Han (7.1%;  $\chi^2 = 19.430$ ,  $P < 0.001$ ) with both high TC and TAG. Thus, the prevalence of hyperlipidaemia in Bai Ku Yao and Han was 24.4% (285/1170) *v.* 33.9% (398/1173;  $\chi^2 = 25.981$ ,  $P < 0.001$ ).

**Table 1** Comparison of demographic, dietary and lifestyle characteristics between the Bai Ku Yao and Han populations

Characteristic	Bai Ku Yao (n 1170)		Han Chinese (n 1173)		t ( $\chi^2$ )	P
	Mean or n	SD or %	Mean or n	SD or %		
Age (years)	37.5	16.7	38.3	16.8	1.120	0.263
Male/female	588/582		589/584		0.000	1.000
Education level (years)	4.4	3.7	6.5	3.7	13.492	0.000
Physical activity (h/week)	48.8	14.5	46.4	13.5	4.151	0.000
Height (cm)	152.7	7.2	155.0	8.1	7.297	0.000
Weight (kg)	50.5	7.1	53.3	8.8	8.430	0.000
BMI (kg/m <sup>2</sup> )	21.7	2.4	22.2	2.9	4.574	0.000
>24 kg/m <sup>2</sup> [n and %]	178	15.2	246	21.0	13.104	0.000
Waist circumference (cm)	70.9	7.6	73.1	9.2	6.183	0.000
Systolic blood pressure (mmHg)	115.7	16.3	120.0	16.5	6.336	0.000
Diastolic blood pressure (mmHg)	74.1	9.4	75.9	10.4	4.278	0.000
Pulse pressure (mmHg)	41.6	12.0	44.2	11.2	5.403	0.000
Hypertension [n and %]	127	10.9	193	16.5	15.571	0.000
Alcohol consumption [n and %]	396	33.9	454	38.7	5.980	0.014
Cigarette smoking [n and %]	294	25.1	347	29.6	5.847	0.016
Energy (MJ/d)	8.85	0.37	8.99	0.39	8.761	0.000
Energy (kcal/d)	2114	88	2148	93	8.761	0.000
Fat (% of energy)	9.7	2.6	12.5	3.9	20.441	0.000
Carbohydrate (% of energy)	77.9	8.3	72.3	7.6	17.033	0.000
Protein (% of energy)	8.6	2.5	10.3	3.8	12.789	0.000
Alcohol (% of energy)	3.8	1.6	4.9	2.1	14.259	0.000
Carbohydrate (g/d)	412.0	17.2	388.3	15.3	35.255	0.000
Protein (g/d)	45.5	5.7	55.3	7.8	34.850	0.000
Animal (%)	26.8	3.2	34.6	4.3	49.799	0.000
Vegetal (%)	73.2	7.2	65.4	6.7	27.146	0.000
Total fat (g/d)	22.8	3.7	29.8	4.2	43.129	0.000
SFA (g/d)	5.7	2.2	9.2	3.3	30.735	0.000
MUFA (g/d)	7.2	2.9	9.7	5.6	13.571	0.000
PUFA (g/d)	9.9	3.9	10.9	4.7	5.810	0.000
Dietary cholesterol (mg/d)	179.4	102.8	198.8	108.5	4.427	0.000
Total dietary fibre (g/d)	9.4	3.1	7.8	2.7	13.582	0.000
Na intake (g/d)	7.2	2.4	7.8	2.6	5.898	0.000

### Associations between general characteristics and hyperlipidaemia in Bai Ku Yao and Han

The associations of sex, BMI, hypertension, alcohol consumption, cigarette smoking and age with hyperlipidaemia in Bai Ku Yao and Han are also shown in Table 2. The prevalence of high TAG in both ethnic groups, high apo A1:apo B ratio in Bai Ku Yao, and low HDL-C and apo A1 in Bai Ku Yao was higher in males than females. The prevalence of high TC, TAG, LDL-C and apo B in both ethnic groups was higher in subjects with BMI > 24 kg/m<sup>2</sup> than those with BMI ≤ 24 kg/m<sup>2</sup>. The prevalence of high TC, TAG and apo B in both ethnic groups was higher in drinkers than in non-drinkers. The prevalence of high apo B and apo A1:apo B ratio in Bai Ku Yao and high TAG in Han was higher in smokers than in non-smokers. There was also a significant difference in the prevalence of high TC, LDL-C and apo B, and of low HDL-C and apo A1 between age subgroups in both ethnic groups.

### Risk factors for hyperlipidaemia in Bai Ku Yao and Han

The results of multiple logistic regression analysis are shown in Table 3. Hyperlipidaemia was positively correlated with BMI, waist circumference, total energy and total fat intakes, and negatively associated with physical

activity and total dietary fibre intake, in Bai Ku Yao ( $P < 0.05$  to 0.001). Hyperlipidaemia was positively associated with age, alcohol consumption, BMI, waist circumference, total energy and total fat intakes, and inversely correlated with physical activity and total dietary fibre intake, in Han ( $P < 0.05$  to 0.001). There was no significant correlation between hyperlipidaemia and sex, education level, dietary cholesterol, cigarette smoking and hypertension in both ethnic groups ( $P > 0.05$ ).

### Comparison of hyperlipidaemia between Libu and Baxu villages

The prevalence of hypertriglycerolaemia, low HDL-C and apo A1 in Bai Ku Yao was higher but the prevalence of hypercholesterolaemia was lower in subjects from Lihu village than from Baxu villages ( $P < 0.05$  to 0.001). There was no significant difference in the prevalence of high apo B, apo A1:apo B ratio and hyperlipidaemia between the two villages ( $P > 0.05$ , Table 4).

### Discussion

The present study shows that the prevalence of hypercholesterolaemia and hyperlipidaemia was lower in

**Table 2** Associations of demographic, dietary and lifestyle characteristics with the prevalence of hyperlipidaemia in the Bai Ku Yao and Han populations

Characteristic	TC > 5.17 mmol/l		TAG > 1.70 mmol/l		HDL-C < 1.00 mmol/l		LDL-C > 3.20 mmol/l		Apo A1 < 1.0 g/l		Apo B > 1.14 g/l		Apo A1:apo B > 2.50	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Bai Ku Yao	1170	12.4	176	15.0	61	5.2	151	12.9	267	22.8	68	5.8	84	7.2
Men	588	12.6	101	17.2	42	7.1	68	11.6	167	28.4	37	6.3	61	10.4
Women	582	12.2	75*	12.9	19**	3.3	83	14.3	100***	17.2	31	5.3	23***	4.0
BMI ≤ 24 kg/m <sup>2</sup>	992	10.0	130	13.1	58	5.9	109	11.0	247	24.9	49	4.9	73	7.4
BMI > 24 kg/m <sup>2</sup>	178	46***	46***	25.8	3*	1.7	42***	23.6	20***	11.2	19**	10.7	11	6.2
Normotensives	1043	11.8	14.4	15.0	58	5.7	123	11.8	256	24.5	57	5.5	70	6.7
Hypertensives	127	27***	21.3	20.5	3	2.4	28**	22.1	11***	8.7	11	4.7	14	11.0
Non-drinkers	774	7.2	9.3	12.7	54	7.0	90	11.6	226	29.2	36	4.7	27	3.5
Drinkers	396	73***	18.4	19.7	7***	1.8	61	15.4	41***	10.4	32*	8.1	57***	14.4
Non-smokers	876	10.0	11.4	13.9	54	6.2	107	12.2	222	25.3	44	5.0	51	5.8
Smokers	294	45	15.3	18.4	7*	2.4	44	15.0	45***	15.3	24*	8.2	33**	11.2
Age (years)														
<20	185	2	1.1	17.3	28	15.1	2	1.1	95	51.4	0		7	3.8
20-29	261	20	7.7	13.8	20	7.7	25	9.6	90	34.5	9	3.5	11	4.2
30-39	239	28	11.7	17.2	6	2.5	29	12.1	31	13.0	18	7.5	19	8.0
40-49	210	42	20.0	16.2	0		42	20.0	20	9.5	16	7.6	22	10.5
50-59	139	30	21.6	10.8	2	1.4	32	23.0	17	12.2	13	9.4	12	8.6
60-69	83	13	15.7	10.8	5	6.0	13	15.7	10	12.1	8	9.6	9	10.8
≥70	53	10	18.9	17.0	0		8	15.1	4	7.6	4	7.6	4	7.6
χ <sup>2</sup> for 7 age subgroups	-	52.151	5.372	17.0	22.916†	22.916†	48.554	161.241	22.318	22.318	0.001	12.405	12.405	0.054
P for 7 age subgroups	-	0.000	0.497	0.000	0.000‡	0.000‡	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Han Chinese	1173	307†††	26.2	14.8	18†††	1.5	191†	16.3	61†††	5.2	143†††	12.2	52††	4.4
Men	589	152†††	25.8	17.5	10†††	1.7	95†	16.1	35†††	5.9	71†††	12.1	25†††	4.2
Women	584	155†††	26.5	12.2	8†	1.4	96	16.4	26†††	4.5	72†††	12.3	27	4.6
BMI ≤ 24 kg/m <sup>2</sup>	927	197†††	21.3	11.3	15†††	1.6	120	13.0	58†††	6.3	88†††	9.5	47	5.1
BMI > 24 kg/m <sup>2</sup>	246	110***,†††	44.7	28.1	3	1.2	71***	28.9	3*,†††	1.2	55***,†††	22.4	5†,†	2.0
Normotensives	980	230†††	23.5	13.2	16†††	1.6	144	14.7	58†††	5.9	101†††	10.3	36††	3.7
Hypertensives	193	77***,†††	39.9	23.3	2	1.0	47***	24.4	3*,†††	1.6	42***,†††	21.8	16**	8.3
Non-drinkers	719	166†††	23.1	12.9	16†††	2.2	124††	17.3	54†††	7.5	75†††	10.4	31	4.3
Drinkers	454	141***,†††	31.1	17.8	2*	0.4	67	14.8	7***,†††	1.5	68*,†††	15.0	21†††	4.6
Non-smokers	826	213†††	25.8	13.0	14†††	1.7	137†	16.6	49†††	5.9	100†††	12.1	33	4.0
Smokers	347	94†††	27.1	19.3	4	1.2	54	15.6	12†††	3.5	43	12.4	19†††	5.5
Age (years)														
<20	180	11††	6.1	10.6	12††	6.7	9†	5.0	49†††	27.2	3	1.7	10	5.6
20-29	241	54†††	22.4	10.0	0†††		33	13.7	4†††	1.7	20†	8.3	6	2.5
30-39	251	61†††	24.3	18.3	5	2.0	33	13.2	6†††	2.4	27	10.8	10	4.0
40-49	208	79†††	38.0	19.7	0		51	24.5	1†††	0.5	38††	18.3	11†	5.3
50-59	139	52††	37.4	15.8	0		40	28.8	0†††		30††	21.6	7	5.0
60-69	85	29††	34.1	17.7	1	1.2	16	18.8	0††		11	12.9	5	5.9
≥70	69	21	30.4	10.1	0		9	13.0	1	1.5	14†	20.3	3	4.4
χ <sup>2</sup> for 7 age subgroups	-	67.235	15.325	10.18	6.046§	6.046§	47.008	210.872	45.422	45.422	0.000	3.704	3.704	0.717
P for 7 age subgroups	-	0.000	0.018	0.018	0.014§	0.014§	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TC, total cholesterol; HDL-C, HDL cholesterol; LDL-C, LDL cholesterol.  
 \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001 for comparison with men, BMI ≤ 24 kg/m<sup>2</sup>, normotensive, non-drinkers or non-smokers of the same ethnic group.  
 †P < 0.05, ††P < 0.01, †††P < 0.001 for comparison with the same subgroup of Bai Ku Yao.  
 ‡Values for the comparison of <20, 20-29 and 30-39 years age subgroups in Bai Ku Yao.  
 §Values for the comparison of <20 and 30-39 years age subgroups in Han Chinese.



**Table 3** Comparison of the risk factors for hyperlipidaemia between the Bai Ku Yao and Han populations

Risk factor	Regression coefficient	SE	Wald	P	OR
<b>Han plus Bai</b>					
Ethnic group	0.386	0.096	16.221	0.000	1.471
Age	0.093	0.032	8.223	0.004	1.098
Physical activity	-0.221	0.105	9.265	0.003	1.212
Alcohol consumption	0.141	0.055	6.699	0.010	1.152
Waist circumference	0.835	0.134	41.558	0.000	2.323
BMI	0.915	0.115	62.937	0.000	2.498
Total energy	0.575	0.143	10.366	0.001	1.324
Total fat	0.551	0.135	17.997	0.000	1.676
Total dietary fibre	-0.443	0.131	12.158	0.000	1.574
<b>Bai Ku Yao</b>					
Physical activity	-0.198	0.093	6.753	0.016	1.301
Waist circumference	0.876	0.177	27.563	0.000	2.396
BMI	0.844	0.179	22.284	0.000	2.326
Total energy	0.631	0.184	12.536	0.000	1.975
Total fat	0.573	0.201	9.433	0.003	1.877
Total dietary fibre	-0.625	0.202	10.369	0.001	1.973
<b>Han Chinese</b>					
Age	0.136	0.043	10.131	0.001	1.146
Physical activity	-0.232	0.078	4.562	0.035	1.274
Alcohol consumption	0.145	0.072	4.020	0.045	1.156
Waist circumference	0.763	0.248	14.674	0.000	2.233
BMI	0.988	0.153	41.469	0.000	2.687
Total energy	0.552	0.215	7.823	0.007	1.824
Total fat	0.477	0.189	8.996	0.003	1.679
Total dietary fibre	-0.424	0.245	4.972	0.036	1.616

For the multiple logistic regression analysis, the data were recorded as follows: ethnic group: Bai Ku Yao = 0, Han = 1; sex: female = 0, male = 1; age (years): <20 = 1, 20–29 = 2, 30–39 = 3, 40–49 = 4, 50–59 = 5, 60–69 = 6, ≥70 = 7; BMI (kg/m<sup>2</sup>): ≤24 = 0, >24 = 1; blood pressure: normotensive = 0, hypertensive = 1; alcohol consumption (g/d): non-drinkers = 0, <25 = 1, 25–49 = 2, 50–99 = 3, ≥100 = 4; cigarette smoking (cigarettes/d): non-smokers = 0, <10 = 1, 10–19 = 2, 20–39 = 3, ≥40 = 4.

**Table 4** Comparison of hyperlipidaemia in Bai Ku Yao between Lihu and Baxu villages

Village	n	TC > 5.17 mmol/l		TAG > 1.70 mmol/l		HDL-C < 1.00 mmol/l		LDL-C > 3.20 mmol/l		Apo A1 < 1.0 g/l		Apo B > 1.14 g/l		Apo A1:apo B > 2.50		Hyperlipidaemia prevalence	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%		
Lihu	602	59	9.8	114	18.9	40	6.6	60	10.0	181	30.1	30	5.0	44	7.3	158	26.3
Baxu	568	86	15.1	62	10.9	21	3.7	91	16.0	86	15.1	38	6.7	40	7.0	127	22.4
χ <sup>2</sup>	–	7.676		14.714		5.137		9.530		36.965		1.555		0.031		2.396	
P	–	0.006		0.000		0.023		0.002		0.000		0.212		0.860		0.121	

TC, total cholesterol; HDL-C, HDL cholesterol; LDL-C, LDL cholesterol; hyperlipidaemia, TC > 5.17 mmol/l and/or TAG > 1.70 mmol/l.

Bai Ku Yao than in Han. This discrepancy may mainly be attributed to the differences in dietary habits and lifestyle choices between the two ethnic groups. The staple and subsidiary foods are more favourable for lipid profiles in Bai Ku Yao than in Han. Corn contains abundant dietary fibre and high-quality plant protein<sup>(23)</sup>. Consumption of dietary fibre, specifically the soluble type such as pectins and guar gum, can result in a decrease of serum cholesterol levels in healthy and hyperlipidaemic subjects<sup>(24,25)</sup>. Plant protein can promote the transportation and excretion of free cholesterol. Corn oil is rich in PUFA and MUFA<sup>(26)</sup>. Suitable intakes of PUFA and MUFA can lower serum levels of cholesterol and LDL-C<sup>(26,27)</sup>. Soya protein intake is effective in reducing TC by 9.3%, LDL-C by 12.9% and TAG by 10.5%, and in increasing HDL-C by 2.4%. Furthermore, the changes in serum TC and LDL-C concentrations are directly related to the initial serum TC

concentration<sup>(28)</sup>. The hypocholesterolaemic activity of buckwheat protein products is far stronger than that of soya protein isolate and corn<sup>(29)</sup>. Daily ingestion of 4 g caiapo (the extract of white-skinned sweet potato) for 6 weeks reduces TC and LDL-C in type 2 diabetic patients previously treated by diet alone<sup>(30)</sup>. The long-term beneficial effects of caiapo on plasma cholesterol levels are further confirmed in a recent report<sup>(31)</sup>. Studies have also demonstrated that pumpkin or pumpkinseed oil may be a useful therapy for hypercholesterolaemia by reducing oxidative stress and cholesterol levels<sup>(32)</sup>. The people of Bai Ku Yao are accustomed to drinking hempseed soup. There are more than twenty-nine fat-soluble constituents in hempseed, with fatty acid methyl esters making up 99.3%<sup>(33,34)</sup>. The main fatty acid components of hempseed oil are palmitic acid (8.4%), γ-linolenic acid (1.3%), linoleic acid (58.7%), linolenic acid (14.0%), oleic acid

(10.1%), stearic acid (3.8%) and arachidic acid (1.0%)<sup>(35)</sup>. A number of experimental and clinical studies have demonstrated that the beneficial effects of hempseed or hempseed oil on lipid profiles include decreasing the levels of TC, TAG and LDL-C, inhibiting lipid peroxidation, reducing the atherogenic index and increasing the levels of HDL-C<sup>(36–38)</sup>. The intake of animal fat, body weight and BMI were all lower in Bai Ku Yao than in Han. For nearly 50 years it has been widely accepted that high-fat diets, particularly those that contain large quantities of SFA, raise blood cholesterol concentrations and predispose individuals to CVD<sup>(39)</sup>. In addition, strict intra-ethnic marriages have been performed from time immemorial in the Bai Ku Yao. Therefore, the hereditary characteristics, genotypes and phenotypes of lipids in the Bai Ku Yao population may be different from those in Han Chinese. However, this still needs to be determined.

It is well documented that androgens induce disadvantageous effects on lipid profiles, whereas oestrogens are held to have opposite effects<sup>(40,41)</sup>. In the present study, we also showed that the prevalence of high TAG in both ethnic groups and low HDL-C and apo A1 in Bai Ku Yao was higher in males than in females. These discrepancies in dyslipidaemia between men and women might be partially attributable to the effects of gonadal hormones.

Epidemiological studies have provided abundant evidence that lipid levels are closely related with age<sup>(42,43)</sup>. In the present study, we showed that there was a significant difference in the prevalence of high TC, LDL-C and apo B, and low HDL-C and apo A1, for the age subgroups in both ethnic groups. This is consistent with previous studies. The exact mechanisms of age on hyperlipidaemia are not well known. They may due, in part, to the hereditary characteristics and ageing of the population.

The link between obesity and dyslipidaemia has been clearly documented<sup>(44,45)</sup>. Obesity not only induces dyslipidaemia, but also is directly associated with diabetes, hypertension and coronary artery disease. The current study also shows that the prevalence of high TC, TAG, LDL-C and apo B in both ethnic groups was higher in subjects with BMI > 24 kg/m<sup>2</sup> than those with BMI ≤ 24 kg/m<sup>2</sup>. This is in agreement with the findings of previous studies<sup>(44–46)</sup>. Dyslipidaemia in obesity may result from insulin resistance<sup>(47,48)</sup>. Insulin resistance can increase plasma NEFA concentrations and stimulate VLDL synthesis and release. At the same time, insulin resistance can also suppress lipoprotein lipase activity and increase plasma VLDL levels.

The association of hyperlipidaemia with hypertension is poorly understood. Studies have shown that the levels of TAG, VLDL cholesterol and apo E are higher and levels of apo AI, apo AII and apo CII are lower in untreated hypertensives than in controls<sup>(49)</sup>. TC and non-HDL-C levels increase significantly with increasing systolic or diastolic blood pressure in both sexes<sup>(50)</sup>. Some authors

believe this relationship may be a kind of random phenomenon<sup>(51)</sup>. In the present study, we found that the prevalence of hypercholesterolaemia in both ethnic groups and hypertriglycerolaemia in Han was higher in hypertensives than in normotensives. These findings suggest that there may be a complicated interrelation between hyperlipidaemia and hypertension<sup>(52)</sup>.

Both case-control and cohort studies have described a J- or U-shaped association between alcohol intake and CHD and between alcohol intake and mortality<sup>(53)</sup>. Although moderate alcohol consumption appears to be protective, heavy consumption of alcohol is associated with subclinical impairment of left ventricular function and occasionally results in overt cardiomyopathy. High alcohol intake and cigarette smoking have disadvantageous effects on lipid profiles. Smoking also disturbs lipoprotein metabolism by raising insulin resistance and lipid intolerance, and is implicated in the production of small dense LDL-C. In the present study, we showed that the prevalence of high TC, TAG and apo B in both ethnic groups was higher in drinkers than in non-drinkers. The prevalence of high apo B and apo A1:apo B ratio in Bai Ku Yao and high TAG in Han was higher in smokers than in non-smokers. These findings indicate that both alcohol consumption and cigarette smoking may be risk factors for hyperlipidaemia in the two populations.

In the present study, we also showed that the prevalence of high TAG, low HDL-C and apo A1 in Bai Ku Yao was higher, but the prevalence of high TC was lower, in subjects from Lihu village than from Baxu village. The reason for this discrepancy is not yet known. The customs and cultures in both villages are similar. One difference is the residential locus. The distance between the two villages is about 50 km. It is unclear whether geographical or other unknown factors might be involved in this discrepancy between the two villages.

## Conclusion

The present study reveals that the prevalence of hypercholesterolaemia and hyperlipidaemia was lower in Bai Ku Yao than in Han. These differences between the two ethnic groups may result from different demographic characteristics, dietary patterns, lifestyle choices and physical activity level, as well as genetic background.

## Acknowledgements

*Conflict of interest:* No conflicts.

*Funding support:* The study was supported by the National Natural Science Foundation of China (No. 30660061).

*Authorship responsibilities:* Y.R. conceived the study hypothesis, participated in the design, carried out the epidemiological survey, collected the samples, supervised data

analyses and drafted the manuscript; Y.D., L.S., P.S. and L.W. carried out the epidemiological survey and biochemical analyses; C.Y. carried out the epidemiological survey and helped to carry out biochemical analyses; F.Q. performed the statistical analyses; Y.H., T.J. and L.Y. carried out the epidemiological survey. All authors read and approved the final manuscript.

## References

1. Grundy SM, Cleeman JI, Merz CN, Brewer HB Jr, Clark LT, Hunninghake DB, Pasternak RC, Smith SC Jr & Stone NJ; National Heart, Lung, and Blood Institute; American College of Cardiology Foundation; American Heart Association (2004) Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation* **110**, 227–239.
2. Third Joint Task Force of European and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (2003) European guidelines on cardiovascular disease prevention in clinical practice. *Eur J Cardiovasc Prev Rehabil* **10**, Suppl. 1, S1–S78.
3. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (2001) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* **285**, 2486–2497.
4. Satoh H, Nishino T, Tomita K & Tsutsui H (2006) Fasting triglyceride is a significant risk factor for coronary artery disease in middle-aged Japanese men: results from a 10-year cohort study. *Circ J* **70**, 227–231.
5. Achari V & Thakur AK (2004) Association of major modifiable risk factors among patients with coronary artery disease – a retrospective analysis. *J Assoc Physicians India* **52**, 103–108.
6. Kwiterovich PO Jr, Coresh J, Smith HH, Bachorik PS, Derby CA & Pearson TA (1992) Comparison of the plasma levels of apolipoproteins B and A-1, and other risk factors in men and women with premature coronary artery disease. *Am J Cardiol* **69**, 1015–1021.
7. Boden WE (2000) High-density lipoprotein cholesterol as an independent risk factor in cardiovascular disease: assessing the data from Framingham to the Veterans Affairs High-Density Lipoprotein Intervention Trial. *Am J Cardiol* **86**, 19L–22L.
8. Sharrett AR, Ballantyne CM, Coady SA, Heiss G, Sorlie PD, Catellier D & Patsch W, Atherosclerosis Risk in Communities Study Group (2001) Coronary heart disease prediction from lipoprotein cholesterol levels, triglycerides, lipoprotein(a), apolipoproteins A-I and B, and HDL density subfractions: The Atherosclerosis Risk in Communities (ARIC) Study. *Circulation* **104**, 1108–1113.
9. Sociedade Brasileira de Cardiologia (2001) III Diretrizes Brasileiras Sobre Dislipidemias e Diretrizes de Prevenção da Aterosclerose. *Arq Bras Cardiol* **77**, Suppl. III, 1–48.
10. Bermudez OI, Velez-Carrasco W, Schaefer EJ & Tucker KL (2002) Dietary and plasma lipid, lipoprotein, and apolipoprotein profiles among elderly Hispanics and non-Hispanics and their association with diabetes. *Am J Clin Nutr* **76**, 1214–1221.
11. Srinivasan SR, Freedman DS, Webber LS & Berenson GS (1987) Black–white differences in cholesterol levels of serum high-density lipoprotein subclasses among children: the Bogalusa Heart Study. *Circulation* **76**, 272–279.
12. Nethononda MR, Essop MR, Mbewu AD & Galpin JS (2004) Coronary artery disease and risk factors in Black South Africans – a comparative study. *Ethn Dis* **14**, 515–519.
13. People's Republic of China–United States Cardiovascular and Cardiopulmonary Epidemiology Research Group (1992) An epidemiological study of cardiovascular and cardiopulmonary disease risk factors in four populations in the People's Republic of China. Baseline report from the PRC–USA Collaborative Study. *Circulation* **85**, 1083–1096.
14. Lyu LC, Yeh CY, Lichtenstein AH, Li Z, Ordovas JM & Schaefer EJ (2001) Association of sex, adiposity, and diet with HDL subclasses in middle-aged Chinese. *Am J Clin Nutr* **74**, 64–71.
15. Yang YX, Wang GY & Pan XC (2002) *The 2002 Chinese Food Composition Table*. Beijing: Medical Publishing House of Beijing University.
16. Paffenbarger RS, Wing AL & Hyde RT (1978) Physical activity as an index of heart attack risk in college alumni. *Am J Epidemiol* **108**, 161–175.
17. Yin R, Chen Y, Pan S *et al.* (2006) Comparison of lipid levels, hyperlipidemia prevalence and its risk factors between Guangxi Hei Yi Zhuang and Han populations. *Arch Med Res* **37**, 787–793.
18. Yin R, Wang Y, Chen G, Lin W, Yang D & Pan S (2006) Lipoprotein lipase gene polymorphism at Pvu II locus and serum lipid levels in the Guangxi Hei Yi Zhuang and Han populations. *Clin Chem Lab Med* **44**, 1416–1421.
19. Ruixing Y, Limei Y, Yuming C, Dezhai Y, Weixiong L, Muyan L, Fengping H, Jinzhen W, Guangqing Y & Zhenbiao N (2006) Prevalence, awareness, treatment, control and risk factors of hypertension in the Guangxi Hei Yi Zhuang and Han populations. *Hypertens Res* **29**, 423–432.
20. Ruixing Y, Jiaqiang D, Dezhai Y, Weixiong L, Shangling P, Jinzhen W, Jiandong H & Xiuyan L (2006) Effects of demographic characteristics, health-related behaviors and lifestyle factors on the prevalence of hypertension for the middle-aged and elderly in the Guangxi Hei Yi Zhuang and Han populations. *Kidney Blood Press Res* **29**, 312–320.
21. Cooperative Meta-analysis Group of China Obesity Task Force (2002) Predictive values of body mass index and waist circumference to risk factors of related diseases in Chinese adult population. *Chin J Epidemiol* **23**, 5–10.
22. von Kamp A & Schuster S (2006) Metatool 5.0: fast and flexible elementary modes analysis. *Bioinformatics* **22**, 1930–1931.
23. Dong W, Ma X, Zhang D & Yu S (2002) Effect of maize embryo on delaying aging. *Food Sci* **23**, 95–97.
24. Jenkins DJ, Kendall CW, Axelsen M, Augustin LS & Vusksan V (2000) Viscous and nonviscous fibres, nonabsorbable and low glycaemic index carbohydrates, blood lipids and coronary heart disease. *Curr Opin Lipidol* **11**, 49–56.
25. Lairon D (1996) Dietary fibres: effects on lipid metabolism and mechanisms of action. *Eur J Clin Nutr* **50**, 125–133.
26. Liu Y, Zhang L & Wu Y (1995) The dietary therapy for hyperlipidemia complicated with NIDDM. *Chin J Clin Nutr* **3**, 174–176.
27. Zhang Y, Zhou Y, Wu F & Zhang M (1996) The comparative effects of maize oil and lard on blood lipids serum glucose and brain lipofuscin in rats. *Acta Nutrimenta Sinica* **18**, 274–279.
28. Anderson JW, Johnstone BM & Cook-Newell ME (1995) Meta-analysis of the effects of soy protein intake on serum lipids. *N Engl J Med* **333**, 276–282.
29. Zhang HW, Zhang YH, Lu MJ, Tong WJ & Cao GW (2007) Comparison of hypertension, dyslipidaemia and hyperglycaemia between buckwheat seed-consuming and non-consuming Mongolian-Chinese populations in Inner Mongolia, China. *Clin Exp Pharmacol Physiol* **34**, 838–844.



30. Ludvik BH, Mahdjoobian K, Waldhaeusl W, Hofer A, Prager R, Kautzky-Willer A & Pacini G (2002) The effect of *Ipomoea batatas* (cayapo) on glucose metabolism and serum cholesterol in patients with type 2 diabetes: a randomized study. *Diabetes Care* **25**, 239–240.
31. Ludvik B, Neuffer B & Pacini G (2004) Efficacy of *Ipomoea batatas* (cayapo) on diabetes control in type 2 diabetic subjects treated with diet. *Diabetes Care* **27**, 436–440.
32. al-Zuhair H, Abd el-Fattah AA & Abd el Latif HA (1997) Efficacy of simvastatin and pumpkin-seed oil in the management of dietary-induced hypercholesterolemia. *Pharmacol Res* **35**, 403–408.
33. Zhang Y & Wang ZZ (2006) GC–MS analysis of fat-soluble constituents of hemp kernels. *Acta Bot Boreal Occident Sin* **26**, 1955–1958.
34. Ross SA, Mehmedic Z, Murphy TP & Elsohly MA (2000) GC–MS analysis of the total  $\Delta^9$ -THC content of both drug- and fiber-type cannabis seeds. *J Anal Toxicol* **24**, 715–717.
35. Zhou YH (2004) Analysis of fatty acid in hemp fruit oil with GC–MS. *Chin Oils Fats* **29**, 72–73.
36. Ren HY, Sun HG, Ma JZ, Zhang Y, Yi CR, Wu MX, Liu WL & Li GL (1997) Experimental study on the effects of hemp fruit oil on serum lipid levels and lipid peroxidation. *Chin J Tradit Med Sci Technol* **4**, 200.
37. Ren HY, Sun HG, Zhang Y, Yi CR, Wu MX, Li GL & Liu WL (1998) Lipid-lowering and antiatherosclerotic effects of hemp fruit oil in partridges. *Henan Tradit Chin Med* **18**, 294–295.
38. Schwab US, Callaway J, Erkkila AT, Gynther J, Uusitupa MI & Jarvinen T (2006) Effects of hempseed and flaxseed oils on the profile of serum lipids, serum total and lipoprotein lipid concentrations and haemostatic factors. *Eur J Nutr* **45**, 470–477.
39. Yu-Poth S, Zhao G, Etherton T, Naglak M, Jonnalagadda S & Kris-Etherton P (1999) Effects of National Cholesterol Education Program's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. *Am J Clin Nutr* **69**, 632–646.
40. Mudali S, Dobs AS, Ding J, Cauley JA, Szklo M & Golden SH, Atherosclerosis Risk in Communities Study (2005) Endogenous postmenopausal hormones and serum lipids: the Atherosclerosis Risk in Communities Study. *J Clin Endocrinol Metab* **90**, 1202–1209.
41. Cheung AP (2000) Acute effects of estradiol and progesterone on insulin, lipids and lipoproteins in postmenopausal women: a pilot study. *Maturitas* **35**, 45–50.
42. Karki DB, Neopane A, Pradhan B & Magar A (2004) Lipid levels in Nepalese population. *Kathmandu Univ Med J (KUMJ)* **2**, 349–353.
43. Guize L, Benetos A, Thomas F, Malmejac A & Ducimetiere P (1998) Cholesterolemia and total, cardiovascular and cancer mortality. Study of a cohort of 220,000 people. *Bull Acad Natl Med* **182**, 631–650.
44. Li D, Yu XM, Zhou XQ, Zhang YH, Yao T & Sinclair AJ (2004) Relationship between BMI and serum and lipoprotein lipids in the Hangzhou region. *Asia Pac J Clin Nutr* **13**, S68.
45. Mataix J, Lopez-Frias M, Martinez-de-Victoria E, Lopez-Jurado M, Aranda P & Llopis J (2005) Factors associated with obesity in an adult Mediterranean population: influence on plasma lipid profile. *J Am Coll Nutr* **24**, 456–465.
46. Kawada T (2002) Body mass index is a good predictor of hypertension and hyperlipidemia in a rural Japanese population. *Int J Obes Relat Metab Disord* **26**, 725–729.
47. Houston MC, Basile J, Bestermann WH *et al.* (2005) Addressing the global cardiovascular risk of hypertension, dyslipidemia, and insulin resistance in the southeastern United States. *Am J Med Sci* **329**, 276–291.
48. Pei D, Kuo SW, Wu DA, Lin TY, Hseih MC, Lee CH, Hsu WL, Chen SP, Sheu WH & Li JC (2005) The relationships between insulin resistance and components of metabolic syndrome in Taiwanese Asians. *Int J Clin Pract* **59**, 1408–1416.
49. Catalano M, Aronica A, Carzaniga G, Seregini R & Libretti A (1991) Serum lipids and apolipoproteins in patients with essential hypertension. *Atherosclerosis* **87**, 17–22.
50. Bona KH & Thelle DS (1991) Association between blood pressure and serum lipids in a population. The Tromso Study. *Circulation* **83**, 1305–1314.
51. Li J, Niu Q, Li P, Wang S & Zhao S (1991) The levels of serum lipids, lipoprotein, and apolipoprotein in patients with essential hypertension. *Chin J Cardiol* **19**, 317–319.
52. Borghi C, Dormi A, Veronesi M, Sangiorgi Z & Gaddi A; Brisighella Heart Study Working Party (2004) Association between different lipid-lowering treatment strategies and blood pressure control in the Brisighella Heart Study. *Am Heart J* **148**, 285–292.
53. Corella D, Tucker K, Lahoz C, Coltell O, Cupples LA, Wilson PW, Schaefer EJ & Ordovas JM (2001) Alcohol drinking determines the effect of the APOE locus on LDL-cholesterol concentrations in men: the Framingham Offspring Study. *Am J Clin Nutr* **73**, 736–745.