

research has also shed light on potential neurobiological mechanisms and genetic factors contributing to shared psychosis. Therapeutically, early intervention and tailored treatment plans are crucial in achieving favorable outcomes. While antipsychotic medications remain a cornerstone of treatment, family therapy and psychoeducation have emerged as valuable adjunctive approaches to address the unique challenges posed by shared psychosis.

Conclusions: In conclusion, “Folie à deux” continues to be a captivating and clinically relevant phenomenon in contemporary psychiatry. This bibliographical review underscores the importance of recognizing and diagnosing shared psychosis in clinical practice. Moreover, it highlights the need for further research to unravel the underlying mechanisms and genetic predispositions associated with this condition. Ultimately, a multidisciplinary approach, including pharmacological, psychotherapeutic, and family-based interventions, holds promise in improving the prognosis of individuals affected by “Folie à deux.”

Disclosure of Interest: None Declared

EPV0968

Ferroptosis affects cognitive dysfunction and the progression of mental illnesses

B. Nycz and K. Krysta**

Department of Rehabilitation Psychiatry, Medical University of Silesia, Katowice, Poland

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1556

Introduction: Ferroptosis is a programmed form of cell death characterized by excessive accumulation of intracellular iron fraction, uncontrolled lipid peroxidation, impairment of glutathione-dependent antioxidant functions, and loss of oxidative-antioxidant balance. Nerve cells are sensitive to excessive amounts of iron, which impairs the functioning of mitochondria and leads to their death. Ferroptosis has been identified in neurological diseases such as stroke, Alzheimer’s disease, Parkinson’s disease. Features of ferroptotic cells have been observed in models expressing cognitive deficits, and ferroptosis-related genes have been associated with the development of mental illnesses.

Objectives: The aim of the study was to analyze the available literature on the relationship between the occurrence of ferroptosis and cognitive impairment occurring in mental diseases, such as schizophrenia

Methods: The publications found in the PubMed database were analyzed after entering the following entries: ferroptosis, mental illness, cognitive functions, schizophrenia.

Results: Ferroptosis occurs in mental illnesses. Increased expression of the TP53 and VEGFA genes, which are associated with ferroptosis, has been identified in patients suffering from schizophrenia. Animal research confirms that disturbed iron homeostasis causes iron overload in nerve cells, which leads to ferroptosis and has a neurodegenerative effect, as well as deepens cognitive deficits. The use of iron chelator has a neuroprotective effect and reduces the occurrence of cognitive disorders.

Conclusions: Genes associated with ferroptosis may influence the development of schizophrenia, which means that ferroptosis may be involved in the pathophysiology of schizophrenia. Excess iron

inside nerve cells, as a feature of ferroptosis, may affect the deterioration of cognitive functioning. Administration of iron chelators protects neurons by reducing the toxic effects of iron.

Disclosure of Interest: None Declared

EPV0969

Determinants of hospital length of stay for patients with schizophrenia

M. Turki, M. Barkallah*, N. Boussaid, A. labyadh, S. Ellouze, N. Halouani and J. Aloulou

Psychiatrie B, CHU Hedi Chaker, Sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1557

Introduction: relapse and frequent rehospitalizations. The length of stay (LOS) of these patients has been a concern of researchers. The ability to identify determinants of LOS at admission – and, thus, identify patients who are likely to need a longer stay early on – may help treatment planning.

Objectives: We aimed to investigate socio-demographic and clinical profile of inpatients with schizophrenia, and to identify factors associated with LOS.

Methods: It was a retrospective study carried out among 90 inpatients with schizophrenia admitted to the psychiatry “B” department, Hedi Chaker university hospital (Sfax, Tunisia), during the period between January 2015 and December 2019. Data collection was performed through the patients’ medical records. Statistical analysis was performed using SPSS v.25.

Results: The mean age of our patients was 32 years. Among them, 57.78% were women. The mean LOS was 28 days. Factors found to be significantly associated with LOS were: the number of admissions ($p < 0.001$, $r = 0.404$), involuntary hospitalization ($p = 0.001$), violence and disturbance of public order as a reason of admission ($p < 0.001$) and the lack of social support ($p = 0.039$). As for the clinical symptoms, hallucinations were significantly associated with a longer LOS ($p = 0.001$).

Conclusions: Our findings highlighted several factors associated with a longer LOS. This may be helpful to the management of hospitalization and ensuring that any periods of liberty deprivation do not last longer than necessary to provide appropriate treatment.

Disclosure of Interest: None Declared

EPV0970

Hydroxychloroquine in systemic lupus erythematosus and psychosis. A case report

M. Martínez Grimal*, A. M. Morales Rivero, P. Rivero Rodríguez, E. E. Morales Castellano and N. Molina Pérez

Hospital Universitario de Gran Canaria Doctor Negrín, Las Palmas, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1558

Introduction: Hydroxychloroquine, an antimalarial drug, is an important therapeutic tool in the management of rheumatic diseases such as Systemic Lupus Erythematosus (SLE) due to its anti-inflammatory action. SLE is a chronic autoimmune inflammatory disease that affects the connective tissue of multiple organs. Neuropsychiatric disturbances in SLE are common; however, lupus psychosis is rare, occurring in 2 to 11% of patients. The literature has described the emergence of neuropsychiatric symptoms as an adverse effect of hydroxychloroquine use, with some patients experiencing clinical depression, anxiety, suicidal ideation, and psychotic symptoms.

Objectives: The aim of this work is to review the available evidence regarding neuropsychiatric symptoms secondary to the use of hydroxychloroquine.

Methods: The case of a 50-year-old woman diagnosed with SLE, with no other relevant medical history, has been evaluated. She was brought to the emergency department due to paranoid and persecutory ideas, as well as self-referentiality, coinciding with the introduction of hydroxychloroquine in her treatment. She was admitted to the University Hospital of Gran Canaria Doctor Negrín with a diagnostic orientation of a first psychotic episode.

Results: The presence of neuropsychiatric symptoms in patients diagnosed with SLE is so common that they constitute a diagnostic criterion for the disease. On the other hand, the medications used for therapeutic management of this disease can lead to the emergence of new neuropsychiatric symptoms or exacerbate preexisting neuropsychiatric clinical manifestations.

Conclusions: The study of this case highlights the challenges in establishing a differential diagnosis between primary SLE symptoms that require an increase in hydroxychloroquine and those caused by its own treatment. It underscores the need for further studies to explore the risk of psychiatric symptoms associated with the use of hydroxychloroquine, as well as its impact on the course of underlying mental disorders.

Disclosure of Interest: None Declared

EPV0971

Clinical experience with once-monthly risperidone ISM in a mental health center. A retrospective study.

A. Merino Iglesias*, N. Rodríguez Ortega, L. Montero Quer and M. R. Rodríguez Campos

Psychiatry, Hospital Universitario de Fuenlabrada, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1559

Introduction: Long-acting injectable antipsychotics have undergone a great development in recent years, becoming useful tools to facilitate therapeutic adherence. Once-monthly risperidone ISM is a new way of treatment which has been commercialized in Spain since September 2022.

Objectives: In this study we will analyze our clinical experience with this treatment, especially in terms of tolerance and efficacy, during its first year of widespread use.

Methods: Longitudinal retrospective study of monthly risperidone users in a mental health center in the Autonomous Community of Madrid (Spain), between September 2022 and September 2023.

A sample of 13 patients was selected, collecting both sociodemographic (age, gender) and clinical variables (diagnosis, dose, time elapsed, number of hospital readmissions, adverse effects and monotherapy or combined use). A descriptive analysis of the collected data was then carried out.

Results: Monthly risperidone was used in 13 patients: 15% (n=2) were women, and 85% (n=11) were male. The mean age of the patients was 43.6 years. The most frequent diagnosis of these patients was “psychotic disorders” (84,6%, n= 11), with other diagnoses such as schizoaffective disorder (7,7%, n=1) and obsessive compulsive disorder (7,7%, n=1).

The doses used of risperidone were 100mg every month in 61,5% of patients (n=8) and 75mg in 38,5% of patients (n=5). The mean time since the first administration was 4.35 months.

Concerning monotherapy, 84,6% (n=11) of patients on monthly risperidone were on antipsychotic monotherapy, while 15,4% (n=2) required more than one antipsychotic. Among the switches made to monthly risperidone, 69,2% (n=9) were previously treated with oral risperidone, 15,4% (n=2) were treated with once-biweekly risperidone long-acting injectable, 7,9% (n=1) with oral paliperidone and 7,9% with aripiprazole monthly injectable.

During the study period, hospital readmissions for psychiatric decompensations occurred in one patient (7,9%, n=1), while the rest of the patients (92,1%, n=12) did not present decompensations that required psychiatric admission.

Moderate or severe effects occurred in one patient (7,9%, n=1), in the form of acute dystonia, which led to the interruption of injectable treatment. The rest of the patients (92,1%, n=12) did not present severe adverse effects. Minor adverse effects appeared in 3 patients (25%); these adverse effects were already present in the previous treatment with oral risperidone and did not condition the suspension of the treatment.

Conclusions: In the sample analyzed, once-monthly Risperidone ISM had reasonable tolerance levels. Also, it's shown to be effective in preventing psychotic decompensations and hospital admissions. Therefore, this new injectable of monthly risperidone represents a therapeutic alternative to consider in order to guarantee therapeutic adherence and improve the quality of life of patients with psychotic symptoms.

Disclosure of Interest: None Declared

EPV0972

Psychometric assessment of patients with treatment-resistant schizophrenia

V. Kaleda* and D. Tikhonov

Mental Health Research Centre, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1560

Introduction: Treatment-resistant schizophrenia (TRS) is one of the most pressing issues in the field of treatment and research of psychotic disorders. The pronounced decline in social and professional functioning in this group of patients as well as high costs of therapy determine high interest in TRS. This is a part of an ongoing study on the clinical and biological features of TRS.