

<sup>1</sup> Faculty of Psychology, Department of Clinical Psychology, Bergen, Norway

<sup>2</sup> Faculty of Medicine, Institute of Clinical Medicine, Oslo, Norway

<sup>3</sup> Solli District Psychiatric Centre DPS, Department of Comprehensive Treatment Programs, Nesttun, Norway

<sup>4</sup> Faculty of Psychology, Department of Psychosocial Science, Bergen, Norway

\* Corresponding author.

**Introduction** Seventy percent of patients with panic disorder (PD) have sleep impairment. Cognitive behavior therapy (CBT) effectively treats PD, but the concomitant effect on sleep is understudied. Physical exercise (PE) improves sleep, but this has not been investigated in patients with PD.

**Objective** To compare the effects of CBT or PE on sleep in PD-patients, and to determine potential mechanisms of action.

**Methods** Thirty-six PD-patients were randomized to either group CBT for PD or regular PE. Sleep was assessed pre/post with the Pittsburgh sleep quality index. Effects were investigated with repeated measures ANOVA and t-tests. Expected mediators were added to the general linear model to assess mediation.

**Results** The effect of time was significant,  $F(1.33)=10.11$ ,  $P=0.003$ , but not the interaction (Time  $\times$  group),  $F(1.33)=.48$ ,  $P=0.49$ . Symptoms were significantly reduced from pre- to post-treatment: PE,  $t(16)=3.03$ ,  $P=.008$ , and CBT,  $t(17)=2.18$ ,  $P=0.044$ . CBT-patients changed significantly ( $P<0.05$ ) on Sleep quality,  $t(17)=2.47$  and Sleep disturbance,  $t(17)=2.38$ . PE-patients changed significantly on sleep duration,  $t(16)=2.58$  and sleep disturbance,  $t(16)=2.58$ . A significant interaction with change in fear of bodily symptoms,  $F(1.16)=5.53$ ,  $P=.032$ , and with change in depression-level,  $F(1.16)=12.13$ ,  $P=0.003$  was only found for CBT. A significant interaction with change in physical fitness,  $F(1.15)=5.01$ ,  $P=.041$ , was only found for PE.

**Conclusion** Both interventions improve sleep in PD-patients, but differently. The findings also suggest that these changes are related to different mechanisms for PE and CBT.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1875>

## EW0007

### Explanatory and confirmatory factor structure of beck anxiety inventory in college sample

S. Kareemi\*, B.M. Alansari

Kuwait University, Faculty of Social Sciences, psychology, kifan, Kuwait

\* Corresponding author.

**Introduction** The Beck anxiety inventory (BAI) is a widely used 21-item self-report inventory used to assess anxiety levels in adults and adolescents in both clinical and non-clinical populations. The values for each item are summed yielding an overall or total score for all 21 symptoms that can range between 0 and 63 points. A total score of 0–7 is interpreted as a “Minimal” level of anxiety; 8–15 as “Mild”; 16–25 as “Moderate”, and; 26–63 as “Severe”. There is no study until this date that examines the Explanatory and confirmatory factor structure of BAI in college student in Kuwait.

**Objectives** The current study investigated the original four-factor structure of the (BAI) in non-clinical sample of college students.

**Methods** Sample one consisted of 540 males and females while sample two consisted of 600 males and females from Kuwait University undergraduates. The Arabic version of BAI was administered to participants. Explanatory factor analysis based on sample one and conformity factor analysis based on sample 2.

**Results** The results revealed four factor structures of BAI in the two samples of Kuwaiti students. Which included neuro-physiological, subjective, autonomic, and panic factors.

**Conclusions** The results of both confirmatory and exploratory factor analysis indicated that the original four-factor structures of the BAI do provide the best fit for the college sample.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1876>

## EW0008

### Playing video games – Psychological threat to adults?

M. Toś<sup>1</sup>, E. Wilk<sup>1</sup>, A. Myszczyk<sup>1</sup>, A. Bratek<sup>2</sup>, K. Krzysztof<sup>1,\*</sup>

<sup>1</sup> Medical University of Silesia, Department of Rehabilitation Psychiatry, Katowice, Poland

<sup>2</sup> Medical University of Silesia, Department of Psychiatry and Psychotherapy, Katowice, Poland

\* Corresponding author.

**Background** Video games become increasingly popular form of spending free time, therefore they are often a research subject. Researchers focus mainly on video games influence over children's psyche and their social interactions, although video games can also have an impact on adult's behaviour.

**Objectives** Incidence of social anxiety disorder and impulsiveness among video game players and non-players.

**Material and methods** An anonymous online survey included 263 students of medical university of Silesia (112 M and 151 W). In study group, 142 people (54%) declared to be players. Questionnaire contained questions about playing time, Barratt Impulsiveness Scale and Liebowitz Social Anxiety Scale. Statistical analyses were performed using the statistical software package Statistica 12.

**Results** Using Liebowitz Scale in 168 (64.86%) all studied people lack of phobia was found, moderate social phobia 39 (15.06%), marked social phobia 26 (10.03%), severe social phobia 17(6.56%) and very severe social phobia 9(3.47%). There was no statistical significance between players and non-players (test  $\chi^2$   $P=0.6521$ ). Also in Barratt Scale statistical significance was not found in attentional impulsiveness (test U M-W  $P=0.3267$ ) and in Motor impulsiveness (test U M-W  $P=0.3140$ ). Statistical significance was observed in Non-planning impulsiveness (players: 23.68 V non-players: 22.02; test U M-W  $P=0.0036$ ).

**Conclusions** The study did not show clear influence adult's video games playing over social phobia and impulsiveness occurrence.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.1877>

## EW0009

### Utilization unspecialized care of patients with anxiety disorder

M. Marachev\*, A. Avedisova

V. Serbsky Federal Medical Research Centre for Psychiatry and Narcology- of the, Therapy of Mental & Behaviour Disorders, Moscow, Russia

\* Corresponding author.

**Introduction** The greatest social and economic burden is divided between the three main anxiety disorders: social phobia (SF), generalized anxiety disorder (GAD) and panic disorder (PD).

**Objectives** To examine the pathways of patients with PD, SF and GAD since the beginning of the first anxiety symptoms and before the first course of a standardized treatment in a specialized mental health facility.

**Aims** To evaluate the period of delay in seeking specialized care and to identify the main ways of seeking medical/non-medical care.