

There is evidence supporting the physiologic benefits of these agents, but outcome data are lacking, so individualization is necessary, and critical time should not be lost administering pretreatment drugs if the patient requires immediate intubation. Despite the lack of outcome studies, there is considerable inferential evidence supporting this approach, and these agents probably provide protection for vulnerable patients against the adverse hemodynamic and intracranial effects of laryngoscopy and intubation.<sup>2</sup>

Research done at our centre has provided evidence supporting the physiologic benefit of pretreatment agents.<sup>3</sup> In addition, we recently published a study of 522 intubations using etomidate, many of which also involved the use of pretreatment agents. This study demonstrated that our approach was associated with hemodynamic stability in a heterogeneous group of patients undergoing RSI in the ED.<sup>4</sup> Our conclusion from the existing literature remains unchanged; premedication should be considered in selected patients undergoing neuroprotective RSI in the ED. The appropriate selection and dosing of medications in such cases provides the best opportunity to minimize post-intubation hypotension and other complications of intubation.

**Peter J. Zed, BSc, BSc(Pharm), PharmD, FCSHP**

**David W. Harrison, MD**

**Nick Kuzak, MD**

CSU Pharmaceutical Sciences and Department of Emergency Medicine  
Vancouver General Hospital  
Faculty of Pharmaceutical Sciences  
and Faculty of Medicine  
University of British Columbia  
Vancouver, BC

**References**

1. Kuzak N, Harrison DW, Zed PJ. Use of lidocaine and fentanyl premedication for neuroprotective rapid sequence intubation in the emergency department. *Can J Emerg Med* 2006;8(2):80-4.
2. Walls RM. Airway. In: Marx JA,

Hockberger RS, Walls RM, et al, editors. *Emergency medicine: concepts and clinical practice*. 6th ed. Missouri: Mosby; 2006. p. 2-26.

3. Zed PJ, Abu-Laban RB, Harrison DW. Effect of fentanyl pretreatment on sympathetic response in patients with cerebrovascular accident undergoing rapid sequence intubation in the emergency department [abstract]. *Can J Emerg Med* 2004;6(3):197.
4. Zed PJ, Abu-Laban RB, Harrison DW. Intubating conditions and hemodynamic effects of etomidate for rapid sequence intubation in the emergency department: an observational cohort study. *Acad Emerg Med* 2006;13:378-83.

**Treatment of patients with severe sepsis and septic shock: real-life lessons**

*To the Editors:* Evidence-based therapies for severe sepsis and septic shock include broad spectrum antibiotics, early goal-directed resuscitation, corticosteroids, glycemic control and recombinant human activated protein C (rhAPC).<sup>1</sup> Prior to dissemination of the Surviving Sepsis Guidelines in 2004,<sup>1</sup> we found that 94% (32/34) of our septic patients received greater than 20 mL/kg intravenous fluid within 6 hours, that 85% (29/34) received low-dose corticosteroids, that 68% (23/34) received antibiotics within 3 hours, and that 82% (29/33) received rhAPC within 24 hours of admission to the intensive care unit. At the same time, only 38% (13/34) received central venous pressure monitoring, and only 6% (2/34) had central venous oximetry performed within 6 hours. This “care-gap” offers a provocative area for research and improvement.

Pharmaceutical companies have provided a great deal of education focused on products such as rhAPC. Unfortunately, educational funding to promote the use of equally efficacious but inexpensive therapies, such as steroids, fluids or pressure monitoring, is lacking. Early goal-directed therapy saves lives, and mortality increases for each hour that appropriate antibiotics and fluid re-

suscitation are delayed.<sup>2,3</sup> With any time-dependant therapy, it is necessary to expedite a continuum of care. The concepts of “chain-of-survival,” “door-to-drug time” and “taking treatment to the patient” are as relevant to sepsis as they are to acute coronary syndromes (ACS) — perhaps more so, given the high incidence, mortality and cost of severe sepsis and septic shock — yet sepsis has not received the same level of attention or funding as ACS.<sup>4,5</sup>

Just as with ACS, the first step is deciding that delays are unacceptable. Comprehensive therapy can only begin once a disease is brought to medical attention. Yet few hospitals triage septic patients in the same aggressive fashion they do for ACS. Pre-hospital sepsis care is unusual; pre-hospital cardiac care is the norm. Early and aggressive treatment of severe sepsis and septic shock will save many lives. Our challenge is to convert guidelines into meaningful clinical practice and change.<sup>6</sup> We have work to do.

**Jonathan S. Davidow, MD**

Departments of Emergency Medicine and Critical Care Medicine

**Peter G. Brindley, MD**

Department of Critical Care Medicine

**Michael J. Jacka, MD, MSc**

Departments of Critical Care Medicine and Anesthesiology & Pain Medicine

**R.T. Noel Gibney, MB**

Department Critical Care Medicine  
University of Alberta  
Edmonton, Alta.

**References**

1. Dellinger RP, Carlet JM, Masur H, et al. Surviving sepsis campaign guidelines for management of severe sepsis and septic shock. *Crit Care Med* 2004;32:858-72.
2. Ronald J, Suppes R, Gulati H, et al. The effect of timing of antimicrobial administration on mortality in septic shock patients. *Can J Anaesth* 2005;54:A4.
3. Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock.

- N Engl J Med 2001;345:1368-77.
4. American Heart Association. Part 1: Introduction to the International Guidelines 2000 for CPR and ECC: a consensus on science. *Circulation* 2000;102:1-11.
  5. Angus DC, Linde-Zwirble WT, Lidicker J, et al. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001;29:1303-10.
  6. Levy M, Pronovost J, Dellinger RP, et al. Sepsis change bundles: converting guidelines into meaningful change in behavior and clinical outcome. *Crit Care Med* 2004;32(suppl):S595-7.

## Prehospital intubation for severe head injury

*To the Editors:* We greatly appreciated the detailed, yet succinct Journal Club summary by Topping and Ducharme<sup>1</sup> of Wang and colleagues' paper<sup>2</sup> on the deleterious association demonstrated by pre-hospital intubation in the seriously head-injured patient versus emergency department intubation of a similar cohort.

Topping and Ducharme<sup>1</sup> carefully defined the population studied; the quality of the database used; the methodology for analysis (including use of a propensity score); the challenges of a possible randomized controlled trial to further delineate causation versus the clear association that has been recently demonstrated

in several emergency medical services (EMS) intubation studies, including this one; and the lessons associated with unbridled enthusiasm for unproven yet seemingly common-sense interventions (i.e., pre-hospital intubation in significantly head-injured patients).

However, one key result from this large study<sup>2</sup> seemed to elude the reviewers. In Wang and colleagues' study one group of pre-hospital providers (air medical transport crews) who used neuromuscular blocking agents had decreased mortality demonstrated in the population studied. Although Wang and colleagues qualify clear conclusions in this regard by pointing out that these 2 elements were used as covariates in the overall regression analysis, the impression is clearly given that this is an area that needs further study before the brush of nihilism for endotracheal intubation in the EMS environment is finalized. Indeed, several EMS air medical studies (observational in nature), where a small cohort of highly trained crew members are given intensive training and reasonable ongoing critical care exposure, have demonstrated exceptional airway management skills.<sup>3,4</sup> Wang and colleagues' findings are consistent with another recent study that also showed an association with improved outcomes using this air medical model.<sup>5</sup>

We feel that Wang and colleagues' suggestive data on air medical rapid sequence intubation management in the seriously ill head-injured patient deserves further consideration and is of key interest to EMS physicians and providers.

**John M. Tallon, MD**

**David Petrie, MD**

Division of EMS

Department of Emergency Medicine  
Dalhousie University, Halifax, NS

## References

1. Topping C, Ducharme J. Prehospital intubation for patients with severe head injury: More is not necessarily better. *Can J Emerg Med* 2006;8(2):116-8.
2. Wang HE, Peitzman AB, Cassidy LD, et al. Out-of-hospital endotracheal intubation and outcome after traumatic brain injury. *Ann Emerg Med* 2004;44(5):439-50.
3. Sing RF, Rotondo MF, Zonies DH, et al. Rapid sequence induction for intubation by an aeromedical transport team: a critical analysis. *Am J Emerg Med* 1998;16(6):598-602.
4. Ma OJ, Atchley RB, Hatley T, et al. Intubation success rates improve for an air medical program after implementing the use of neuromuscular blocking agents. *Am J Emerg Med* 1998;16(2):125-7.
5. Davis DP, Peay J, Serrano JA, et al. The impact of aeromedical response to patients with moderate to severe traumatic brain injury. *Ann Emerg Med* 2005;46(2):115-22.

Letters will be considered for publication if they relate to topics of interest to emergency physicians in urban, rural, community or academic settings. Letters responding to a previously published *CJEM* article should reach *CJEM* head office in Vancouver (see masthead for details) within 6 weeks of the article's publication. Letters should be limited to 400 words and 5 references. For reasons of space, letters may be edited for brevity and clarity.

Les lettres seront considérées pour publication si elles sont pertinentes à la médecine d'urgence en milieu urbain, rural, communautaire ou universitaire. Les lettres en réponse à des articles du *JCMU* publiés antérieurement devraient parvenir au siège social du *JCMU* à Vancouver (voir titre pour plus de détails) moins de six semaines après la parution de l'article en question. Les lettres ne devraient pas avoir plus de 400 mots et cinq références. Pour des raisons d'espace et par souci de concision et de clarté, certaines lettres pourraient être modifiées.