

Fish intake, erythrocyte *n*-3 fatty acid status and metabolic health in Danish adolescent girls and boys

Lotte Lauritzen^{1*}, Laurine B. S. Harsløf¹, Lars I. Hellgren², Maiken H. Pedersen^{1,2}, Christian Mølgaard¹ and Kim F. Michaelsen¹

¹Department of Human Nutrition, Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark

²Department of Systems Biology, the Technical University of Denmark, Lyngby, Denmark

(Received 16 March 2010 – Revised 30 June 2010 – Accepted 26 July 2010 – First published online 7 June 2011)

Abstract

Marine *n*-3 long-chain PUFA (*n*-3 LCPUFA) may have a beneficial effect on several aspects of the metabolic syndrome (dyslipidaemia, insulin resistance, hypertension and abdominal obesity). The metabolic syndrome is increasing in prevalence during adolescence, but only few studies have investigated the effects of *n*-3 LCPUFA in adolescence. The present study examines associations between fish intake (assessed by a 7 d pre-coded food diary), erythrocyte (RBC) DHA status (analysed by GC) and metabolic syndrome measures (anthropometry, blood pressure and plasma lipids, insulin and glucose) in 109 17-year-old children from the Copenhagen Birth Cohort Study. Of the children, 8% were overweight or obese and few showed signs of the metabolic syndrome, but all the metabolic syndrome variables were correlated. Median fish intake was 10.7 (interquartile range 3.6–21.2) g/d. Boys tended to have a higher fish intake ($P=0.052$), but girls had significantly higher RBC levels of DHA ($P=0.001$). Sex and fish intake explained 37% of the variance in RBC-DHA ($P<0.001$). After adjusting for confounders, high DHA status was found to be significantly correlated with higher systolic blood pressure ($P=0.014$) and increased fasting insulin ($P=0.018$), but no adverse association was observed with the mean metabolic syndrome *z*-score. Overall, the present study showed the expected association between fish intake and RBC-DHA, which in contrast to our expectations tended to be associated with a poorer metabolic profile. Whether these results reflect the physiological function of *n*-3 LCPUFA, lifestyle factors associated with fish intake in Denmark, or mere chance remains to be investigated.

Key words: *n*-3 PUFA: Blood pressure: Insulin sensitivity: Plasma lipid profile

The incidence of metabolic dysregulation is an increasing problem, now even occurring during adolescence. Metabolic dysregulation is characterised by dyslipidaemia, insulin resistance, hypertension and abdominal obesity – the so-called metabolic syndrome – which predisposes toward the development of CVD and type 2 diabetes. Dietary *n*-3 long-chain PUFA (*n*-3 LCPUFA) from marine foods have in adults been shown to have a beneficial effect on blood pressure (BP)⁽¹⁾ and blood lipid profile, specifically plasma TAG⁽²⁾. Furthermore, *n*-3 LCPUFA have been shown to prevent the development of insulin resistance induced by a high-fat or high-sucrose diet in rodents^(3,4). Some studies have shown associations between *n*-3 LCPUFA and better insulin sensitivity in human subjects^(5,6) and recent randomised trials have also shown beneficial effects of fish oil supplementation^(7–10). However, the results concerning the relationship between *n*-3 LCPUFA and insulin sensitivity in human subjects are not at present convincing^(11,12). A possible explanation for the

inconsistencies could be that the effect may be different in established insulin resistance and during its development.

It is at present believed that infants have higher needs for *n*-3 PUFA due to their effect on the development of the central nervous system⁽¹³⁾. In Denmark, it is recommended that infants ingest a minimum of 1% of their energy (percentage of energy; E%) as *n*-3 PUFA, as opposed to 0.5 E% from 2 years of age. We have previously shown that fish oil supplementation of 9-month-old infants have a beneficial effect on plasma lipid profile and BP⁽¹⁴⁾. This result surprised us since the infants had an overall low cardiovascular risk profile and the adult studies tend only to find an effect of fish oil in subjects with a high risk profile⁽¹⁾. It is our hypothesis that the results could reflect a greater susceptibility towards effects of *n*-3 LCPUFA intake in the infant growth phase due to a higher *n*-3 PUFA need and that effects of *n*-3 LCPUFA intake would also be detectable in other growth phases and on other characteristics of the metabolic syndrome.

Abbreviations: BP, blood pressure; E%, percentage of energy; FA%, percentage of all fatty acids in the chromatogram; HDL-C, HDL-cholesterol; HOMA-IR, homeostasis model assessment of insulin resistance; LCPUFA, long-chain PUFA; RBC, erythrocyte; SBP, systolic blood pressure.

* **Corresponding author:** Dr Lotte Lauritzen, fax +45 3533 2483, email ll@life.ku.dk

In the present study, we examined if *n*-3 LCPUFA status or intake in adolescents was associated with biomarkers of the metabolic syndrome in a cross-sectional analysis within the prospective Copenhagen Birth Cohort Study. The prospective design enabled us to take prognostic variables (for example, birth weight, duration of breast-feeding and socio-economic factors) into account as well as possible confounding lifestyle factors (diet and physical activity). Furthermore, we examined associations between erythrocyte (RBC) *n*-3 LCPUFA and fish intake.

Methods

The Copenhagen Cohort Study is a prospective observational study where a random sample of Danish infants born from 1987 to 1988 was followed during the first year of life and later followed up at 10 and 17 years. The inclusion criteria were: parents of Danish origin, singleton birth, 37–42 weeks' gestation, normal birth weight (10th–90th percentiles for gestational age), no neonatal disease or malformation. Of the 251 infants who fulfilled the inclusion criteria, 145 infants completed the study, fifty-nine of which were in a control group only examined at 9 months, whereas the rest were examined several times during the first year of life⁽¹⁵⁾. They were all invited to take part in this 17-year follow-up and 109 (75%) agreed to participate. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki and was approved by the Ethics Committee for Copenhagen and Frederiksberg (no. KF 01-226/97). Written informed consent was obtained from all participants and their parents.

Dietary intake was determined in the week before the examination visit by a 7 d food record with pre-coded response categories and open supplementary alternatives as previously described⁽¹⁶⁾. Intake was registered in household measures and portion sizes based on an accompanying photograph series. From these records, intake of energy, macronutrients and micronutrients and that of specific food categories (fish, meat, dairy products, fruits and vegetables, cereals and sweets) was calculated using the national food database. Fish intake was assessed by twelve individual variables; six for lunch and six for dinner which distinguished between fatty and lean fish and seafood: lean fish for dinner, fatty fish for dinner, fish balls, sushi, fish stews, shrimps as main course, sandwich with breaded fillet of sole or similar, tuna in salad or on bread, herring, mackerel and other kinds of fatty fish on bread, cod roe, and seafood as main dish or garnish. Physical activity was determined by a 24 h recall questionnaire (habitual activity estimation scale) as previously described⁽¹⁷⁾.

Height was measured to the nearest 1 mm by a stadiometer (Chasmore, London, UK) and weight was measured to the nearest 0.1 kg by an electronic scale (Lideltronic 8000; Lindells Inc., Kristianstad, Sweden). Waist and hip circumferences were measured in triplicate according to standard procedures. Body fat percentage was determined by dual-energy X-ray absorptiometry on a Hologic QDR 1000/W with software version 5.73 (Hologic Inc., Waltham, MA, USA). Brachial BP was

assessed after 10 min of supine rest by oscillometry using the average of three BP recordings.

RBC isolated from a fasting heparinised blood sample was used to assess the RBC fatty acids composition by GLC as previously described⁽¹⁸⁾. Fatty acid composition is given as the percentage of individual fatty acids relative to the total chromatogram (FA%). Metabolism ratios were calculated by ratios between the fatty acids immediately before a Δ 6-desaturation step and those after (also including the last elongation and post-desaturation oxidation step in the LCPUFA synthesis, 18:2*n*-6:18:3*n*-6, 18:3*n*-3:18:4*n*-3, 22:4*n*-6:22:5*n*-6, 22:5*n*-3:22:6*n*-3 and the combined ratio).

Total cholesterol, HDL-cholesterol (HDL-C) and TAG concentrations of fasting EDTA plasma were measured with enzymic kits from Roche (CHOD-PAP, HDL-C-plus 2nd generation, and GPO-PAP, respectively) on a Cobas Mira Plus analyser (Roche Diagnostic, Basel, Switzerland). The LDL-cholesterol concentrations were calculated using the Friedewald formula⁽¹⁹⁾. Fasting glucose was measured with the use of the hexokinase endpoint method (Gluco-quant Glucose/HK; Roche Diagnostics, Basel, Switzerland) using an ABX Pentra 400 chemistry analyser (ABX Pentra, Horiba ABX, Montpellier, France). Fasting serum insulin was measured by a solid-phase, two-site chemiluminescent immunometric assay with the use of an Immulite 1000 analyzer (Diagnostic Products Corp., Los Angeles, CA, USA). Homeostasis model assessment of insulin resistance (HOMA-IR) was calculated by the HOMA2 model using the HOMA Calculator[®] (version 2.2.2; University of Oxford, 2004, <http://www.dtu.ox.ac.uk/homacalculator/>). We used standard adult cut-offs according to the American National Institute of Health's National Cholesterol Education Program⁽²⁰⁾ to determine the presence of the metabolic syndrome using three out of the five following criteria: waist > 102 cm for boys and > 88 cm for girls, plasma TAG > 1.7 mmol/l, HDL-C < 0.9 mmol/l for boys and < 1.0 mmol/l for girls, glucose > 6.1 mmol/l and BP > 130/85 mmHg. Furthermore, for each of the metabolic syndrome variables (waist, systolic BP (SBP), TAG, HDL-C and HOMA-IR) *z*-scores were calculated based on the mean and the standard deviation in the relevant sex subgroup of the participants in such a way that a positive *z*-score for all variables indicates higher risk. A combined metabolic syndrome *z*-score was calculated as the mean of the *z*-scores of all metabolic syndrome variables (the *z*-score of HDL-C was used with an opposite sign since a high HDL-C concentration is considered beneficial).

Statistics

All statistical analysis was performed by SPSS 16.0 (SPSS Inc., Chicago, IL, USA). Normal distribution of all variables was checked by Kolmogorov–Smirnov and by visual inspection of histograms. The following variables were not normally distributed and thus log-transformed before analysis: plasma cholesterol, TAG, RBC 20:3*n*-9, RBC-EPA, intake of energy, fish, Na, dietary fibres and sugar. Fasting insulin was used as a reverse value. All descriptive results are given as mean values and standard deviations or median (25th and 75th percentile) and outcome parameters are given as mean values

with their standard errors. Group comparisons (participants *v.* non participants and boys *v.* girls) were performed by Student's *t* test or, for non-normally distributed variables, by the Mann–Whitney *U* test. Associations between fish intake (total or individual fish intake parameters) and RBC *n*-3 LCPUFA were checked by backward multiple regression analysis and the final model was chosen as the one with maximum explained variance and the fewest independent variables. The participants were divided in tertiles according to RBC-DHA status with an even distribution of girls and boys by a separate division of boys and girls and a subsequent pooling of the sex subgroups in each of the tertiles. Associations between RBC-DHA status (high, medium and low as determined by tertiles in the population) and various outcome parameters (diet, body composition parameters, and the other metabolic syndrome variables) were assessed by logistic regression analysis. All models were adjusted for sex, duration of breast-feeding and diet (and body composition for all non-body composition variables) plus for the individual blood lipids with further adjustment of the corresponding blood lipid during the first year of life (mean of values at 2, 4 and 9 months). Outcomes that were found to be significantly associated with RBC-DHA status were also analysed in relation to RBC-DHA as a continuous variable and the same adjustments.

Results

The 109 children, who took part in this follow-up substudy, were examined at a mean age of 17.3 (SD 0.2) years. The characteristics of the included children are shown in Table 1. The participating children did not differ from the non-participating children with respect to any of the given characteristics (data not shown). Only approximately 8% of the children were overweight or obese according to waist

circumference cut-offs or BMI and few exceeded the cut-off for one or more of the other metabolic syndrome criteria: nine had high plasma TAG, eleven had low HDL-C (four of whom also had high TAG), seven had total plasma cholesterol > 5 mmol/l, none were hypertensive, five had SBP > 130 mmHg, and none was hyperglycaemic. Of the children, two had the metabolic syndrome according to the 'three out of five' National Cholesterol Education Program definition⁽²⁰⁾. There were significant (sex-adjusted) bivariate associations between all of the metabolic syndrome variables.

Table 2 gives the overall fatty acid composition of the RBC of the children. Girls had significantly higher levels of DHA and the PUFA composition generally points to a better Δ 6-desaturation capacity with a significant lower ratio between pre- and post- Δ 6-desaturase fatty acids (2.4 (SD 0.5) *v.* 3.0 (SD 0.7) in boys; $P < 0.001$), especially the 22:5*n*-3:DHA ratio (0.4 (SD 0.1) *v.* 0.6 (SD 0.1); $P < 0.001$). After adjustment for fish intake, RBC-DHA status was estimated to be 1.0 (SD 0.2) FA% ($P < 0.001$) higher in girls.

Median fish intake was 10.7 (interquartile range 3.6–21.2) g/d and tended to be higher in boys (14.3 *v.* 8.0 g/d in girls; $P = 0.052$; Mann–Whitney). Overall, fish intake was distributed as follows: 21% of the total amount of fish came from fatty fish for dinner, 25% from dishes with lean fish for dinner, 9% from mixed fish dishes (sushi, etc.), 10% from fatty fish sandwiches for lunch, 10% from sandwiches with lean fish and 14% from seafood – maximally 14% of the fish was breaded. There was a significant association between the intake of fish at 17 years and that at 10 years ($r = 0.195$; $P = 0.007$; $n = 92$) and also a slight tendency towards an association with intake of fish at 9 months of age ($r = 0.124$; $P = 0.101$; $n = 91$).

RBC-DHA was correlated with total fish intake adjusted for sex ($r = 0.488$; $P < 0.001$) and so was the RBC content of *n*-3 LCPUFA ($r = 0.341$; $P = 0.001$). Of the variance in RBC-DHA,

Table 1. Characteristics of the participating 17-year-old boys and girls (Mean values and standard deviations or medians and 25th–75th percentiles)

	Boys		Girls		<i>P</i>
	Mean	SD	Mean	SD	
Subjects (<i>n</i>)	44		65		
Birth weight (kg)	3.52	0.38	3.36	0.35	0.021
Length at birth (cm)	52.1	1.8	51.2	2.0	0.296
Duration of breast-feeding (months)					0.454
Median	5.0		7.0		
25th–75th percentile	2.0–8.8		3.0–9.0		
Intake of fish at 9 months (g/d)					0.381
Median	2.9		1.6		
25th–75th percentile	0–11.8		0–7.6		
Maternal BMI (kg/m ²)	20.9	2.6	21.5	2.8	0.298
Paternal BMI (kg/m ²)	23.6	3.0	23.8	2.7	0.773
Maternal smoking in first year (cigarettes/d)					0.540
Median	3		0		
25th–75th percentile	0–12		0–12		
Mean plasma cholesterol in first year (mmol/l)	4.1	0.6	4.3	0.7	0.192
Mean LDL-cholesterol in first year (mmol/l)	2.1	0.7	2.6	1.2	0.052
Mean HDL-cholesterol in first year (mmol/l)	0.98	0.26	1.28	1.25	0.239
Mean TAG in first year (mmol/l)	2.2	1.1	1.7	0.6	0.019
Height at 17 years (m)	1.83	0.06	1.67	0.06	<0.001
Weight at 17 years (kg)	72.6	10.6	59.4	10.6	<0.001
BMI at 17 years (kg/m ²)	21.62	3.05	21.30	3.15	0.597

Table 2. Fatty acid composition of erythrocytes from 17-year-old boys and girls*
(Mean values and standard deviations)

	Boys		Girls		P
	Mean	SD	Mean	SD	
Subjects (n)	44		65		
SFA	41.1	0.55	41.3	0.90	0.175
MUFA	17.1	0.83	16.8	0.82	0.066
18:2n-6	10.8	1.15	10.2	0.99	0.003
18:3n-3	0.19	0.05	0.18	0.04	0.784
20:2n-6	0.24	0.03	0.26	0.04	0.014
20:3n-9	0.11	0.03	0.11	0.02	0.835
20:3n-6	1.97	0.38	1.94	0.41	0.686
20:4n-6	15.5	1.34	15.7	0.98	0.324
20:5n-3	0.79	0.28	0.74	0.32	0.160
22:4n-6	3.16	0.60	3.21	0.59	0.632
22:5n-6	0.61	0.13	0.70	0.23	0.008
22:5n-3	2.73	0.21	2.31	0.31	0.001
22:6n-3	4.99	1.21	5.82	0.99	0.001
PUFA	41.3	0.93	41.43	0.93	0.390
n-3 Long-chain PUFA	8.62	1.51	9.00	1.23	0.152
n-6:n-3 PUFA	3.70	0.78	3.50	0.61	0.161

* Values are given as the percentages of individual fatty acids relative to the total chromatogram area (FA%).

37% was explained by sex and the last week's fish intake, with the most significant contributions from lean fish for dinner, tuna, sushi, sole fillet sandwich, fish balls, and fatty fish for dinner. Inclusion of the overall intake of PUFA (essentially equivalent to that of linoleic and α -linolenic acids) did not change this pronouncedly (39% explained variance).

When divided according to RBC-DHA status by sex-adjusted tertiles, the RBC-DHA groups were found to differ significantly with respect to duration of breast-feeding, but not in any of

the other prognostic or lifestyle-dependent variables at birth (Table 3). Furthermore, sex-adjusted dietary intake in the high, medium and low RBC-DHA status groups differed significantly with respect to PUFA E% ($P=0.041$) and there was a tendency towards a difference in the Na intake ($P=0.077$) and the protein:carbohydrate ratio of the diet ($P=0.123$). The RBC-DHA groups did not differ significantly with respect to parental BMI (data not shown) or body composition in analysis adjusted for sex, breast-feeding and dietary factors (total energy intake, PUFA E% and protein:carbohydrate) (Table 4). However, after adjusting for sex, breast-feeding, dietary factors and body fat percentage there were significant differences in SBP and fasting insulin and tendencies in HOMA-IR, heart rate and plasma total cholesterol after further adjustment for plasma cholesterol during first year of life (Table 4). The degree of explained variance in the models was 20–30% for the HOMA-IR and insulin parameters, about 20% for the blood lipids, and 43% for SBP. The overall outcome of these analyses was not changed by inclusion of physical activity or food patterns (for example, intake of fruits and vegetables). SBP was also associated with RBC-DHA as a continuous variable in a multiple regression analysis including the same adjusting factors as above ($B = 1.7$ (SEM 0.6) mmHg/FA% ($P=0.005$) and HOMA-IR tended to be associated with RBC-DHA as well ($P=0.052$)). However, no association was observed between RBC-DHA and the mean metabolic syndrome z -score ($P=0.60$). The overall picture and conclusions were the same (although less significant) in similar analyses with RBC-EPA and fish intake.

Table 3. Diet and lifestyle at 17 years of age and infant characteristics in sex-adjusted tertiles according to erythrocyte DHA status
(Mean values with their standard errors or medians and 25th–75th percentiles)

Tertile...	Low		Medium		High		P
	Mean	SEM	Mean	SEM	Mean	SEM	
Subjects (n)	36		37		36		
Sex (% girls)	58		62		58		
Birth weight (kg)	3.35	0.06	3.50	0.06	3.41	0.06	0.227
Duration of breast-feeding (months)							0.002
Median	4		5		9		
25th–75th percentile	1–8		3–8		5–9		
Fish intake at 9 months (g/d)							0.276
Median	0		1		4		
25th–75th percentile	0–4		0–10		0–11		
HAES physical activity score	4.1	0.2	4.1	0.2	3.8	0.2	0.333
Energy intake (MJ/d)	9.3	0.5	10.2	0.6	10.4	0.6	0.122
Carbohydrate (E%)	53.1	0.7	51.3	0.8	52.1	0.7	0.314
Protein (E%)	13.1	0.3	13.9	0.3	13.9	0.4	0.137
Fat (E%)	30.9	0.8	32.0	0.8	31.7	0.6	0.419
Fish and fish products (g/d)	8	2	16	3	23	3	0.001
Girls (g/d)	8	3	10	2	20	4	
Boys (g/d)	8	2	25	8	27	6	
Meat and poultry (g/d)	136	9	142	10	142	9	0.539
Dairy products (g/d)	461	64	461	83	452	64	0.916
Fruits and vegetables (g/d)	532	48	423	32	596	72	0.462
Cereals and bread (g/d)	210	13	238	15	246	18	0.082
Fats (g/d)	30	3	33	3	36	4	0.193
Sweets and sugar products (g/d)	50	5	44	4	51	5	0.911

HAES, habitual activity estimation scale; E%, percentage of energy.

Table 4. Metabolic syndrome variables at 17 years of age in sex-adjusted tertiles according to erythrocyte DHA status

(Mean values with their standard errors or medians and 25th–75th percentiles)

Tertile...	Low		Medium		High		P*
	Mean	SEM	Mean	SEM	Mean	SEM	
Subjects (n)	36		37		36		
Height (m)	1.74	0.02	1.72	0.02	1.74	0.02	0.662
Weight (kg)	64.4	1.8	62.4	2.0	67.5	2.3	0.252
BMI (kg/m ²)	21.3	0.5	20.9	0.4	22.1	0.6	0.332
Waist circumference (cm)	79.0	1.4	77.1	1.4	80.2	1.8	0.734
Body fat (%)	20.9	1.4	20.3	1.0	22.7	1.3	0.319
Fasting glucose (mmol/l)	4.48	0.06	4.52	0.05	4.56	0.06	0.649
Fasting insulin (pmol/l)							0.018
Median	51		52		62		
25th–75th percentile	40–67		43–78		49–86		
HOMA-IR	1.03	0.09	1.02	0.07	1.44	0.21	0.052
Diastolic blood pressure (mmHg)	53.5	0.8	52.5	0.9	54.5	1.0	0.333
Systolic blood pressure (mmHg)	113.8	1.4	113.6	1.3	117.9	1.4	0.014
Heart rate (beats per min)	63.7	9.3	63.2	7.8	69.0	8.5	0.062
Cholesterol (mmol/l)							0.064
Median	3.31		3.42		3.58		
25th–75th percentile	2.93–3.90		3.16–4.06		3.15–4.12		
LDL-cholesterol (mmol/l)	2.26	0.18	2.23	0.10	2.37	0.11	0.365
HDL-cholesterol (mmol/l)	1.29	0.05	1.46	0.05	1.46	0.07	0.287
TAG (mmol/l)							0.474
Median	0.86		0.89		0.87		
25th–75th percentile	0.62–1.12		0.61–1.13		0.66–1.06		
Mean MS z-score	0.01	0.09	–0.18	0.05	0.04	0.09	0.958

HOMA-IR, homeostasis model assessment of insulin resistance; MS z-score, mean metabolic syndrome z-score;

* The statistical analysis of anthropometric variables was adjusted for sex, duration of breast-feeding and diet (percentage of energy from PUFA and proteins:carbohydrates ratio). These covariates were also used in the models for the other metabolic syndrome markers along with body fat percentage, and blood lipids in the first year for plasma lipids and intake of Na for blood pressure.

Discussion

The results of the present study showed that teenage girls have higher levels of DHA in their RBC than boys and that fish intake in both sexes was strongly associated with RBC *n-3* LCPUFA status. In contrast to our expectations, higher RBC-DHA was not found to be associated with lower markers of the metabolic syndrome, but rather a higher BP and lower insulin sensitivity compared with adolescents with a lower RBC DHA status. However, after adjustment for confounders, no adverse association was observed with the mean metabolic syndrome z-score.

The positive association between intake of fish and RBC *n-3* LCPUFA status was in accordance with our expectations. Intake of fish explained approximately 40% of the variation in RBC-DHA, which is in accordance with that in previous studies, for example, in pregnant women⁽²¹⁾. To our knowledge, no studies have verified the fish intake–RBC biomarker relationship in children, but the present study indicates only minor potential for effect modification by age and sex. The low degree of explained variance may primarily be because the periods of diet registration were short compared with the day-to-day variation in fish intake. Furthermore, the 7 d food record was only semi-quantitative. The uncertainty of the present study is therefore presumed to be mainly related to the dietary registration, while assessment of RBC *n-3* LCPUFA is considered a stable measure of fish intake during the last couple of months. We therefore believe that RBC

n-3 LCPUFA is the most relevant measure to use when associations with the functional outcomes are examined. Surprisingly, we found that intake of lean fish had a significant impact on DHA status at 17 years. This could be explained by the fact that the fish intake in this group of teenagers was relatively low (approximately half the amount compared with Danish adults) and primarily consumed for dinner (66% was fish for dinner), which typically consists of lean fish in Denmark while fatty fish such as herring and mackerel more often are consumed for lunch. Although the average fish intake in this group of adolescents is low, the range of fish intake was quite wide (total intake of fish ranged between 0 and 120 g/d) and spans the whole spectrum of fish intake that has previously been reported to have beneficial effects. The surprising adverse association between fish intake and some metabolic syndrome markers in the present study may be explained by high fast food intakes (for example, fish and chips) during adolescence. However, that was not supported by the food records. Furthermore, breaded fish filets also contributed to the DHA status, which probably means that the filets were only lightly breaded and not deep-fried, since most of the fat in commercially breaded deep-fried fish products would be SFA, MUFA and *n-6* PUFA. It has been hypothesised that intake of linoleic acid relative to that of PUFA may blunt the tissue incorporation of *n-3* LCPUFA⁽²²⁾. However, adjustment for intake of PUFA did not play a role in relation to the association between fish intake and RBC-DHA. This is in agreement with the incorporation

results from a recent two-factorial trial where we supplemented adult men simultaneously with both *n*-3 LCPUFA and linoleic acid⁽²³⁾; nevertheless, such an effect has been shown in plasma from children⁽²⁴⁾. Interestingly, this cohort study results show significant associations between fish intake throughout childhood, indicating stability in the dietary pattern and validity in the dietary assessments. This, however, also means that we cannot distinguish between whether the associations we found between status and health reflect a relationship with the present fish intake or a programming effect. The observed association between duration of breast-feeding and DHA status at 17 years of age could be a programming effect but is more likely due to lifestyle associations (between breast-feeding and fish intake). No significant association was observed of breast-feeding on RBC fatty acid composition after adjustment for sex and fish intake (data not shown).

Previous studies have shown that fish oil lowers plasma TAG and BP, although the effect on BP only seems to occur among people who have an unfavourable risk profile⁽¹⁾ and among infants⁽¹⁴⁾. Furthermore, it has been shown that *n*-3 PUFA inhibit development of obesity and insulin resistance induced by a high-fat and -sucrose diet in rodents^(3,4). A high serum *n*-3 PUFA status has also in adult men been shown to decrease the relative risk for developing the metabolic syndrome over the following 20 years, independent of smoking habits, physical activity and BMI⁽²⁵⁾. Much to our surprise, the present cross-sectional study showed a potentially inverse relationship between DHA status and BP and insulin sensitivity in adolescents. A lack of an association might be anticipated due to lack of power, the overall health status of the subjects or too low a dose. However, we have previously in a follow-up study to a randomised controlled trial with fish oil supplementation during lactation seen similar adverse long-term effects on BP⁽²⁶⁾, but this is in contrast to the programming effect of perinatal intake of *n*-3 PUFA observed in rodents⁽²⁷⁾ and formula-fed infants⁽²⁸⁾. The children in our randomised controlled trial were also healthy Danish adolescents, but the *n*-3 LCPUFA dose was in the higher end of the population intake range and the difference in intake was limited to a period of the first 4 months of lactation. Most of the children in the present study had BP, plasma lipid profile and markers of glucose homeostasis within the normal range and the overall negative effects were small. The long-term implications may, however, be clinically relevant as these risk markers have been shown to track and a high value could then translate into a clinically increased level later in life. The observed adverse associations could in theory be due to various pollutants in the fish⁽²⁹⁾. The fish intake in the examined population was, however, not very high and other studies have also found beneficial association between plasma levels of marine pollutants and CVD risk factors, indicating that the benefits outweigh the potential risks⁽³⁰⁾. Fish intake is typically related to a healthy diet and lifestyle and in the present study there was also a positive correlation with intake of fruit and vegetables, indicating that fish consumption is also related to a generally healthy diet among these teenagers, which makes the lack

of a beneficial association between RBC-DHA and markers of the metabolic syndrome even more odd. It seems implausible that the active components of a healthy diet should have markedly different physiological effects in teenagers compared with in adults. The adverse relationship between RBC-DHA and BP and markers of insulin sensitivity could, however, be due to hidden confounding factors that we did not measure or that other aspects of the teenage lifestyle were unhealthy – too much food and too little physical activity. Inclusion of the possible confounding factors that we did have information on (for example, physical activity and intake of fruits and vegetables) did not affect the adverse association. We have also taken the protein:carbohydrate ratio in the diet into account, as this has been found to modulate the effect of PUFA in rats^(31,32). This factor played a significant role for some of the outcomes but not as an effect modifier of the effect of DHA status.

There was a significant difference in the RBC fatty acid composition between boys and girls, with higher DHA levels in girls in spite of a lower fish intake. There was also a significant higher RBC incorporation of 22:5*n*-6, the *n*-6 LCPUFA synthesis endproduct, and a lower RBC Δ 6-desaturase substrate:product-ratio, specifically in the ratio between 22:5*n*-3 and DHA. Thus, overall, the sex difference could reflect a higher Δ 6-desaturase capacity in girls. This is the first study to show sex differences in PUFA levels in adolescents, but similar differences in fatty acid composition have previously been observed in blood lipids of men and women^(33–35). Furthermore, women have been shown to have a higher rate of conversion from α -linolenic acid to *n*-3 LCPUFA⁽³⁶⁾. Sex differences have also been observed in the tissue fatty acid composition in rats, but Burdge *et al.* found no sex difference in rat hepatic Δ 6-desaturase expression and no association between the proportion of individual PUFA in hepatic phospholipids and Δ 6- or Δ 5-desaturase expression^(37–39). Genetic differences in fatty acid desaturase (FADS) 1–2, the gene cluster that codes for Δ 5- and Δ 6-desaturases in man, have been shown to be reflected in tissue PUFA levels, mostly so in the variation in 20:2*n*-6 and 20:4*n*-6 (arachidonic acid)^(40,41). The final LCPUFA synthesis step to DHA and 22:5*n*-6 involves more than a Δ 6-desaturation and it could be these other steps that were up-regulated. What ever the mechanism, tissue PUFA composition has been found to be manipulated by sex hormones^(33,36,42). The fact that girls have a higher endogenous capacity for synthesis of DHA could presumably mean that they were less dependent or less affected by intake. This idea is in line with the fact that people with a high genetically determined capacity (a specific allele composition in the FADS2 gene) have been shown to be less affected by intake – for example, the effect of breast-feeding *v.* formula feeding on intelligence quotient (IQ)⁽⁴³⁾.

This is among the first studies to examine the relationship between intake of fish and *n*-3 LCPUFA on markers of health in adolescents. The strength of our cohort study is that it had an initial random selection and a relatively high rate of follow-up (i.e. 75% of all the children who were recruited at birth). However, since it is an observational study we cannot exclude that our findings could be due to

residual confounding or atypical associations between fish intake and lifestyle in this particular group of teenagers. In summary, the results from this cross-sectional analysis in our cohort showed that sex and fish intake explain much of the variance in adolescent RBC-DHA status, boys tend to have a higher fish intake, but girls had higher RBC-DHA status. Furthermore, in contrast to expectations, DHA status was found to be associated with higher BP and lower insulin sensitivity, but not with the overall metabolic syndrome z -score. Whether these results are a mere chance finding or whether they reflect a truly different effect of n -3 LCPUFA during growth needs to be investigated by more studies, preferably intervention studies, on the relationship between n -3 LCPUFA intake and health in adolescence.

Acknowledgements

We gratefully acknowledge Majken Ege and Helga Kristensen Ingstrup for collecting the data, Vivian Anker and Birgitte Hermansen for technical assistance and Janni Felskov Agersten for performing the RBC fatty acid analysis.

L. L., K. F. M. and C. M. were responsible for the study idea and design, and L. I. H. for the fatty acid analysis. L. L. and L. B. S. H. conducted the statistical analyses and L. L. prepared the first draft of the manuscript. All authors contributed to the interpretation of the results and were involved in preparing the final manuscript. The present study was funded by The Danish Heart Association.

None of the authors had any financial or personal conflicts of interest.

References

- Geleijnse JM, Giltay EJ, Grobbee DE, *et al.* (2002) Blood pressure response to fish oil supplementation: meta-regression analysis of randomized trials. *J Hypertens* **20**, 1493–1499.
- Hooper L, Thompson RL & Harrison RA, *et al.* (2004) Omega 3 fatty acids for prevention and treatment of cardiovascular disease. *Cochrane Database of Systematic Reviews, Issue 4*, CD003177. <http://www.mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD003177/frame.html>
- Ghafoorunissa, Ibrahim A, Rajkumar L, *et al.* (2005) Dietary (n -3) long-chain polyunsaturated fatty acids prevent sucrose-induced insulin resistance in rats. *J Nutr* **135**, 2634–2638.
- Storlien LH, Kraegen EW, Chisholm DJ, *et al.* (1987) Fish oil prevents insulin resistance induced by high-fat feeding in rats. *Science* **237**, 885–888.
- Haugaard SB, Madsbad S, Høy CE, *et al.* (2006) Dietary intervention increases n -3 long-chain polyunsaturated fatty acids in skeletal muscle membrane phospholipids of obese subjects. Implications for insulin sensitivity. *Clin Endocrinol* **64**, 169–178.
- Clore JN, Harris PA, Li J, *et al.* (2000) Changes in phosphatidylcholine fatty acid composition are associated with altered skeletal muscle insulin responsiveness in normal man. *Metab Clin Exp* **49**, 232–238.
- Delarue J, Li CH, Cohen R, *et al.* (2006) Interaction of fish oil and a glucocorticoid on metabolic responses to an oral glucose load in healthy human subjects. *Br J Nutr* **95**, 267–272.
- Spadaro L, Magliocco O, Spampinato D, *et al.* (2008) Effects of n -3 polyunsaturated fatty acids in subjects with nonalcoholic fatty liver disease. *Digest Liver Dis* **40**, 194–199.
- Ramel A, Martinez A, Kiely M, *et al.* (2008) Beneficial effects of long-chain n -3 fatty acids included in an energy-restricted diet on insulin resistance in overweight and obese European young adults. *Diabetologia* **51**, 1261–1268.
- Browning LM, Krebs JD, Moore CS, *et al.* (2007) The impact of long-chain n -3 polyunsaturated fatty acid supplementation on inflammation, insulin sensitivity and CVD risk in a group of overweight women with an inflammatory phenotype. *Diabet Obes Metab* **9**, 70–80.
- Riserus U, Willett WC & Hu FB (2009) Dietary fats and prevention of type 2 diabetes. *Prog Lipid Res* **48**, 44–51.
- Hartweg J, Farmer AJ, Holman RR, *et al.* (2009) Potential impact of omega-3 treatment on cardiovascular disease in type 2 diabetes. *Curr Opin Lipidol* **20**, 30–38.
- Lauritzen L, Hansen HS, Jørgensen MH, *et al.* (2001) The essentiality of long-chain n -3 fatty acids in relation to development and function of the brain and retina. *Prog Lipid Res* **40**, 1–94.
- Damsgaard CT, Schack-Nielsen L, Michaelsen KF, *et al.* (2006) Fish oil affects blood pressure and the plasma lipid profile in healthy Danish infants. *J Nutr* **136**, 94–99.
- Michaelsen KF (1997) Nutrition and growth during infancy. The Copenhagen Cohort Study. *Acta Paediatr Suppl* **420**, 1–36.
- Budek AZ, Hoppe C, Ingstrup H, *et al.* (2007) Dietary protein intake and bone mineral content in adolescents – The Copenhagen Cohort Study. *Osteoporos Int* **18**, 1661–1667.
- Mølgaard C, Thomsen BL & Michaelsen KF (2001) The influence of calcium intake and physical activity on bone mineral content and bone size in healthy children and adolescents. *Osteoporos Int* **12**, 887–894.
- Lauritzen L, Jørgensen MH, Mikkelsen TB, *et al.* (2004) Maternal fish oil supplementation in lactation: effect on visual acuity and n -3 fatty acid content of infant erythrocytes. *Lipids* **39**, 195–206.
- Friedewald WT, Levy RI & Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* **18**, 499–502.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (2001) Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* **285**, 2486–2497.
- Olsen SE, Hansen HS, Sandström B, *et al.* (1995) Erythrocyte levels compared with reported dietary intake of marine n -3 fatty acids in pregnant women. *Br J Nutr* **73**, 387–395.
- Hibbeln JR, Nieminen LR, Blasbalg TL, *et al.* (2006) Healthy intakes of n -3 and n -6 fatty acids: estimations considering worldwide diversity. *Am J Clin Nutr* **83**, S1483–S1493.
- Damsgaard CT, Frøkiær H & Lauritzen L (2008) The effects of fish oil and high or low linoleic acid intake on fatty acid composition of human peripheral blood mononuclear cells. *Br J Nutr* **99**, 147–154.
- Hoyos C, Almqvist C, Garden F, *et al.* (2008) Effect of omega 3 and omega 6 fatty acid intakes from diet and supplements on plasma fatty acid levels in the first 3 years of life. *Asia Pacific J Clin Nutr* **17**, 552–557.
- Warensjö E, Sundström J, Lind L, *et al.* (2006) Factor analysis of fatty acids in serum lipids as a measure of dietary fat quality in relation to the metabolic syndrome in men. *Am J Clin Nutr* **84**, 442–448.

26. Asserhøj M, Nehammer S, Matthiessen J, *et al.* (2009) Maternal fish oil supplementation during lactation may adversely affect long-term blood pressure, energy intake, and physical activity of 7-year-old boys. *J Nutr* **139**, 298–304.
27. Armitage JA, Pearce AD, Sinclair AJ, *et al.* (2003) Increased blood pressure later in life may be associated with perinatal *n*-3 fatty acid deficiency. *Lipids* **38**, 459–464.
28. Forsyth JS, Willatts P, Agostoni C, *et al.* (2003) Long-chain polyunsaturated fatty acid supplementation in infant formula and blood pressure in later childhood: follow up of a randomised controlled trial. *BMJ* **326**, 953–957.
29. Lee DH, Lee IK, Porta M, *et al.* (2007) Relationship between serum concentrations of persistent organic pollutants and the prevalence of metabolic syndrome among non-diabetic adults: results from the National Health and Nutrition Examination Survey 1999–2002. *Diabetologia* **50**, 1841–1851.
30. Mozaffarian D (2009) Fish, mercury, selenium and cardiovascular risk: current evidence and unanswered questions. *Int J Environ Res Public Health* **6**, 1894–1916.
31. Madsen L, Petersen RK & Kristiansen K (2005) Regulation of adipocyte differentiation and function by polyunsaturated fatty acids. *Biochim Biophys Acta-Mol Basis Dis* **1740**, 266–286.
32. Madsen L, Pedersen LM, Liaset B, *et al.* (2008) cAMP-dependent signaling regulates the adipogenic effect of *n*-6 polyunsaturated fatty acids. *J Biol Chem* **283**, 7196–7205.
33. Crowe FL, Skeaff CM, Green TJ, *et al.* (2008) Serum *n*-3 long-chain PUFA differ by sex and age in a population-based survey of New Zealand adolescents and adults. *Br J Nutr* **99**, 168–174.
34. Bakewell L, Burdge GC & Calder PC (2006) Polyunsaturated fatty acid concentrations in young men and women consuming their habitual diets. *Br J Nutr* **96**, 93–99.
35. Giltay EJ, Gooren LJ, Toorians AW, *et al.* (2004) Docosahexaenoic acid concentrations are higher in women than in men because of estrogenic effects. *Am J Clin Nutr* **80**, 1167–1174.
36. Burdge GC & Wootton SA (2002) Conversion of α -linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women. *Br J Nutr* **88**, 411–420.
37. Gerster H (1998) Can adults adequately convert α -linolenic acid (18:3*n*-3) to eicosapentaenoic acid (20:5*n*-3) and docosahexaenoic acid (22:6*n*-3)? *Int J Vitam Nutr Res* **68**, 159–173.
38. Burdge GC & Calder PC (2005) α -Linolenic acid metabolism in adult humans: the effects of gender and age on conversion to longer-chain polyunsaturated fatty acids. *Eur J Lipid Sci Technol* **107**, 426–439.
39. Burdge GC, Slater-Jefferies JL, Grant RA, *et al.* (2008) Sex, but not maternal protein or folic acid intake, determines the fatty acid composition of hepatic phospholipids, but not of triacylglycerol, in adult rats. *Prostaglandins Leukot Essent Fatty Acids* **78**, 73–79.
40. Schaeffer L, Gohlke H, Muller M, *et al.* (2006) Common genetic variants of the FADS1 FADS2 gene cluster and their reconstructed haplotypes are associated with the fatty acid composition in phospholipids. *Hum Mol Genet* **15**, 1745–1756.
41. Malerba G, Schaeffer L, Xumerle L, *et al.* (2008) SNPs of the FADS gene cluster are associated with polyunsaturated fatty acids in a cohort of patients with cardiovascular disease. *Lipids* **43**, 289–299.
42. Giltay EJ, Duschek EJ, Katan MB, *et al.* (2004) Raloxifene and hormone replacement therapy increase arachidonic acid and docosahexaenoic acid levels in postmenopausal women. *J Endocrinol* **182**, 399–408.
43. Caspi A, Williams B, Kim-Cohen J, *et al.* (2007) Moderation of breastfeeding effects on the IQ by genetic variation in fatty acid metabolism. *Proc Natl Acad Sci U S A* **104**, 18860–18865.