

NOTES FOR AUTHORS

Papers submitted for publication in the *Bulletin of Entomological Research* should be sent, in duplicate, to the Director, C·A·B International Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maevé O'Connor & E. Peter Woodford). 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991, Part 1, 1967*, and are alike in singular and plural; stops will be used only where ambiguity might occur, e.g. 'in.' not 'in' for inch, but 'in²' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on Bristol-board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12·7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones, Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

Offprints

Free copies of offprints are allowed as follows: One author—20 copies, two or more authors—30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

BULLETIN OF ENTOMOLOGICAL RESEARCH

(World List Abbreviation: *Bull. ent. Res.*)

ISSN 0007-4853

| | PAGE |
|--|------|
| WRIGHT, A. D., BOLAND, N. P. & KERR, J. D. Changes in leaf hardness and moisture content of water hyacinth, <i>Eichhornia crassipes</i> , following feeding by <i>Neochetina eichhorniae</i> Warner (Coleoptera: Curculionidae) | 529 |
| VAN DER LINDE, T. C. DE K., HEWITT, P. H., VAN DER WESTHUIZEN, M. C. & MITCHELL, J. The use of ¹³¹ I, ¹²⁵ I and aggressive behaviour to determine the foraging area of <i>Hodotermes mossambicus</i> (Hagen) (Isoptera: Hodotermitidae) | 537 |
| DIKE, M. C. A key for the identification of Afrotropical species of the shoot-fly subgenus <i>Atherigona</i> of <i>Atherigona</i> (Diptera: Muscidae), with a description of some new species from Africa | 545 |
| WHEATLEY, A. R. D., WIGHTMAN, J. A., WILLIAMS, J. H. & WHEATLEY, S. J. The influence of drought stress on the distribution of insects on four groundnut genotypes grown near Hyderabad, India | 567 |
| NEUENSCHWANDER, P., HAMMOND, W. N. O., GUTIERREZ, A. P., CUDJOE, A. R., ADJAKLOE, R., BAUMGÄRTNER, J. U. & REGEV, U. Impact assessment of the biological control of the cassava mealybug, <i>Phenacoccus manihoti</i> Matile-Ferrero (Hemiptera: Pseudococcidae), by the introduced parasitoid <i>Epidinocarsis lopezi</i> (De Santis) (Hymenoptera: Encyrtidae) | 579 |
| JACKAI, L. E. N. & OGHIAKHE, S. Pod wall trichomes and resistance of two wild cowpea, <i>Vigna vexillata</i> , accessions to <i>Maruca testulalis</i> (Geyer) (Lepidoptera: Pyralidae) and <i>Clavigralla tomentosicollis</i> Stål (Hemiptera: Coreidae) | 595 |
| MARTIN, J. H. Identification, occurrence and pest status of <i>Toxoptera odinae</i> (van der Goot) (Hemiptera: Aphididae) in Africa | 607 |
| JOHNSTONE, D. R., COOPER, J. F., DOBSON, H. M. & TURNER, C. R. The collection of aerosol droplets by resting tsetse flies, <i>Glossina morsitans</i> Westwood (Diptera: Glossinidae) | 613 |
| SULAIMAN, S., SOHADI, A. R. & JEFFERY, J. Human helminth parasite burdens on cyclorrhaphan flies (Diptera) trapped at an aboriginal settlement in Malaysia | 625 |
| DUNCAN, F. D. & HEWITT, P. H. Observations on the foraging behaviour of the harvester termite, <i>Hodotermes mossambicus</i> (Hagen) (Isoptera: Hodotermitidae) | 631 |
| WILLIAMS, D. J. A new genus and two new species of mealybugs (Hemiptera: Pseudococcidae) from South Africa on bulbs and corms imported into Great Britain | 643 |
| ADAIR, R. F. & SCOTT, J. K. The life-history and host specificity of <i>Comostolopsis germana</i> Prout (Lepidoptera: Geometridae), a biological control agent of <i>Chrysanthemoides monilifera</i> (Compositae) | 649 |
| HEMINGWAY, J., CALLAGHAN, A. & KURTAK, D. C. Temephos resistance in <i>Simulium damnosum</i> Theobald (Diptera: Simuliidae): a comparative study between larvae and adults of the forest and savanna strains of this species complex .. | 659 |
| AGRICOLA, U., AGOUNKÉ, D., FISCHER, H. U. & MOORE, D. The control of <i>Rastrococcus invadens</i> Williams (Hemiptera: Pseudococcidae) in Togo by the introduction of <i>Gyranusoidea tebygi</i> Noyes (Hymenoptera: Encyrtidae) .. | 671 |
| EL-SAYED, S., HEMINGWAY, J. & LANE, R. P. Susceptibility baselines for DDT metabolism and related enzyme systems in the sandfly <i>Phlebotomus papatasi</i> (Scopoli) (Diptera: Psychodidae) | 679 |
| THOMSON, M. C., DAVIES, J. B. & WILSON, M. D. A portable allozyme electrophoresis kit used to identify members of the <i>Simulium damnosum</i> Theobald complex (Diptera: Simuliidae) in the field | 685 |
| HOBBS, S. E. & WOLF, W. W. An airborne radar technique for studying insect migration | 693 |

© C.A.B International, 1989

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.