

# Reconstructing the spatial distribution of the Galactic stellar halo

M. Cignoni<sup>1,2†</sup>, V. Ripepi<sup>1</sup>, M. Marconi<sup>1</sup>, J. M. Alcalá<sup>1</sup>,  
M. Capaccioli<sup>1</sup>, M. Pannella<sup>3</sup> and R. Silvotti<sup>1</sup>

<sup>1</sup>Osservatorio Astronomico Di Capodimonte, Via Moiariello 16, 80131 Napoli, Italy  
email: cignoni@df.unipi.it

<sup>2</sup>Dipartimento di Astronomia, Università di Bologna, Via Ranzani 1, I-40127 Bologna, Italy

<sup>3</sup>Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse, Postfach 1312,  
D-85741 Garching, Germany

**Abstract.** The VLT Survey Telescope (VST) is going to offer a unique chance to collect deep and wide field photometry in multi-directions, opening the door to a definitive mapping of the Galactic halo. In this shortcoming scenario, we present a pilot study aimed at recovering the halo stellar density using the Capodimonte Deep Field (OACDF, Alcalá *et al.* 2004). Turn-off stars are isolated and the relative color-magnitude diagram (CMD) is compared with synthetic CMDs. Our result is consistent with a power law exponent  $n \approx 3$  over a range of Galactocentric distances from 8 to 40 kpc.

**Keywords.** Galaxy: halo, Galaxy: structure, Galaxy: stellar content, Stars: Hertzsprung-Russell diagram, Methods: statistical

---

## 1. Introduction

The observed star counts from the OACDF photometric survey<sup>‡</sup> are investigated. This survey covers about 0.5 square degree in the  $B$ ,  $V$ ,  $R$  optical filters and is located at a high galactic latitude ( $l \sim 293$ ,  $b \sim 50$ ), making available a sample of halo stars with low contamination of disk and thick-disk stars.


To derive the spatial distribution of the Galactic halo, one needs to identify stars in a precise evolutionary phase and use them to map the observed CMD density. Turn-off stars have many advantages:

- They can be univocally selected (definitely bluer than halo red giants);
- They are readily distinguishable from the disk stars in the main sequence (only very young and massive disk stars may have similar colors, but they are rare);
- They are numerous, in fact HB stars and red giants are less frequent than turn-off stars of a factor larger than  $\approx 100$ .

On this basis, we have explored the OACDF color-magnitude diagram selecting turn-off stars by color ( $0.3 \lesssim B - V \lesssim 0.6$ ). Contamination by galaxies is reduced by choosing objects with stellarity index higher than 0.9 (simultaneously in  $B$  and  $V$  filters). The adoption of a  $V$ -magnitude range of 18 to 23 ensures respectively thick disk removal and completeness. Finally, the most probable spatial distribution is obtained by comparing the turn-off density in the observed CMD with that in model CMD.

† Present address: Dipartimento di Astronomia, Via Ranzani 1, I-40127 Bologna, Italy

‡ Acquired with the Wide Field Imager (WFI) at the ESO 2.2-m telescope.



**Figure 1.** The recovered density distribution. For comparison, different power law densities are also shown (with the labeled exponents).

## 2. Method

Following the Monte Carlo framework, an artificial CMD is populated with a large number of stars randomly built by means of a set of theoretical ingredients (see Cignoni *et al.* 2006 for a complete description). This synthetic halo population is modeled as originating from a short period of star formation (constant between 10 and 12 Gyr). Moreover, the metallicity is fixed at  $[Fe/H] \sim -1.6$  with a Gaussian spread ( $\sigma_{[Fe/H]} = 1.0$ ) and the IMF is a power law with a Salpeter index. For each stellar model of a given mass, age and chemical composition, colors and magnitudes are estimated by interpolating a library of stellar tracks (Pisa Evolutionary Library, Cariulo, Degl’Innocenti & Castellani 2004). In order to reconstruct the halo structure, this synthetic population is placed at different distances, randomly placing stars in the  $j$ -th heliocentric interval  $[d_{\odot,j}, d_{\odot,j} + 4 \text{ kpc}]$ †. The final product is a base of partial CMDs, each one representing the same population but at different distance moduli. The code searches (through a simplex algorithm) for the linear combination of these basic CMDs which best matches the observed CMD.

## 3. Results

We have combined a star count model with an updated stellar library to reconstruct the Galactic halo stellar density in the direction of the Capodimonte Deep Field. Our results confirm a steep decline. In particular, a power-law index  $n \approx 3$  (see Fig. 1) gives a reasonable fit out to  $\approx 40$  Kpc from the galactic center.

## Acknowledgements

Financial support for this study was provided by MIUR, under the scientific project “On the evolution of stellar systems: fundamental step toward the scientific exploitation of VST” (P.I. Massimo Capaccioli).

## References

- Alcalá, J. M. *et al.* 2004, *A&A*, 428, 339  
 Cariulo, P., Degl’Innocenti, S., & Castellani 2004, *A&A*, 421, 1121  
 Cignoni, M., Ripepi, V., Marconi, M., Alcalá, J. M., Capaccioli, M., Pannella, M., & Silvotti, R. 2006, *A&A*, DOI: 10.1051/0004-6361:20066598

†  $d_{\odot,j}$  is the heliocentric distance of the  $j$ -th volume bounded by the field of view.