

# What Disc Brightness Profiles Can Tell us about Galaxy Evolution

John Beckman<sup>1,2,3</sup>, Peter Erwin<sup>4</sup> and Leonel Gutiérrez<sup>5</sup>

<sup>1</sup>Instituto de Astrofísica de Canarias, c/ Vía Láctea, s/n, E38205, La Laguna, Tenerife, Spain.  
email: jeb@iac.es

<sup>2</sup>Departamento de Astrofísica. Universidad de La Laguna, Tenerife, Spain.

<sup>3</sup>Consejo Superior de Investigaciones Científicas, Spain.

<sup>4</sup>Max Planck Institut für Extraterrestrische Physik, Germany

<sup>5</sup>UNAM, Ensenada, México.

**Abstract.** Azimuthally averaged surface brightness profiles of disc galaxies provide a most useful practical classification scheme which gives insights into their evolution. Freeman (1970) first classified disc profiles into Type I, with a single exponential decline in surface brightness, and Type II, having a split exponential profile, whose inner radial portion is shallower than its outer section. Van der Kruit & Searle, (1981) drew attention to sharply truncated profiles of outer discs observed edge-on, but more recently Pohlen *et al.* (2004) showed that if these same galaxies were observed face-on their profiles would be of Type II. Finally in Erwin, Beckman and Pohlen (2005) we found a significant fraction of profiles with inner portion steeper than the outer portion, which we termed “antitruncations” or Type III profiles. In Erwin, Pohlen and Beckman (2008), we produced a refined classification, taking into account those Type II’s produced by dynamical effects at the outer Lindblad resonance, and those Type III’s caused by the presence of an outer stellar halo. In Gutiérrez *et al.* (2011) we showed the distribution of the three main profile types along the Hubble sequence. In early type discs Types I and III predominate, while in late types, Sc and later, Type II predominates.

The evolution of Type II’s over cosmic time was studied by Azzollini *et al.* (2008a, 2008b) who obtained four key results: (a) between  $z = 1$  and  $z = 0$  the break radius between the inner (shallower) and outer (steeper) profile has increased systematically, by a factor 1.3; (b) the inner profile has steepened while the outer profile is shallower at lower  $z$ ; (c) the extrapolated central surface brightness has fallen by over two magnitudes; (d) the discs in the full redshift interval are always bluest at the break radius. While this behaviour can be qualitatively explained via evolutionary models including stellar migration plus gas infall, such as that by Roskar *et al.* (2008), and while Type III profiles may have a qualitative explanation via mergers and/or accretion, the widespread existence of Type I’s is still a major conceptual challenge.

---

## References

- Azzollini, R., Trujillo, I., & Beckman, J. E. 2008a, *ApJ. Letters* 679, L69  
Azzollini R., Trujillo, I. & Beckman, J. E. 2008b, *ApJ* 684, 1026  
Erwin, P. E., Beckman, J. E., & Pohlen, M. 2005, *ApJ. Letters* 626, L81  
Erwin, J. E., Pohlen, M., & Beckman, J. E. 2008, *AJ* 135, 20  
Freeman, K. C. 1970, *ApJ* 160, 767  
Van der Kruit, P. & Searle, L. 1981 *A&A* 95, 105  
Gutiérrez, L., Erwin, P., Aladro, R., & Beckman, J. E. 2011, *AJ* 142, 145  
Pohlen, M., Beckman, J. E., *et al.* 2004, in: Block, D., *et al.* (eds.), *ASSL*, Lecture Notes in Physics (Kluwer, Dordrecht), vol. 317, p. 713  
Roskar, R. & Debattista, P. 2008, *ApJ. Letters* 675, L65