

Quantifying associations of the dietary share of ultra-processed foods with overall diet quality in First Nations peoples in the Canadian provinces of British Columbia, Alberta, Manitoba and Ontario

Malek Batal^{1,*}, Louise Johnson-Down², Jean-Claude Moubarac¹, Amy Ing¹, Karen Fediuk³, Tonio Sadik⁴, Constantine Tikhonov⁵, Laurie Chan⁶ and Noreen Willows⁷

¹Département de Nutrition, Faculté de Médecine, Université de Montréal, Pavillon Liliane de Stewart, CP 6128 succ. Centre-Ville, Montréal, QC, Canada, H3T 1A8; ²School of Dietetics and Human Nutrition, McGill University, Ste Anne de Bellevue, QC, Canada; ³First Nations Food, Nutrition and Environment Study, University of Ottawa, Ottawa, ON, Canada; ⁴Assembly of First Nations, Ottawa, ON, Canada; ⁵Environmental Public Health Division, First Nations and Inuit Health Branch, Health Canada, Ottawa, ON, Canada; ⁶Department of Biology, University of Ottawa, Ottawa, ON, Canada; ⁷Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, AB, Canada

Submitted 28 October 2016: Final revision received 8 June 2017: Accepted 13 June 2017: First published online 25 July 2017

Abstract

Objective: To quantify associations of the dietary share of ultra-processed foods (UPF) with the overall diet quality of First Nations peoples.

Design: A cross-sectional analysis of data from the First Nations Food, Nutrition and Environment Study, designed to contribute to knowledge gaps regarding the diet of First Nations peoples living on-reserve, south of the 60th parallel. A multistage sampling of communities was conducted. All foods from 24 h dietary recalls were categorized into NOVA categories and analyses were performed to evaluate the impact of UPF on diet quality.

Setting: Western and Central Canada.

Subjects: First Nations participants aged 19 years or older.

Results: The sample consisted of 3700 participants. UPF contributed 53.9% of energy. Compared with the non-UPF fraction of the diet, the UPF fraction had 3.5 times less vitamin A, 2.4 times less K, 2.2 times less protein, 2.3 times more free sugars and 1.8 times more Na. As the contribution of UPF to energy increased so did the overall intakes of energy, carbohydrate, free sugar, saturated fat, Na, Ca and vitamin C, and Na:K; while protein, fibre, K, Fe and vitamin A decreased. Diets of individuals who ate traditional First Nations food (e.g. wild plants and game animals) on the day of the recall were lower in UPF.

Conclusions: UPF were prevalent in First Nations diets. Efforts to curb UPF consumption and increase intake of traditional First Nations foods and other fresh or minimally processed foods would improve diet quality and health in First Nations peoples.

Keywords
First Nations
Indigenous
Aboriginal
Food intake
Food processing
Traditional food
Ultra-processed foods
Diet quality
Dietary intake
Dietary guidelines
Colonialism

Indigenous peoples throughout the world, including in Canada, suffer a disproportionate share of health problems compared with their non-Indigenous counterparts⁽¹⁾. In Canada in 2011, a total of 859 970 individuals reported being First Nations peoples of Indigenous ancestry, representing 3% of the Canadian population⁽²⁾. Almost half of these individuals lived on reserves and they were younger than the non-Indigenous share of the Canadian population⁽³⁾.

Compared with non-Indigenous peoples, First Nations peoples have a lower life expectancy⁽¹⁾, higher prevalence

of chronic and infectious diseases, and are more prone to have mental health issues^(4–6). Rates of being overweight or obese in First Nations communities are also consistently higher than among other Canadians⁽⁷⁾ and the worldwide prevalence of 39%⁽⁸⁾. First Nations peoples are much more likely than other Canadians to have obesity-related chronic diseases such as type 2 diabetes mellitus and CVD^(9,10). These health disparities result from a complex intersection of factors, many of which have their origin in a long history of colonialism and assaults on indigenous

*Corresponding author: Email malek.batal@umontreal.ca

ways of being and knowing. The alienation of lands and resources and the disempowering of Indigenous peoples from self-determination with respect to the stewardship of their environment have contributed to the socio-economic and political marginalization of these peoples⁽¹¹⁾.

Traditional food (TF) is the collective name for the fresh or minimally processed foods obtained from the local environments of First Nations peoples. First Nations diets in the pre-colonial and early-colonial periods consisted of a wide range of wild foods such as fish, shellfish, game, fowl, and roots, berries and other plants that were gathered from the land. Many First Nations peoples developed mariculture or horticulture to enhance food production, such as the creation of shellfish and root beds, the burning of forest and prairies, and the active collection and storage of seeds for future crops (corn, beans and squash). TF continues to be important to First Nations peoples for multiple reasons⁽¹²⁾. In addition to providing a cost-effective alternative to buying food in a store, TF can contribute to better health and a greater sense of well-being by providing a rich source of nutrients; social cohesion through food preparation and sharing practices; a sense of identity from engaging in customary food procuring traditions; and enhanced physical activity from the exertion required to obtain such foods^(12–18). The introduction of market or store-bought foods (MF) into the diets of First Nations peoples contributed to a nutrition transition that resulted in the adoption of high-energy, nutrient-poor MF and the corresponding erosion of some customary practices associated with TF harvesting and consumption^(12,19). Because MF are quite heterogeneous in terms of nutritional quality (e.g. fruits *v.* carbonated beverages), there is a need to analyse MF with more specificity to better understand their impacts on the quality of the diet of First Nations peoples.

As the study of nutritional sciences evolves, it becomes apparent that one needs to look at more than just nutrients in a diet, and hence better methods of investigating the types of food (e.g. whole *v.* processed) that are consumed have been developed^(20,21). The NOVA classification enables researchers to classify foods according to the nature, extent and purpose of food processing, and has been used to evaluate the impact of food processing on diet quality and health-related outcomes^(22–27).

Studies using the NOVA classification have shown that consumption of ultra-processed foods (UPF) negatively impacts the nutritional quality of diets^(22,24–26,28) and is associated with weight gain, obesity^(24,26,29) and non-communicable diseases (NCD)^(30,31). UPF are industrial food and drink formulations made mostly from refined substances extracted or derived from other food sources and additives. They include sweet or savoury snacks, carbonated soft drinks, ready-to-eat meals, mass-produced packaged breads and buns, poultry and fish nuggets and other reconstituted meat products⁽³²⁾. International bodies such as the Pan American Health Organization and WHO have proposed using the energy share of UPF as an overall indicator of diet quality⁽³³⁾.

An important determinant of obesity and NCD is the proportion of sugar in the diet^(34,35). The WHO introduced the term ‘free sugars’ in 2002 to refer to monosaccharides (such as glucose, fructose) and disaccharides (such as sucrose or table sugar) added to foods and drinks by the manufacturer, cook or consumer, and sugars naturally present in honey, syrups, fruit juices and fruit juice concentrates⁽³⁶⁾. The WHO’s latest recommendation states that free sugars be less than 10% of total energy intake to reduce the risk of overweight, obesity and tooth decay⁽³⁷⁾.

The objective of the present study was to investigate diet quality by estimating the consumption of UPF in the diets of adult First Nations living on reserves in four provinces in Canada and the impact of UPF on their free sugar consumption and overall diet quality. To our knowledge, this type of study has not been done previously.

Methods

Study

Following the passage of a resolution by the Assembly of First Nations in 2007, calling for the generation of more knowledge on the diet and environment of First Nations living on-reserve south of the 60th parallel, the First Nations Food, Nutrition and Environment Study (FNFNES) was developed to address gaps in these areas with respect to First Nations peoples^(15–18). FNFNES is a joint project between the Assembly of First Nations, University of Northern British Columbia, University of Ottawa and Université de Montréal, with funding and technical support from Health Canada, to investigate First Nations’ total diet and food-related exposures to environmental contaminants. This study was guided by the Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans, and the First Nations data-related principles of Ownership, Control, Access and Possession (OCAPTM)⁽³⁸⁾. All research activities were planned conjointly with individual communities through a methodology workshop in each targeted region (using Assembly of First Nations’ regions which largely correspond to the Canadian provinces), and original data, tabulated results and reports were returned to the communities mainly via face-to-face meetings with community members and workers before aggregate reports were provided for each region.

Sampling

A random multistage sampling strategy was established based on communities, households and individuals in four Canadian provinces (British Columbia, Alberta, Manitoba and Ontario)^(15–18). There are 442 First Nation communities in total in these four provinces. Community sampling was conducted based on ecological zones, which are naturally occurring divisions of the earth’s surface based on the distribution and systemic interdependence of plants and animals and their physical environment⁽³⁹⁾.

Households were then randomly selected within communities. In households where more than one individual aged 19 years or older was present, the person with the next birthday was selected to participate in the research.

From 2008 to 2013, 5355 households in fifty-eight communities were sampled using the aforementioned process. Of the fifty-eight communities included in the analyses, eleven (19%) were remote (fly-in or winter road only) and an additional twenty-five (43%) were rural and ≥ 60 km away from the nearest urban centre. Of the 5355 households, fifty were not eligible (not First Nations, 19 years or older, or living on-reserve; or for health reasons such as deafness, cognitive impairment, etc.) and 138 homes were vacant. A total of 3847 inhabitants of the 5167 households completed an interview for a participation rate of 74.5%. These participants were aged 19 years or older, living on-reserve and self-identified as First Nations. Pregnant and breast-feeding women were excluded (n 143) as well as individuals with no food intake the prior day (n 4). The final sample size was 3700 individuals from households in participating First Nations communities in the four provinces.

Data collection

Questionnaires were administered to each household participant by trained community members, with the guidance of a trained dietitian, to collect information on dietary patterns, lifestyle, general health status, environmental concerns and food security. Height and weight measurements were both self-reported and measured for individuals who agreed to have these values recorded. Where measured, participants' weight was recorded using a Seca 803 digital scale (Seca Measuring Systems and Scales, Hanover, MD, USA) with the participant lightly clothed, and height was obtained with participants shoeless on an even surface using a measuring tape. BMI was calculated as weight/height² (kg/m²) for 3362 participants. In 27.4% (n 922) of cases, BMI was calculated from measured height and weight; in 20.6% (n 691) of cases it was calculated from a combination of measured and reported data; and in 52.0% (n 1749) of cases, it was calculated from reported measures. To test for bias in reported anthropometric values, paired t tests by gender were used to test for differences in BMI values using reported and measured heights and weights for participants who provided both these values. To adjust for a possible reporting bias in each province, if the difference between the reported and measured BMI was significant, then the estimated difference between the means (or estimated bias value) was added to the reported BMI where participants provided only reported values. This was necessary for Ontario and Alberta but not for British Columbia and Manitoba. BMI values were missing for 339 individuals who did not consent to provide anthropometric measures. Participants were considered overweight if their BMI was ≥ 25.0 kg/m² and obese if it was ≥ 30.0 kg/m²(40).

The questionnaires included a 24 h diet recall conducted in the autumn. The recall used a three-stage multiple-pass method, i.e. quick list, detailed description and review(41). Portion sizes were estimated using three-dimensional food models (Santé Québec, Montréal, QC, Canada). Because 24 h recalls give reliable estimates of the nutrient intake of groups of individuals, no analyses requiring the evaluation of nutrient cut-offs were done(42). A question on self-reported diabetes was considered in the analysis. This question was not present in the first year of the study in British Columbia and is thus considered in the analysis for only 3314 participants.

Data processing and analysis

All collected data were entered into a database by the research coordinators, except for information derived from the 24 h recalls, which were entered by research nutritionists at the Université de Montréal. Questionnaire data were entered using Epi-Info version 3.4 (Centers for Disease Control and Prevention, Atlanta, GA, USA). The 24 h recalls were analysed using CANDAT (Godin, London, ON, Canada) which is a nutrient analysis software that uses the 2010 Canadian Nutrient File (CNF)(43). In addition to the CNF, a file created by the FNFNES analyst and containing nutritional information on some TF and other MF not available in the CNF was used to complete the CNF and estimate food and nutrient intakes. The accuracy of data entry of the 24 h recalls was ensured using several steps. First, a sub-sample of 10% of the records were cross-checked and discrepancies reconciled. If many errors were found during this first process, a further 10% check was conducted. Preliminary analyses were then performed to review for outliers such as unusual foods and intakes that were ± 2 SD of the mean for energy and selected nutrients. Finally, all data entry errors were reconciled before finalizing the file for data analyses.

Food processing, as described by the NOVA classification, involves the physical, biological and chemical processes used to modify foods in their natural state. All foods and drinks reported in the 24 h recall data were classified into four NOVA groups: (i) fresh or minimally processed foods (fresh, dried or frozen fruits and vegetables, meats, grains and pasteurized milk); (ii) processed culinary ingredients (sugar, oils, fats and salt used in cooking); (iii) processed foods (canned foods, artisanal breads, cheese and smoked or fermented foods); and (iv) UPF (industrial formulations based on refined substances and additives such as flavours, emulsifiers, etc.)(22,44,45). Next, mean estimates and SE for energy intake and proportion of energy (percentage of energy) were calculated from each NOVA food group and selected subgroups within them. TF consumption was investigated (based on whether TF was reported eaten at least once on the 24 h recall) to determine whether it was associated with the consumption of NOVA food groups.

Free sugar content as a percentage of energy was estimated using a publicly available database developed at

the University of Toronto which provided estimates for free sugars for more than 4000 food and drink products sold in Canadian food markets⁽⁴⁶⁾. Each NOVA food group was matched manually with a corresponding item found in the free sugars database to estimate the proportion of free sugars from total sugar (i.e. both free sugars and naturally occurring sugars in fresh fruits, vegetables and milk). Products were matched by brand name when available, or by using a mean value for the various brands of the product. For items not found in the database, free sugars were estimated using the US Department of Agriculture's database on added sugars⁽⁴⁷⁾. Added sugars are sugars and syrups incorporated into foods during preparation or processing, or added at the table, but not naturally occurring sugars in foods⁽⁴⁷⁾. Food items (*n* 320) in the food database with missing values for total sugars were also corrected using the University of Toronto and the US Department of Agriculture databases.

Diet quality was estimated by comparing the nutrient content of fresh or minimally processed foods, processed culinary ingredients and processed foods (collectively referred to as non-UPF) with that of UPF. Thereafter, the population was divided into quintiles of the proportion of energy from UPF. The first quintile thus included the lowest consumers of UPF and the fifth quintile included the highest consumers. The average dietary content of macronutrients (expressed as % of total energy), energy density (kJ/g) and energy-adjusted micronutrients and fibre (expressed as µg, mg or g/4184 kJ (1000 kcal)) was compared across quintiles of dietary share of UPF. Trend and χ^2 analyses were used to identify differences in the dietary contribution between quintiles of UPF and non-UPF.

The 95 % CI were used to evaluate differences in intake between the NOVA categories and trend analyses (using logistic regression) and χ^2 for differences between quintiles of energy from UPF, with significance at $P < 0.05$. Unweighted data analysis used the statistical software packages SAS/STAT version 9.2 (2009) and IBM SPSS Statistics version 22.0 (2013).

Results

Table 1 summarizes some sociodemographic and health characteristics of the participants. Participants were, on average, 45.1 (95 % CI 44.6, 45.6) years old, had a BMI of 30.2 (95 % CI 30.0, 30.4) kg/m² and were 62.7% female. The majority (80%) of participants were overweight, of whom 46% were obese. One-fifth (20.5%) self-reported having diabetes, of which 70.9% of cases were type 2, 10.8% were type 1 and 18.3% were unknown type.

First Nations participants had a mean energy intake of 8166 (95 % CI 8029, 8303) kJ/d (1952 (95 % CI 1920, 1984) kcal/d). UPF accounted for 53.9 (95 % CI 53.1, 54.7)% of total energy, fresh or minimally processed foods for 36.4 (95 % CI 35.7, 37.2)%, processed culinary ingredients for 6.1 (95 % CI 5.89, 6.42)% and processed foods for 3.5 (95 % CI 3.20, 3.70)% of total energy. Table 2 presents the mean energy intake and dietary share of the NOVA food subgroups in the First Nations diet. MF and TF are distinguished in the table. Energy in the diet from UPF came mostly from fast food and ready-to-eat dishes (15.9%), commercial breads (9.3%), carbonated, energy and fruit drinks, and fruit juices (8.0%), deli and processed meats

Table 1 Demographic, anthropometric, health and lifestyle characteristics of adults aged 19 years or older from fifty-eight on-reserve First Nations communities in British Columbia, Alberta, Manitoba and Ontario (Canada), 2008–2013

Characteristic	Men		Women		All	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Completed a 24 h food recall	1379	37.3	2321	62.7	3700	
Traditional food on day of 24 h recall	341	24.7	499	21.5	840	22.7
Age (years), mean and 95 % CI	45.9	45.1, 46.8	44.6	44.0, 45.2	45.1	44.6, 45.6
19–30 years	255	18.6	434	18.8	689	18.7
31–50 years	602	43.9	1112	48.2	1718	46.4
51–70 years	432	31.5	626	27.1	1058	28.6
≥71 years	84	6.12	135	5.81	219	5.92
Anthropometry recorded†, <i>n</i>	1315		2047		3362	
BMI (kg/m ²), mean and 95 % CI	29.6	29.3, 29.9	30.6	30.3, 30.9	30.2	30.0, 30.4
BMI ≥ 25.0 kg/m ²	506	38.5	626	30.6	1132	33.7
BMI ≥ 30.0 kg/m ²	545	41.5	1001	48.9	1546	46.0
Education‡, <i>n</i>	1227		2075		3302	
Years of schooling, mean and 95 % CI	10.3	10.1, 10.4	10.9	10.8, 11.0	10.7	10.6, 10.8
Less than high school	609	49.7	895	43.1	1504	45.5
High school or equivalent	522	42.5	855	41.2	1377	41.7
Post-secondary degree	95	7.70	325	15.7	420	12.7
Reported diabetes status§, <i>n</i>	1233		2080		3314	
Diabetes, self-reported§	248	20.1	431	20.7	679	20.5

First Nations Food, Nutrition and Environment Study (FNFNES). Data are unweighted, and presented as *n* and % unless indicated otherwise.

†Number of individuals with self-reported or measured anthropometry was less than those who were interviewed for a 24 h recall.

‡Number of individuals who reported their education on the questionnaire was less than those who were interviewed for a 24 h recall.

§Self-reported prevalence based on the question: 'Have you been diagnosed with diabetes?' Number of individuals who self-reported having been diagnosed with diabetes or not on the questionnaire was less than those who were interviewed for a 24 h recall.

Table 2 Mean energy contribution of NOVA food groups and subgroups to the diet of 3700 adults aged 19 years or older from fifty-eight on-reserve First Nations communities in British Columbia, Alberta, Manitoba and Ontario (Canada), 2008–2013

Food group/subgroup	Absolute energy intake (kJ/d)		Relative energy intake (% of total)	
	Mean	95% CI	Mean	95% CI
Fresh or minimally processed foods	2900	2822, 2979	36.4	35.7, 37.2
Meat and poultry – market	839	795, 883	10.0	9.55, 10.5
Meat – traditional	209	184, 233	2.64	2.36, 2.91
Grain products and flours	484	454, 514	6.20	5.84, 6.56
Eggs	254	238, 269	3.41	3.20, 3.62
Roots and tubers	214	199, 229	2.75	2.56, 2.94
Pasta	215	194, 235	2.62	2.37, 2.86
Home-made dishes†	222	197, 246	2.61	2.34, 2.87
Milk and plain yoghurt	145	134, 156	1.88	1.75, 2.02
Fruits – market	117	108, 126	1.66	1.53, 1.80
Fruits – traditional	1.27	0.54, 1.99	0.02	0.00, 0.03
Fish – market	27.3	19.5, 35.0	0.35	0.26, 0.44
Fish – traditional	53.1	41.3, 64.9	0.60	0.48, 0.73
Vegetables – market	61.8	56.9, 66.8	0.87	0.80, 0.95
Vegetables – traditional	0.28	0.00, 0.57	0.00	0.00, 0.01
Other fresh or minimally processed foods‡	57.9	47.9, 37.8	0.80	0.67, 0.94
Other fresh or minimally processed – traditional food§	0.14	0.00, 0.31	0.00	0.00, 0.01
Processed culinary ingredients	492	468, 516	6.15	5.89, 6.42
Sugars	230	215, 245	2.96	2.79, 3.14
Animal fats – market	147	136, 159	1.84	1.69, 1.98
Animal fats – traditional	5.52	0.48, 10.6	0.05	0.01, 0.09
Plant oils	109	97.2, 120	1.30	1.18, 1.42
Other ingredients¶	0.37	0.15, 0.60	0.00	0.00, 0.01
Processed foods	289	265, 312	3.45	3.20, 3.70
Canned meat and fish – market	75.2	65.0, 85.3	1.01	0.88, 1.14
Canned meat and fish – traditional	7.05	3.45, 10.60	0.10	0.05, 0.15
Cheese	74.8	63.0, 86.6	0.84	0.72, 0.95
Canned fruits, vegetables and legumes	49.5	43.6, 55.4	0.62	0.55, 0.69
Other processed foods††	82.1	67.3, 96.8	0.88	0.74, 1.03
Ultra-processed foods	4485	4376, 4593	53.9	53.1, 54.7
Fast food and ready-to-eat dishes‡‡	1340	1278, 1402	15.9	15.3, 16.5
Mass-produced packaged breads	685	659, 709	9.27	8.95, 9.60
Carbonated, sports and energy drinks	321	298, 343	3.82	3.57, 4.07
Fruit juices and fruit drinks	358	330, 386	4.17	3.88, 4.45
Reconstituted meat products§§	387	361, 413	4.61	4.33, 4.89
Chips, crackers and other salty snacks	342	311, 372	3.61	3.33, 3.89
Chocolate, candies and other sweets	255	232, 278	2.87	2.64, 3.10
Cookies, cakes and baked goods	205	184, 227	2.29	2.07, 2.51
Margarine	176	160, 193	2.08	1.92, 2.23
Breakfast cereals	149	135, 163	1.98	1.79, 2.17
Sauces and spreads	138	122, 153	1.60	1.47, 1.73
Other ultra-processed foods	129	119, 139	1.74	1.61, 1.87
Total	8166	8029, 8303	100.0	–

†Soups, salads, sauces, baked goods, stews and other dishes made from fresh or minimally processed foods.

‡Spices and herbs, pulses, unsalted and unsweetened nuts; coffee, tea.

§Labrador tea, wild ginger tea and peppermint.

||White and brown sugar, icing sugar, molasses, honey and maple syrup.

¶Vinegar, leavening agents, unsweetened cocoa powder, corn starch.

††Salted, sweetened or oil-roasted nuts or seeds, plain tortillas and potato chips, tofu, soya sauce, condensed milk, peanut butter, French and pita breads, bannock and dumplings.

‡‡Hot dogs, hamburgers, pizzas and sandwiches from fast-food outlets, commercial pies, canned or dehydrated soups; frozen and prepared French fries and onion rings and frozen meals.

§§Sausages, luncheon meats, meat spreads, bacon, corned beef, beef jerky and fish sticks.

|||Cheese products, fish or seafood imitations, meal replacements, sweeteners, supplements and coffee whitener.

(4.6%), and salty snacks (3.6%). Most of the energy in the diet from fresh or minimally processed foods came from meat and poultry (12.6% of energy), grains and flours (6.2%), eggs (3.4%), and roots and tubers (2.8%). Energy from market meats was four times that from traditional game and birds (839 kJ/d (201 kcal/d) *v.* 209 kJ/d (50 kcal/d)), whereas traditional fish provided more energy than market

fish (53.1 kJ/d (12.7 kcal/d) *v.* 27.3 kJ/d (6.52 kcal/d)). Processed culinary ingredients that contributed the most to total energy were sugars (3.0% of energy) and animal fats (1.9%), mostly from market sources (6.1%).

Table 3 compares the energy and nutrient content of NOVA food groupings in the First Nations diet. The diet fraction from UPF was more energy-dense compared with

Table 3 Mean energy and nutrients from non-ultra-processed foods (non-UPF)[†] and ultra-processed foods (UPF)[†] of 3700 adults aged 19 years or older from fifty-eight on-reserve First Nations communities in British Columbia, Alberta, Manitoba and Ontario (Canada), 2008–2013

Nutrient	Fresh or minimally processed foods		Processed culinary ingredients		Processed foods		Ultra-processed foods	
	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
Energy (kJ)	2900	2822, 2979	492	468, 516	289	265, 312	4485	4376, 4593
Energy density (kJ/g) [†]	6.18	6.08, 6.28	20.3	20.0, 20.7	7.90	7.59, 8.22	12.2	12.0, 12.4
Protein (% energy)	29.2	28.7, 29.7	0.67	0.61, 0.73	10.2	9.65, 10.7	11.1	10.9, 11.3
Carbohydrates (% energy)	37.8	37.0, 38.5	39.4	38.0, 40.8	13.9	13.0, 14.8	54.5	54.0, 55.1
Free sugars (% energy)	0.74	0.61, 0.86	37.6	36.2, 38.9	1.83	1.58, 2.09	20.7	20.2, 21.3
Total fats (% energy)	30.9	30.4, 31.5	32.5	31.2, 33.8	14.2	13.4, 15.0	32.7	32.2, 33.2
Saturated fats (% of energy)	9.42	9.22, 9.61	12.8	12.0, 13.3	9.20	5.78, 6.62	10.2	9.98, 10.4
Fibre (g/4184 kJ)	8.08	7.73, 8.42	0.05	0.00, 0.13	6.35	5.82, 6.87	7.52	7.34, 7.71
Na (mg/4184 kJ)	961	859, 1062	2225	1552, 2898	3972	3428, 4515	2279	2203, 2355
K (mg/4184 kJ)	3933	3102, 4765	85.1	76.7, 93.6	799	739, 859	917	896, 939
Na:K	0.51	0.47, 0.56	43.6	31.3, 55.9	1.89	74, 2.04	3.79	3.06, 4.52
Ca (mg/4184 kJ)	607	506, 708	66.8	59.1, 74.5	269	250, 288	304	295, 313
Fe (mg/4184 kJ)	9.29	8.95, 9.63	0.17	0.13, 0.20	4.27	3.95, 4.59	7.43	7.27, 7.59
Vitamin A (µg/4184 kJ)	548	497, 600	165	155, 175	146	120, 172	132	125, 139
Vitamin C (mg/4184 kJ)	54.7	49.6, 59.9	0.39	0.33, 0.45	32.7	23.4, 42.1	47.3	44.0, 50.7

First Nations Food, Nutrition and Environment Study (FNFNES). Data are unweighted.

[†]Energy density calculated only for the solid fraction of the diet, referring to the sum of energy provided by solid foods divided by the amount of these foods in grams.

the non-UPF fraction (i.e. the sum of minimally processed foods, culinary ingredients and processed foods). Compared with the non-UPF fraction of the diet, the UPF fraction had 3.5 times less vitamin A, 2.4 times less K, 2.2 times less protein, 2.3 times more free sugars and 1.8 times more Na. Saturated fats were equally high in both fractions at over 10% of energy.

Table 4 presents how the mean nutrient content of the diet varied across quintiles of energy from UPF. It also reports the total energy and nutrients across categories. As the dietary contribution of UPF to energy increased, the contribution of energy density, carbohydrate, free sugar, fibre, fat, saturated fat, vitamin C, Na and Ca increased, as did Na:K, while protein, K, Fe and vitamin A intakes decreased. Diets of individuals who ate TF on the day of the recall were higher in fresh and minimally processed foods, processed culinary ingredients and processed foods, and lower in UPF, compared with the diets of individuals who did not eat TF. The percentage of those who ate TF decreased from 42.3% in the first quintile of energy from UPF to 5.5% in the highest quintile of UPF intake.

Discussion

Organizations such as the FAO have recognized the need to incorporate the collection of information on processed foods into food consumption surveys⁽⁴⁵⁾. UPF sales are increasing over time, especially rapidly in low- and middle-income countries and populations around the world⁽⁴¹⁾, and may be related to the rising prevalence of obesity^(23,26,48) and related NCD^(21,44,49,50).

Given the legacy of colonialism in Canada, many Indigenous peoples remain caught in a vicious cycle of

poverty and food insecurity, violence, educational failure and ill health^(51,52). For First Nations peoples, TF systems have vital nutritional, cultural, symbolic and spiritual value. Obstacles to attaining TF and fresh and minimally processed MF are the high cost of healthy, nutritious foods relative to income, along with numerous barriers to the harvesting, sharing and consumption of TF⁽¹²⁾. A consequence of the involuntary transition from TF to market-based diets is increased consumption of UPF and a greater burden of obesity and obesity-related diseases⁽¹⁾. The prevalence of obese (46%) and overweight individuals (34%) was very high among First Nations participants in the present study: there is some concern that cut-offs used for determining obesity are different for First Nations peoples⁽⁵³⁾, but others have confirmed that they are in fact the same as for those of European ancestry⁽⁵⁴⁾. The proportion of participants who reported having diabetes in our study (21.5%) is 2.2 times higher than the national average in Canada of 9.3%⁽⁵⁵⁾. In this sample of First Nations peoples, we found high intakes of UPF (54% of energy), while other studies have consistently shown the benefits of TF^(13,14,56). We have also shown that the consumption of TF on the day of the recall was associated with lower energy from UPF.

One striking finding was the diminished contribution of protein to energy as the contribution of UPF to energy increased. Percentages of energy as protein, carbohydrate and fat were within the acceptable macronutrient distribution ranges⁽⁵⁷⁾. As with other studies, we found that free sugar increased with higher energy from UPF^(24,58). These sugars have been associated with weight gain and obesity⁽³⁴⁾. The contribution of free sugar to energy in the highest quintile of UPF intake was double the WHO recommendation of less than 10% of energy from free sugars. Saturated fat

Table 4 Mean nutrients across quintiles of the dietary contribution of ultra-processed foods (UPF) to energy of 3700 adults aged 19 years or older from fifty-eight on-reserve First Nations communities in British Columbia, Alberta, Manitoba and Ontario (Canada), 2008–2013

Indicators	Quintiles of the contribution of UPF to energy										Total energy and nutrient intakes	
	Q1 (n 740)		Q2 (n 740)		Q3 (n 740)		Q4 (n 740)		Q5 (n 740)		Mean	95 % CI
	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI	Mean	95 % CI		
Energy (kJ)†	7538	7254, 7821	8102	7812, 8391	8230	7943, 8516	8657	8357, 8986	8387*	8048, 8726	8166	8029, 8303
Energy density(kJ/g)†	7.34	7.16, 7.56	7.98	7.81, 8.16	8.45	8.25, 8.66	8.93	8.75, 9.10	10.4*	10.2, 10.9	8.65	8.54, 8.76
Protein (% energy)†	24.7	23.9, 25.5	19.9	19.4, 20.5	17.4	17.0, 17.8	15.2	14.8, 15.5	12.8*	12.5, 13.1	18.0	17.7, 18.3
Carbohydrates (% energy)†	41.1	40.0, 42.1	46.0	45.1, 46.8	48.2	47.4, 49.1	50.5	49.7, 51.4	52.6*	51.7, 53.5	47.7	47.2, 48.1
Free sugars (% energy)†	8.32	7.71, 8.94	13.0	12.2, 13.7	15.3	14.5, 16.0	17.4	16.5, 18.3	19.8*	18.8, 20.8	14.8	14.4, 15.1
Fat (% energy)†	34.2	33.3, 35.1	34.1	33.3, 34.9	34.4	33.6, 35.1	34.3	33.5, 35.0	34.6	33.8, 34.4	34.3	34.0, 34.7
Saturated fat (% energy)†	10.6	10.2, 11.0	10.7	10.4, 11.0	10.9	10.6, 11.2	11.1	10.8, 11.4	11.2*	10.9, 11.5	10.9	10.8, 11.1
Fibre (g/4184 kJ)†	7.36	7.00, 7.71	7.48	7.17, 7.79	7.14	6.86, 7.42	7.13	6.86, 7.41	6.62*	6.37, 6.86	7.14	7.01, 7.28
Na (mg/4184 kJ)†	1308	1246, 1369	1509	1453, 1565	1693	1638, 1749	1815	1751, 1878	2034*	1949, 2118	1672	1641, 1702
K (mg/4184 kJ)†	1669	1546, 1793	1467	1421, 1513	1301	1268, 1335	1237	1203, 1271	1101*	1066, 1137	1355	1326, 1385
Na:K ratio†	0.95	0.90, 1.00	1.18	1.12, 1.24	1.46	1.40, 1.52	1.67	1.59, 1.74	2.23*	1.94, 2.51	1.49	1.43, 1.56
Ca (mg/4184 kJ)†	298	282, 313	319	305, 333	332	319, 346	333	319, 347	342*	328, 355	325	318, 331
Fe (mg/4184 kJ)†	8.93	8.56, 9.30	8.11	7.82, 8.41	7.54	7.28, 7.79	7.25	7.00, 7.50	6.80*	6.57, 7.02	7.73	7.60, 7.85
Vitamin A (µg/4184 kJ)†	305	261, 349	314	263, 366	286	244, 329	242	208, 276	184*	164, 204	266	249, 284
Vitamin C (mg/4184 kJ)†	35.0	31.3, 38.6	38.3	34.8, 41.8	43.5	39.1, 47.9	49.5	43.2, 55.8	46.9*	41.3, 52.5	42.6	40.5, 44.8
Traditional food eater (%),‡,§		42.3		29.6		20.3		15.8		5.54**		22.7

First Nations Food, Nutrition and Environment Study (FNFNES). Data are unweighted.

* $P < 0.05$, ** $P < 0.01$.

†Trend analyses across quintiles.

‡Differences using χ^2 test.

§Based on whether First Nations traditional food was consumed at least once on the 24 h recall.

intake in our study was greater than 10% of energy in all quintiles of UPF intake in First Nations individuals; higher saturated fat intakes potentially increase CVD⁽⁵⁹⁾.

UPF in our study of dietary intake from 2008 to 2013 represented 54% of energy compared with other studies of intake of non-indigenous populations in Brazil from 2008–2009 (21.5%)⁽²⁴⁾, the USA from 2009–2010 (57.9%)⁽⁵⁸⁾ and the UK from 2008–2012 (53%)⁽²²⁾. Because Brazil is classified as middle-income country and UPF are relatively more expensive, a lower intake of UPF would be expected⁽²¹⁾, while our results are consistent with other upper-income countries such as the USA and the UK despite the fact that many First Nations in Canada live in low-income households⁽⁶⁰⁾. Studies based on food expenditure data, which consider household purchases only without accounting for food waste, show 54% of UPF in Canada (2011)⁽⁴⁴⁾, 28% in Brazil (2002–2003)⁽²⁵⁾ and 63% in the UK (2008–2012)⁽²²⁾.

Our results agree with population-based studies in Canada⁽⁴⁴⁾, the USA⁽⁵⁸⁾, the UK⁽²²⁾ and Brazil⁽²⁴⁾ that have consistently shown that UPF are nutritionally inferior compared with non-UPF and are linked to an overall deterioration of diet quality. These foods were found to have a higher content of free sugars, saturated fats and Na, and were more energy-dense compared with non-UPF, whereas UPF were also lower in protein, minerals, most vitamins and fibre. The nutritional quality of the diet fraction of UPF consumed by First Nations peoples in our study was mostly consistent with previous studies, with the exception of higher fibre and vitamin C as UPF increased. The higher content of vitamin C in UPF in Canada can be attributed to fortification practices.

Since the NOVA food categories are a relatively recent development, few studies have been able to evaluate associations of intake with mortality and morbidity⁽²⁵⁾. Studies in Brazil indicate the existence of associations between UPF and metabolic syndrome in adolescents⁽³¹⁾, dyslipidaemias in children⁽³⁰⁾ and obesity in all age groups^(23,26). A cross-sectional time-series analysis performed on thirteen Latin American countries revealed an association between increased sales of UPF and increased BMI in adults from 2000 to 2009⁽⁴⁸⁾. A recent prospective cohort study in Spain found that consumption of UPF was associated with higher risk of overweight and obesity⁽²⁹⁾. In future studies, we intend to examine the association of UPF intake with NCD and BMI of First Nations peoples who participated in the FNFNES.

As was evident from our data, decreasing K and increasing Na intakes with higher UPF may lead to increased risk of hypertension in an already susceptible population⁽⁶¹⁾. Reduced Na intakes have long been recommended as a public health initiative to reduce blood pressure and CVD^(62,63). Concurrently, higher K has a protective effect on blood pressure and related NCD⁽⁶¹⁾. Furthermore, the increasing Na:K is a concern as this has also been associated with blood pressure and CVD^(64,65).

TF provides a very important food source for Indigenous populations in Canada⁽¹²⁾ and some of the TF most

common in such diets are clearly beneficial to health⁽⁶⁶⁾. First Nations peoples in Canada have been going through a nutrition transition where MF are increasingly replacing TF in their diets⁽¹²⁾. Increased health challenges in some First Nations communities, such as high rates of obesity, diabetes and CVD⁽¹⁾, make it imperative to identify ways to slow down these epidemics. There is plenty of evidence that diets including TF are more nutrient-dense^(67,68); therefore, it can be speculated that diets higher in TF may result in a lower prevalence of obesity and diet-related NCD⁽¹⁾. An inverse association between TF intake and energy from UPF in our results is an interesting and novel finding that shows the positive benefit of TF consumption for First Nations people. Given the many co-benefits of TF for Indigenous populations (e.g. social cohesion, sense of identity, physical exertion), there is simply no good reason to avoid stressing the importance of TF to decision makers so that access to and concomitant conservation of TF become increasingly possible.

Public health recommendations are made targeting nutrient intakes that may not be easily understood and applied by the general population. Food-based advice is generally easier to follow. A recommendation to decrease UPF intake in First Nations participants could significantly increase protein and K intakes and decrease free sugar, Na and overall energy intakes. A reduction of UPF would also lead to a healthier diet as these foods would potentially be replaced by foods with a better nutritive profile. Doing this would require policy changes that reduce the availability of UPF and allow all Canadians, regardless of income or geography, to choose a diet rich in fresh and minimally processed foods, and additionally address both climate change and pollution of natural resources. Support is also needed for individuals to make these changes through educational initiatives to promote the optimal use of MF and TF, when available.

One of the limitations of our study was missing data, mainly due to a refusal to answer questions about years of education (12.2% missing data), highest degree obtained (10.8%), age (7.8%) and BMI (9.2%). However, the proportion of missing data was greatest with respect to education and diabetes (10.5% missing data) because these questions were not included during the first year of data collection (i.e. in British Columbia).

However, the large sample allowed us to identify trends that may not have been evident with a smaller population. While the study may not be representative of all First Nations in Canada, as data from only four out of the ten provinces were represented, and there may be systematic differences in dietary intake across other regions in Canada, further data from across the country will be available when FNFNES is completed in 2018.

A potential limitation of our study is that, due to social desirability bias (i.e. valuing fresh foods more than processed foods), the intake of minimally processed foods and TF may have been over-reported by participants

whereas the intake of UPF may have been under-reported⁽⁶⁹⁾; consequently energy intake may have been underestimated. The NOVA food categories were created from the self-reported dietary data, so social desirability bias may have been minimized compared with using an FFQ containing the different NOVA food groups. The standardized methods that we used may help better estimate intake⁽⁴¹⁾. This method may also lead to the overestimation of healthy foods and the underestimation of unhealthy foods that may have influenced our results. The FAO considers 24 h recalls as one of the best ways of collecting data for this type of analysis, as more information is gathered on the types of foods using an open-ended approach⁽⁴⁵⁾. However, with one 24 h recall, only the intake of the group can be considered, not the proportion of the population meeting nutrition recommendations. National nutrient databases evolve over time and there is always some possibility of error and missing information on nutrients.

As the NOVA food classification was applied *a posteriori* from the entry and coding of the recalls, it is possible that some foods were misclassified due to limited food choices in the CNF and a lack of brand names in the database. If data entry had been designed to correspond with the NOVA classification, more accurate distribution may have been obtained and other studies have noted this possible limitation^(22,45,58). Since the NOVA food classification does not group foods according to their nutritional quality, but rather based on the degree and nature of processing, some groupings might appear unusual from a nutritional perspective. However, the current study and others have shown the linear relationship between the increased contribution to energy from UPF and reduced diet quality, regardless of the inclusion/exclusion of specific food items in the UPF group and their individual nutrient profile. In addition, CANDAT, our nutrient analysis software, does not allow us to disaggregate mixed foods based on recipes, so it is possible that some UPF used in culinary preparations, such as margarine used in making a traditional First Nations' quick bread called bannock, may have been included in the fresh and minimally processed foods.

Future studies could benefit from standard guidelines for assignment to NOVA categories, such as brand names, home prepared *v.* ready-to-consume, and name of restaurant where food is purchased⁽²²⁾. Future analyses might identify the MF that, when consumed in combination with TF, will contribute to a diet for First Nations peoples that is more congruent with the current dietary recommendations and more protective against the development of NCD. This, in turn, will allow public health initiatives to better target dietary advice to this particular population, thereby supporting not only better health and nutrition, but also a wide range of other factors that can contribute to the overall well-being and cultural resilience of Indigenous peoples in Canada.

Acknowledgements

Acknowledgements: The authors would like to acknowledge the efforts of the following individuals in the collection of the data used in this study: Olivier Receveur, Harold Schwartz, William David, Laverne Gervais, Lisa Wabegijig, Judy Mitchell and Kathleen Lindhorst. They would like to thank all members of the communities who participated in and worked on the study. *Financial support:* Funding for this analysis was provided by an operating grant from the Canadian Institutes for Health Research (CIHR) for the research project 'Pulling Together for Health Research: Food Security in First Nations Communities' (grant numbers 348833 and 334049). The data used in this article originate from the First Nations Food, Nutrition and Environment Study (FNFNES) funded by Health Canada. At the time of this research, N.W. was the recipient of a Health Scholar award from Alberta Innovates Health Solutions. The funders had no role in the design, analysis or writing of this article. *Conflict of interest:* No conflict of interest. *Authorship:* M.B. and N.W. designed the concept, guided the analyses and planned the manuscript. L.C., M.B. and T.S. are Principal Investigators and C.T. is Co-Investigator of FNFNES which provided the data for the current analyses. K.F. helped in the study design and coordinated data collection of FNFNES. M.B. and N.W. wrote the manuscript with the help of L.J.D. T.S., C.T. and L.C. edited the manuscript for content and clarity. J.-C.M. contributed to the literature review, categorized the foods into NOVA groups and designed the analyses. J.-C.M., L.J.D. and A.I. analysed the data. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and was guided by the principles of the Canadian Institutes of Health Research Guidelines for Health Research Involving Aboriginal People, and the Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans. All procedures involving human participants were approved by the Research Ethics Board of Health Canada, University of Northern British Columbia, University of Ottawa and Université de Montréal. Written informed consent was obtained from all participants. The present study was approved by ethics boards at the Université de Montréal and University of Alberta.

References

1. Anderson I, Robson B, Connolly M *et al.* (2016) Indigenous and tribal peoples' health (The Lancet-Lowitja Institute Global Collaboration): a population study. *Lancet* **388**, 131–157.
2. Statistics Canada (2015) Aboriginal Peoples: Fact Sheet for Canada. <http://www.statcan.gc.ca/pub/89-656-x/89-656-x2015001-eng.htm> (accessed September 2016).
3. Statistics Canada (2016) Aboriginal Peoples in Canada: First Nations People, Métis and Inuit. <https://www12.statcan.gc.ca/nhs-enm/2011/as-sa/99-011-x/99-011-x2011001-eng.cfm#a3> (accessed September 2016).

4. Public Health Agency of Canada (2011) *Diabetes in Canada: Facts and Figures from a Public Health Perspective*. Ottawa, ON: Public Health Agency of Canada.
5. Public Health Agency of Canada (2012) *Tuberculosis in Canada 2008*. Ottawa, ON: Minister of Public Works and Government Services Canada.
6. Public Health Agency of Canada (2011) *Obesity in Canada*. Ottawa, ON: Public Health Agency of Canada and Canadian Institute for Health Information.
7. Statistics Canada (2009) CANSIM table 105-0507. In *Measured Adult Body Mass Index (BMI) by Age Group and Sex, Household Population Age 18 and Over Excluding Pregnant Females, Canada (Excluding Territories)*. Ottawa, ON: Statistics Canada.
8. World Health Organization (2016) Obesity and overweight: Fact sheet. <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed September 2016).
9. Ayach BB & Korda H (2010) Type 2 diabetes epidemic in First Nations people of Canada. *Ethn Dis* **20**, 300–303.
10. Belanger-Ducharme F & Tremblay A (2005) Prevalence of obesity in Canada. *Obes Rev* **6**, 183–186.
11. Adelson N (2005) The embodiment of inequity: health disparities in aboriginal Canada. *Can J Public Health* **96**, Suppl. 2, S45–S61.
12. Willows ND (2005) Determinants of healthy eating in Aboriginal peoples in Canada: the current state of knowledge and research gaps. *Can J Public Health* **96**, Suppl. 3, S32–S36 (S36–S41).
13. Batal M, Gray-Donald K, Kuhnlein HV *et al.* (2005) Estimation of traditional food intake in indigenous communities in Denendeh and the Yukon. *Int J Circumpolar Health* **64**, 46–54.
14. Johnson-Down LM & Egeland GM (2013) How is nutrition transition affecting dietary adequacy in Eeyouch (Cree) adults of Northern Quebec, Canada? *Appl Physiol Nutr Metab* **38**, 300–305.
15. Chan L, Receveur O, Batal M *et al.* (2014) *First Nations Food, Nutrition and Environment Study (FNFNES): Results from Ontario (2011/2012)*. Ottawa, ON: University of Ottawa.
16. Chan L, Receveur O, Batal M *et al.* (2012) *First Nations Food, Nutrition and Environment Study (FNFNES): Results from Manitoba (2010)*. Prince George, BC: University of Northern British Columbia.
17. Chan L, Receveur O, Batal M *et al.* (2016) *First Nations Food, Nutrition and Environment Study (FNFNES): Results from Alberta 2013*. Ottawa, ON: University of Ottawa.
18. Chan L, Receveur O, Batal M *et al.* (2011) *First Nations Food, Nutrition and Environment Study (FNFNES): Results from British Columbia (2008/2009)*. Prince George, BC: University of Northern British Columbia.
19. Egeland G & Harrison GG (2013) Health disparities: promoting Indigenous Peoples' health through traditional food systems and self-determination. In *Indigenous Peoples' Food Systems & Well-Being: Interventions & Policies for Healthy Communities* [HV Kuhnlein, B Erasmus, D Spigelski *et al.*, editors]. Rome: FAO and Centre for Indigenous Peoples' Nutrition and Environment.
20. Fardet A, Rock E, Bassama J *et al.* (2015) Current food classifications in epidemiological studies do not enable solid nutritional recommendations for preventing diet-related chronic diseases: the impact of food processing. *Adv Nutr* **6**, 629–638.
21. Monteiro CA, Moubarac JC, Cannon G *et al.* (2013) Ultra-processed products are becoming dominant in the global food system. *Obes Rev* **14**, Suppl. 2, 21–28.
22. Adams J & White M (2015) Characterisation of UK diets according to degree of food processing and associations with socio-demographics and obesity: cross-sectional analysis of UK National Diet and Nutrition Survey (2008–12). *Int J Behav Nutr Phys Act* **12**, 160.
23. Louzada ML, Baraldi LG, Steele EM *et al.* (2015) Consumption of ultra-processed foods and obesity in Brazilian adolescents and adults. *Prev Med* **81**, 9–15.
24. Louzada ML, Martins AP, Canella DS *et al.* (2015) Ultra-processed foods and the nutritional dietary profile in Brazil. *Rev Saude Publica* **49**, 38.
25. Monteiro CA, Levy RB, Claro RM *et al.* (2011) Increasing consumption of ultra-processed foods and likely impact on human health: evidence from Brazil. *Public Health Nutr* **14**, 5–13.
26. Canella DS, Levy RB, Martins AP *et al.* (2014) Ultra-processed food products and obesity in Brazilian households (2008–2009). *PLoS One* **9**, e92752.
27. Moubarac JC, Batal M, Louzada ML *et al.* (2017) Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* **108**, 512–520.
28. Moubarac JC, Martins AP, Claro RM *et al.* (2013) Consumption of ultra-processed foods and likely impact on human health. Evidence from Canada. *Public Health Nutr* **16**, 2240–2248.
29. Mendonca RD, Pimenta AM, Gea A *et al.* (2016) Ultraprocessed food consumption and risk of overweight and obesity: the University of Navarra Follow-Up (SUN) cohort study. *Am J Clin Nutr* **104**, 1433–1440.
30. Rauber F, Campagnolo PD, Hoffman DJ *et al.* (2015) Consumption of ultra-processed food products and its effects on children's lipid profiles: a longitudinal study. *Nutr Metab Cardiovasc Dis* **25**, 116–122.
31. Tavares LF, Fonseca SC, Garcia Rosa ML *et al.* (2012) Relationship between ultra-processed foods and metabolic syndrome in adolescents from a Brazilian Family Doctor Program. *Public Health Nutr* **15**, 82–87.
32. Moodie R, Stuckler D, Monteiro C *et al.* (2013) Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *Lancet* **381**, 670–679.
33. Vandevijvere S, Monteiro C, Krebs-Smith SM *et al.* (2013) Monitoring and benchmarking population diet quality globally: a step-wise approach. *Obes Rev* **14**, Suppl. 1, 135–149.
34. Te Morenga L, Mallard S & Mann J (2013) Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* **346**, e7492.
35. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Report of a Joint WHO/FAO Expert Consultation*. WHO Technical Report Series no. 916. Geneva: WHO.
36. World Health Organization (2002) *The World Health Report 2002: Reducing Risks, Promoting Healthy Life*. Geneva: WHO.
37. World Health Organization (2015) *Guideline: Sugars Intake for Adults and Children*. Geneva: WHO.
38. Schnarch B (2004) Ownership, control, access, and possession (OCAP) or self-determination applied to research: a critical analysis of contemporary First Nations research and some options for First Nations communities. *J Aboriginal Health* **1**, 80–95.
39. Wiken EB (1986) Terrestrial Ecozones of Canada. Ecological Land Classification Series No. 19. <http://ecozones.ca/english/introduction.html> (accessed April 2017).
40. World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation*. WHO Technical Report Series no. 894. Geneva: WHO.
41. Raper N, Perloff B, Ingwersen L *et al.* (2004) An overview of USDA's dietary intake data system. *J Food Compos Anal* **17**, 545–555.

42. Institute of Medicine, Food and Nutrition Board (2000) *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academies Press.
43. Health Canada (2010) Canadian Nutrient File. <https://www.canada.ca/en/health-canada/services/food-nutrition/healthy-eating/nutrient-data/canadian-nutrient-file-compilation-canadian-food-composition-data-2010-database-structure.html> (accessed September 2016).
44. Moubarac JC, Batal M, Martins AP *et al.* (2014) Processed and ultra-processed food products: consumption trends in Canada from 1938 to 2011. *Can J Diet Pract Res* **75**, 15–21.
45. Food and Agriculture Organization of the United Nations (2015) *Guidelines on the Collection of Information on Food Processing Through Food Consumption Surveys*. Rome: FAO.
46. Kennedy M (2015) One Sweet App. <http://childnutrition.utoronto.ca/news/one-sweet-app> (accessed August 2016).
47. US Department of Agriculture (2006) USDA Database for the Added Sugars Content of Selected Foods, Release 1. <http://www.megaheart.com/pdf/added sugars.pdf> (accessed July 2016).
48. Pan American Health Organization (2015) *Ultra-Processed Food and Drink Products in Latin America*. Washington, DC: PAHO.
49. Monteiro CA, Cannon G, Moubarac JC *et al.* (2015) Dietary guidelines to nourish humanity and the planet in the twenty-first century. A blueprint from Brazil. *Public Health Nutr* **18**, 2311–2322.
50. Moubarac J-C, Parra DC, Cannon G *et al.* (2014) Food classification systems based on food processing: significance and implications for food policies: a systematic literature review and assessment. *Curr Obes Rep* **3**, 256–272.
51. Kendall J (2001) Circles of disadvantage: aboriginal poverty and underdevelopment in Canada. *Am Rev Can Stud* **2001**, 43–49.
52. Dietitians of Canada (2016) *Prevalence, Severity and Impact of Household Food Insecurity: A Serious Public Health Issue. Background Paper Dietitians of Canada*. Toronto, ON: Dietitians of Canada.
53. Manjoo P, Dannenbaum D, Joseph L *et al.* (2015) Utility of current obesity thresholds in signaling diabetes risk in the James Bay Cree of Eeyou Istchee. *BMJ Open Diabetes Res Care* **3**, e000114.
54. Lear SA, Humphries KH, Frohlich JJ *et al.* (2007) Appropriateness of current thresholds for obesity-related measures among Aboriginal people. *CMAJ* **177**, 1499–1505.
55. Canadian Diabetes Association (2015) Diabetes in Canada. <https://www.diabetes.ca/getmedia/513a0f6c-b1c9-4e56-a77c-6a492bf7350f/diabetes-charter-background-under-national-english.pdf.aspx> (accessed September 2016).
56. Downs SM, Arnold A, Marshall D *et al.* (2009) Associations among the food environment, diet quality and weight status in Cree children in Québec. *Public Health Nutr* **12**, 1504–1511.
57. Institute of Medicine, Food and Nutrition Board (2005) *Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. Washington, DC: National Academies Press.
58. Martinez Steele E, Baraldi LG, Louzada ML *et al.* (2016) Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* **6**, e009892.
59. Food and Agriculture Organization of the United Nations (2010) *Fats and Fatty Acids in Human Nutrition. Report of an Expert Consultation*. FAO Food and Nutrition Paper no. 91. Rome: FAO.
60. Gerber L (2014) Education, employment, and income polarization among aboriginal men and women in Canada. *Can Ethn Stud* **46**, 121–144.
61. World Health Organization (2012) *Effect of Increased Potassium Intake on Cardiovascular Disease, Coronary Heart Disease and Stroke*. Geneva: WHO Document Production Services.
62. Johnson C, Raj TS, Trieu K *et al.* (2016) The science of salt: a systematic review of quality clinical salt outcome studies June 2014 to May 2015. *J Clin Hypertens (Greenwich)* **18**, 832–839.
63. World Health Organization (2009) *Global Health Risks. Mortality and Burden of Disease Attributable to Selected Major Risks*. Geneva: WHO.
64. Zhang Z, Cogswell ME, Gillespie C *et al.* (2013) Association between usual sodium and potassium intake and blood pressure and hypertension among US adults: NHANES 2005–2010. *PLoS One* **8**, e75289.
65. Cook NR, Obarzanek E, Cutler JA *et al.* (2009) Joint effects of sodium and potassium intake on subsequent cardiovascular disease: the Trials of Hypertension Prevention follow-up study. *Arch Intern Med* **169**, 32–40.
66. Kwon YI, Apostolidis E, Kim YC *et al.* (2007) Health benefits of traditional corn, beans, and pumpkin: *in vitro* studies for hyperglycemia and hypertension management. *J Med Food* **10**, 266–275.
67. Johnson-Down L, Labonte ME, Martin ID *et al.* (2015) Quality of diet is associated with insulin resistance in the Cree (Eeyouch) indigenous population of northern Quebec. *Nutr Metab Cardiovasc Dis* **25**, 85–92.
68. Johnson-Down L & Egeland GM (2013) How is the nutrition transition affecting the dietary adequacy in Eeyouch (Cree) adults of Northern Quebec, Canada? *Appl Physiol Nutr Metab* **38**, 300–305.
69. Subar AF, Freedman LS, Tooze JA *et al.* (2015) Addressing current criticism regarding the value of self-report dietary data. *J Nutr* **145**, 2639–2645.