

to 64, but not in those aged 65 to 75 years; the latter might be a function of the relatively small number of patients in this age group. The increased prevalence of cataract found in those in the 45- to 64-year-old age group may be the result of increased levels of CuZnSOD, in turn resulting from the location of the associated five exons of SOD1 on chromosome 21. These elevated levels of superoxide dismutase may give rise to increased levels of reactive species, including hydrogen peroxide and hydroxyl radicals, which may increase the risk of cataractogenesis.

P0298

Factor structure of the Barcelona immigration stress scale

A. Qureshi, F. Collazos, H.W. Revollo, S. Valero, M. Ramos. *Servei de Psiquiatria, Hospital Universitari Vall D'Hebron, Barcelona, Spain*

Background: The relationship between immigration and psychopathology is increasingly understood to be indirect and moderated by the stressors related to the migratory process. Empirical study of acculturative stress has been hindered by the lack of an instrument sensitive to the local immigration context. The Barcelona Immigration Stress Scale (BISS) was developed for as a response to this lack. A four factor model is hypothesized as constituting acculturative stress: (1) stress related to the migratory process (homesickness); (2) stress related to the acculturative process (culture shock); (3) stress related to perceived discrimination; and (4) general psychosocial stress. The objective of this study was to evaluate the internal validity of the BISS.

Methods: The BISS was administered to 500 immigrant patients who participated in a study concerned with psychopathology and immigration. Internal consistency was estimated on the basis of the Cronbach alpha and a maximum likelihood factor analysis was carried out with a prior direct oblim rotation.

Results: The BISS has satisfactory internal consistency. Preliminary findings confirmed the four factor model which explained a significant proportion of the common variance.

Discussion: The BISS is internally valid and supports the hypothesized model of acculturative stress. Future studies will establish the external validity of the BISS comparing the scale to related instruments and to determine if acculturative stress is predictive of psychopathology.

P0299

Suicide in urban and rural regions of Belarus, 1990-2005

Y.E. Razvodovsky^{1,2}, A. Stickley^{1,2}. ¹Grodno State Medical University, Grodno, Belarus ²Stockholm Centre On Health of Societies in Transition, Grodno, Belarus

Objective: To examine the occurrence of suicide in urban and rural regions of Belarus in the post-Soviet period. Design: Unlinked cross-sectional study using data drawn from four time points.

Methods: Age and sex-specific suicide data for urban and rural regions of Belarus were obtained from the Ministry of Statistics for the years 1990, 1995, 2000 and 2005. The data were recalculated into seven age categories and then directly standardized. Poisson regression models were used to assess changes in urban-rural suicide rate ratios.

Results: Between 1990 and 2000 the suicide rate rose sharply in Belarus and by 2005 it remained much higher than its initial level. After 1995 there was a divergence between sex-specific rates in urban

and rural areas. By 2005 although suicide rates had fallen from their 2000 level for both sexes in urban and rural locations, the fall was much smaller in rural areas. These changes resulted in a deteriorating rural/urban suicide ratio across the period 1990 to 2005 with suicide rates among nearly every rural male age group remaining extreme after 1995. Although it is probable that a deteriorating social and economic situation has underpinned increasing suicide rates in all regions, there may be factors that are specific to rural locations such as social isolation, alcohol abuse and the poor provision of medical services.

Conclusion: By 2005 Belarus had one of the highest suicide rates in the world. This now requires urgent intervention by the necessary authorities to ameliorate this situation in urban and especially rural locations.

P0300

Inpatient suicidal behaviour during admission: Prevalence and risk factors

T. Neuner, R. Schmid, H. Spiessl. *University of Regensburg, Department of Psychiatry and Psychotherapy, Regensburg, Germany*

Aim: The study aimed at exploring prevalence and risk factors of attempted suicide and suicidal thoughts before hospital admission.

Methods: 40.451 episodes of inpatient care from 1995 to 2004 were included. Last admission of each patient (n=20.543) was extracted to investigate attempted suicide and suicidal thoughts before admission, which were recorded by the German Basic Documentation System (BADO). Univariate analyses and multivariate logistic regression analyses were performed to explore predictors of suicidality before hospital admission.

Results: 3.643 (17.7%) of 20.543 inpatients attempted suicide or had suicidal thoughts before admission. 1.478 (7.2%) attempted suicide before admission. Most had diagnoses F43 (36.5%), F32/33 (19.2%) or F60/61 (8.8%). According to regression analysis risk of attempted suicide increases with life time suicide attempt, diagnosis F43, admission due to doctor's recommendation, diagnosis F32/33, diagnosis F60/61 and period of onset of present episode less than one week. 2.165 (10.5%) had suicidal thoughts before admission without concomitant suicide attempt. Most frequent diagnoses were F43 (23.5%), F32/33 (22.7%) and F10 (11.7%). Risk of suicidal thoughts increases with diagnosis F43, diagnosis F32/33, life time suicide attempt, diagnosis F60/61, supportive psychotherapy before admission, period of onset of present episode less than one week, being younger and being not rehospitalized.

Conclusion: Basic rate of suicidality before admission of psychiatric inpatients is high. Therefore, risk of suicide should be carefully examined at admission but also during inpatient care. Especially, inpatients with adjustment disorder and depression seem to be a high risk groups regarding suicidality before admission and require special attention.

P0301

Sleep disorders in patients with end-stage renal disease. Preliminary report

A. Sieradzki¹, T. Okolow², T. Adamowski¹, A. Kiejna¹, E. Trafidlo², W. Weyde³, P. Biecek⁴, N. Morawska⁵. ¹Department of Psychiatry Medical University of Wroclaw, Wroclaw, Poland ²International Center for Hemodialysis, Wroclaw, Poland ³Department of Nephrology and Transplantation Medicine Medical University of Wroclaw, Wroclaw, Poland ⁴Polish Academy

of Science, Wrocław, Poland⁵ General Public Psychiatric Hospital, Wrocław, Poland

Background and Aims: Sleep disorders are significant problem in hemodialysis patients, contributing to psychosocial disability and decline in quality of life. Aim of this study was to assess the quality of sleep disorders in relation to various clinical and sociodemographic factors in patients undergoing hemodialysis.

Methods: The preliminary study group included 55 persons selected at random from 150 patients treated by hemodialysis. The mean age was $60,31 \pm 12,14$, mean time of dialysis $3,49 \pm 4,53$ years. 20 women and 35 men were interviewed using the Schedule for Clinical Assessment in Neuropsychiatry (SCAN) sleep disorders section and additional sociodemographic interview. In each case psychiatric examination has been performed. Medical comorbidity including restless legs syndrome (RLS), suffering from pain, blood and urine tests were analyzed for potential impact on sleep disturbance.

Results: In studied group sleep disturbances did not show significant correlation with presence of mental disorders, somatic comorbidity other than RLS or most laboratory data (blood or urine tests). The following differences were observed:

1. Positive correlation between overall sleep impairment, night wakefulness and nPCR index related to nutrition and metabolism (p-value 0.047),
2. Greater sleep satisfaction in the group with higher calcium level before hemodialysis (p-value 0.045),
3. Higher nightmares occurrence in the group with lower calcium level before hemodialysis (p-value 0.023).

Conclusions: In the population of patients with end-stage renal disease after adaptation period to hemodialysis more important role in sleep disorders may play factors related to the somatic state and metabolism rather than psychiatric comorbidity.

P0302

Domestic violence against the Iranian pregnant women

A. Soltanifar¹, H.R. Behnam², V. Moghadam Hoseini³.
¹ Psychiatry Department, Mashad University, Avicenna Hospital, Mashad, Iran ² Nursing Department, Mashad University, Mashad, Iran ³ Department, Mashad University, Mashad, Iran

Background: Domestic violence especially from intimate partner, is the most common type of violence against women that has many psychological, social and economic adverse effects. Domestic violence in pregnancy causes serious problems for the infant and mother. Considering the importance of this matter, the present study investigated the frequency and intensity of intimate partner violence against pregnant women.

Method: This study is a cross-sectional research. Two hundred and ninety inpatient mothers in post-delivery sections in educational hospitals were assessed using Iranian version of "Conflict Tactics Scale" (CTS2) after establishing the validity and reliability of this scale.

Results: Results showed that 92.4%, 6.9% and 0.7% of women experienced very mild, mild and moderate intimate partner violence in their pregnancy, respectively. Verbal violence was rated as 40.9% and 0.7%, emotional violence as 81.1% and 1.7%, physical violence as 93.1% and 0.3% and sexual violence as 95.2% and 0.3%, as very mild and severe, respectively. Findings showed that there was a significant difference between the violence against pregnant women and their neonates' low birth weight ($W < 2500$ grams). There was not a significant difference between education of men and violence against their wives ($P = 0.614$).

Discussion: This research shows that many pregnant women experience the very mild forms of violence and mostly as the sexual violence. The rate of neonates' low birth weight has a significant relationship with the intimate partner's violence against Iranian pregnant women. These findings show the importance of this problem and the necessity of considering preventive strategies for that.

P0303

Prevalence of burnout syndrome among medical service providers and hospital employees of Mashhad, Iran

A. Talaei, N. Mokhber. Department of Psychiatry, Ebn-E-Sina Psychiatric Hospital, Mashhad University of Medical Sciences, Mashad, Iran

Background and Aims: This study aimed to evaluate prevalence rate of burnout syndrome in three aspects dimensions of emotional exhaustion, depersonalization, and decrement of personal accomplishment and also determining degrees of vocational satisfaction, work load, social support, and self-respect in medical service providers.

Methods: In this descriptive study we listed all of the hospitals of Mashhad University of Medical Sciences and selected 660 staffs by stratified cluster sampling. They have filled up demographic information forms and self-respect, social support, work load, vocational satisfaction and burnout questionnaires of Maslach Burnout Inventory (MBI). Data has been analyzed by spss and statistical tests (Pearson Chi-Square ...).

Results: Results have shown that 91.2% of study units have high self-respect, 67.1% have high social support, 54.8% have moderate vocational satisfaction, and 67.7% have high work load. In the light of three burnout aspects, 38.6% have low level of dimensions of emotional exhaustion, 58% have low levels of depersonalization, and 62.9% have high levels of decrement of personal accomplishment. Also, findings have depicted considerable direct relation between dimensions of emotional exhaustion and depersonalization feeling and work load, between decrement of personal accomplishment feeling and social support, and between emotional exhaustion feeling and self-respect degree.

Conclusions: Rate of burnout in staffs in dimensions of emotional exhaustion and depersonalization and decrement of personal accomplishment was low but because most of dimensions of burnout were higher among women it is important to increase support of them.

P0304

Psychiatric home care: A descriptive study

M. Thalassinos¹. ¹ Service de Psychiatrie, CASH, Nanterre, France

The present « naturalistic » survey was carried out among a sample of 336 patients who were admitted in the Home Care Psychiatric Unit (HCPU), during 5 years (2001-2006).

Definition: the HCPU provides intensive care with 2 consulting visits per day by the psychiatric staff (including family interviews), for 5 patients at a time, with a maximum duration of 40 days.

Socio demographic data:

58% of the patients were female (n=195).

Mean age was 40,2 years with a range 18-80.

76,5% of the patients had family support at home.

The mean duration of home care was 24,2 days.

Diagnosis: According to ICD 10 criteriae, 46% (n=150) of the patients were assessed a diagnostic of psychotic disorder, 42% (n=141)